# LebanonTurf

# Country Club® Blended Fertilizer

# 17-0-5 METH-EX®



### **GUARANTEED ANALYSIS**

Total Nitrogen (N)
2.5% Water insoluble Nitrogen*
10.1% Urea Nitrogen
4.4% Other Water Soluble Nitrogen*
Soluble Potash (K <sub>2</sub> 0)
Sulfur (S)
1.6% Combined Sulfur (S)
Iron (Fe)
0.01% Water Soluble Iron (Fe)
Manganese (Mn)
0.1% Water Soluble Manganese (Mn)

Derived From: Methylene Ureas, Urea, Sulfate of Potash, Iron

Sucrate, Manganese Sucrate.

\*6.9% Slowly available Nitrogen from Methylene Ureas.

Chlorine (CI) not more than ......

### GIIARANTEEN MINIMIIM ANALYSIS

GUANANI EED WINIMOW ANALISIS	
Total Nitrogen (N)	17%
2.5% Water Insoluble Nitrogen*	
10.1% Urea Nitrogen	
4.4% Other Water Soluble Nitrogen*	
Soluble Potash (K <sub>2</sub> 0)	5%
Sulfur (S)	1.6%
1.6% Combined Sulfur (S)	
Iron (Fe) (actual)	1.0%
0.01% Water Soluble Iron (Fe)	
Manganese (Mn) (actual)	0.7%
0.1% Water Soluble Manganese (Mn)	
Chlorine (CI) (maximum)	2.0%

Derived From: Methylene Ureas, Urea, Sulfate of Potash, Iron Sucrate, Manganese Sucrate.

\*6.9% Slowly available Nitrogen from Methylene Ureas. CANADA

NOTICE: This product contains the secondary nutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.

U.S.A

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

# Application Rates:

To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 5.3 lbs per 1,000 sq. ft. or 230 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.9 lbs per 1,000 sq. ft. or 126 lbs. per acre.

ine ber 1,000 edi iti	от тео прот ро		
SUGGESTED SPREADER SETTINGS			
Spreader 0.9#N	/1,000 sq. ft.	0.5#N/1,000 sq. ft	
Lebanon Turf	41/2	11/2	
Andersons AccuPro	K	F	
Earthway Rotary	15	11	
Gandy	26	21	
Lely	5 II	2½ II	
Lesco (letter/numeric	) I/17	E/15	
PennMulch HVO	L	G	
ProScape SS	L	G	
Spyker .	41/2	11/2	
Vicon	24	16	
Those cottings were	solibrated a	nd field teeted	

These settings were calibrated and field tested.
However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

# WARNING



Hazard Statements

May be harmful if inhaled repeatedly over prolonged periods.

Suspected of causing cancer by prolonged/repeated inhalation.

Precautionary Statements
Avoid breathing dust.

Avoid breathing dust.
Use appropriate personal protective equipment as required to avoid

If exposed or concerned, seek medical advice.

Dispose in accordance with all federal, state and local regulations.

# KEEP OUT OF REACH OF CHILDREN.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations of the commendations of the commendations

tions in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:
To feed at the rate of 0.43 kg Nitrogen (N)
Equivalent to 1.6 kg of product per 100 sq. meters.

Equivalent to 1.6 kg of product per 100 sq. meters

To feed at the rate of 0.22 kg Nitrogen (N)

Equivalent to 0.9 kg of product per 100 sq. meters

Equivalent to	0 U.9 kg of product	per 100 sq. meters		
SI	SUGGESTED SPREADER SETTINGS			
Spreader	0.43 kg #N/100 m <sup>2</sup> .	0.22 kg #N/100 m <sup>2</sup>		
Lebanon Turf	41/2	11/2		
Andersons Acc	uPro K	F		
Earthway Rotar	ry 15	11		
Gandy	26	21		
Lely	5 II	<b>2</b> ½ II		
Lesco (letter/ni	umeric) I/17	E/15		
PennMulch HV	0 L	G		
ProScape SS	L	G		
Cnukar	416	11/6		

Vicon 24 16
These settings were calibrated and field tested.
However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



<sup>F699</sup> **22-35742** 

\_\_\_ j/1

LebanonTurf

For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation

1600 E. Cumberland St • Lebanon, PA 17042 800-233-0628 • (717) 273-1685

50 Lbs. (22.7 kg) Net Weight

# 17-0-5 METHEX®



### **GUARANTEED ANALYSIS**

Total Nitrogen (N)	17%
2.5% Water Insoluble Nitrogen*	
10.1% Urea Nitrogen	
4.4% Other Water Soluble Nitrogen*	
Soluble Potash (K <sub>2</sub> 0)	<b>5</b> %
Sulfur (S)	1.6%
1.6% Combined Sulfur (S)	
Iron (Fe)	1.0%
0.01% Water Soluble Iron (Fe)	
Manganese (Mn)	0.7%
0.1% Water Soluble Manganese (Mn)	
Derived From: Methylene Ureas, Urea, Sulfate of Potash, Iro Sucrate, Manganese Sucrate.	n
*6.9% Slowly available Nitrogen from Methylene Ureas.	

Chlorine (CI) not more than ...... 2.0%

Total Nitrogen (N)2.5% Water Insoluble Nitrogen*	17%
10.1% Urea Nitrogen	
4.4% Other Water Soluble Nitrogen*	
Soluble Potash (K <sub>2</sub> 0)	5%
Sulfur (S)	1.6%
1.6% Combined Sulfur (S)	
Iron (Fe) (actual)	1.0%
0.01% Water Soluble Iron (Fe)	
Manganese (Mn) (actual)	0.7%
0.1% Water Soluble Manganese (Mn)	
Chlorine (CI) (maximum)	2.0%

Derived From: Methylene Ureas, Urea, Sulfate of Potash, Iron Sucrate. Manganese Sucrate.

\*6.9% Slowly available Nitrogen from Methylene Ureas.

CANADA

NOTICE: This product contains the secondary nutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.

U.S.A

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

### **Application Rates:**

To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 5.3 lbs per 1,000 sq. ft. or 230 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.9 lbs per 1,000 sq. ft. or 126 lbs. per acre.

SUGGESTED SPREADER SETTINGS		
Spreader 0.9#N/	1,000 sq. ft.	0.5#N/1,000 sq. ft.
Lebanon Turf	41/2	1½
Andersons AccuPro	K	F
Earthway Rotary	15	11
Gandy	26	21
Lely	5 II	2½ II
Lesco (letter/numeric)	I/17	E/15
PennMulch HVO	L	G
ProScape SS	L	G
Spyker	41/2	1½
Vicon	24	16

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

### WARNING



### **Hazard Statements**

May be harmful if inhaled repeatedly over prolonged periods. Suspected of causing cancer by prolonged/repeated inhalation.

### **Precautionary Statements**

Avoid breathing dust.

Use appropriate personal protective equipment as required to avoid breathing dust.

If exposed or concerned, seek medical advice.

Dispose in accordance with all federal, state and local regulations.

### KEEP OUT OF REACH OF CHILDREN.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

### WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m<sup>2</sup>, applied every 4 to 8 weeks during the local growing season.

### Application Rates:

To feed at the rate of 0.43 kg Nitrogen (N)
Equivalent to 1.6 kg of product per 100 sq. meters.

To feed at the rate of 0.22 kg Nitrogen (N) Equivalent to 0.9 kg of product per 100 sq. meters.

m².

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



F699 I

22-35742



For technical assistance or more information about our products visit www.LebanonTurf.com

# Lebanon Seaboard Corporation

1600 E. Cumberland St • Lebanon, PA 17042 800-233-0628 • (717) 273-1685

50 Lbs. (22.7 kg) Net Weight