

Country Club® Blended Fertilizer





GUARANTEED ANALYSIS Total Nitrogen (N) 3.4% Ammoniacal Nitrogen 5.0% Water Insoluble Nitrogen* 2.4% Urea Nitrogen	19%	GUARANTEED MINIMUM ANALYSIS Total Nitrogen (N) 3.4% Ammoniacal Nitrogen 5.0% Water Insoluble Nitrogen* 2.4% Urea Nitrogen	19%
8.2% Other Water Soluble Nitrogen* Soluble Potash (K ₂ 0)	19%	8.2% Other Water Soluble Nitrogen* Soluble Potash (K ₂ O)	19%
Sulfur (S)	10.2%	Sulfur (S) 10.2% Combined Sulfur (S)	10.2%
10.2% Combined Sulfur (S) Iron (Fe) 0.03% Water Soluble Iron (Fe)	3.0%	Iron (Fe) (actual) 0.03% Water Soluble Iron (Fe)	3.0%

0 00/

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.			
NOTICE: This product contains the micronutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.			
	U.S.A	CANADA	
Chlorine (CI) not more than	2.0%	*13.2% Slowly available Nitrogen from Methylene Ureas.	
*13.2% Slowly available Nitrogen from Methylene Ureas.		Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Sulfate of Potash, Iron Sucrate.	
of Potash, Iron Sucrate.	C	Desired Frame Associate Outfale, Mathematical Users Users Outfale	

WARNING

Application Rates: To feed at the rate of 0.9 lb. Nitro sq. ft., apply this product at 4.7 or 196 lbs. per acre. To feed at th Nitrogen (N) per 1,000 sq. ft., apply lbs per 1,000 sq. ft. or 109 lbs. per	lbs per 1,000 sq. ft. e rate of 0.5 lb. y this product at 2.6	Hazard Statements: Causes skin irritation. May be harmful if inhaled. Causes serious eye irritation. Precautionary Statements for han Wash hands and exposed skin thorou Wear protective gloves/protective clot Take off contaminated clothing and wa IF ON SKIN: Wash with plenty of soar
SUGGESTED SPREADER Spreader 0.9#N/1,000 sq. ft. Lebanon Turf 4½ Andersons AccuPro K Earthway Rotary 15 Gandy 26 Lely 5 II Lesco (letter/numeric) 1/17 PennMulch HVO K ProScape SS K Sovker 4½		IF INHALED: Call » POISON CENTER IF IN EYES: Rinse cautiously with wat lenses, if present and easy to do. Con If skin irritation occurs: Get medical at If eye irritation persists: Get medical a Keep out of reach of children Precautionary Statements for disposa state and local regulations.
Spyker 4½ Vicon 24 These settings were calibrated an However, age and condition of spi operation and evenness of terrain different settings for desired cover	reader, speed of 1 may require slightly erage.	Fertilizer may cause irritation of eyes, n skin or eyes, flush with plenty of water; Use in accordance with recommendat such as, but not limited to a certified c extension publication, or apply accord nutrient management plan.

Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com lazard Statements: Jauses skin irritation. Aay be harmful if inhaled. Jauses serious eye irritation. **Frecautionary Statements for handling:** Vash hands and exposed skin thoroughly after handling. Vash hands and exposed skin thoroughly after handling. Vash protective gloves/protective clothing/eye protection/face protection. ^{Take} off contaminated clothing and wash before reuse. F ON SKIN: Wash with plenty of soap and water. F INHALED: Call » POISON CENTER or doctor/physician if you feel unwell. F IN EYES: Rinse cautiously with water for several minutes. Remove contact enses, if present and easy to do. Continue rinsing. ^I skin irritation occurs: Get medical advice/attention. ^I eye irritation persists: Get medical advice/attention. ^I eye or freach of children Precautionary Statements for disposal - Dispose in accordance with all federal tate and local regulations.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

Do not apply to frozen ground.

WARRANTY


NOTE: To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates: To feed at the rate of 0.43 kg Nitrogen (N) Equivalent to 1.6 kg of product per 100 sq. meters.					
	To feed at the rate of 0.22 kg Nitrogen (N)				
Equivalent to 0.9 kg of product per 100 sq. meters.					
SUGGESTED SPREADER SETTINGS					
Spreader 0.43 k	(g #N/100 m².	0.22 kg #N/100 m².			
Lebanon Turf	41/2	3			
Andersons AccuPro	K	G			
Earthway Rotary	15	13			
Gandy	26	23			
Lely	5 II	3 ½ II			
Lesco (letter/numeric)		F/13			
PennMulch HVO	K	G			
ProScape SS	K	G			
Spyker	4 ½	3			
Vicon	24	21			
These settings were calibrated and field tested.					

However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.







For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation

1600 E. Cumberland St • Lebanon, PA 17042 800-233-0628 • (717) 273-1685

50 Lbs. (22.7 kg) Net Weight





GUARANTEED ANALYSIS Total Nitrogen (N)	19%	GUARANTEED MINIMUM ANALYSIS Total Nitrogen (N)	19%
3.4% Ammoniacal Nitrogen		3.4% Ammoniacal Nitrogen	
5.0% Water Insoluble Nitrogen*		5.0% Water Insoluble Nitrogen*	
2.4% Urea Nitrogen		2.4% Urea Nitrogen	
8.2% Other Water Soluble Nitrogen*		8.2% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ 0)	19%	Soluble Potash (K ₂ O)	19%
Sulfur (S) 10.2% Combined Sulfur (S)	10.2%	Sulfur (S) 10.2% Combined Sulfur (S)	10.2%
Iron (Fe) 0.03% Water Soluble Iron (Fe)	3.0%	Iron (Fe) (actual) 0.03% Water Soluble Iron (Fe)	3.0%
Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Sulfate of Potash, Iron Sucrate.		Chlorine (Cl) (maximum)	2.0%
*13.2% Slowly available Nitrogen from Methylene Ureas.		Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Sulfate of Potash, Iron Sucrate.	
Chlorine (CI) not more than	2.0%	*13.2% Slowly available Nitrogen from Methylene Ureas.	
	U.S.A		CANADA

NOTICE: This product contains the micronutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Application Rates: To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 4.7 lbs per 1,000 sq. ft. or 196 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.6 lbs per 1,000 sq. ft. or 109 lbs. per acre.			
SUGGESTED SPREADER SETTINGS			
Spreader 0.9#N	/1, 000 s	q. ft. 0.5#N/1,000 sq. ft.	
Lebanon Turf	4 ½	3	
Andersons AccuPro	K	G	
Earthway Rotary	15	13	
Gandy	26	23	
Lely	5 II	3 ½ II	
Lesco (letter/numeric)) I/17	F/13	
PennMulch HVO	K	G	
ProScape SS	K	G	
Spyker	4½	3	
Vicon	24	21	
These settings were calibrated and field tested.			
However, age and condition of spreader, speed of			
operation and evenness of terrain may require slightly different settings for desired coverage.			

Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com

WARNING Hazard Statements: Causes skin irritation. May be harmful if inhaled. Causes serious eye irritation. **Precautionary Statements for handling:** Wash hands and exposed skin thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection. Take off contaminated clothing and wash before reuse. IF ON SKIN: Wash with plenty of soap and water. IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If skin irritation occurs: Get medical advice/ attention. If eve irritation persists: Get medical advice/attention. Keep out of reach of children Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

Do not apply to frozen ground.

WARRANTY

NOTE: To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates: To feed at the rate of 0.43 kg Nitrogen (N) Equivalent to 1.6 kg of product per 100 sq. meters.			
To feed at the rate	•		
Equivalent to 0.9 k	g of produc	t per 100 sq. meters.	
SUGGES	SUGGESTED SPREADER SETTINGS		
Spreader 0.43 kg #N/100 m². 0.22 kg #N/100 m².			
Lebanon Turf	4 ½	3	
Andersons AccuPro	K	G	
Earthway Rotary	15	13	
Gandy	26	23	
Lely	5 II	3 ½ II	
Lesco (letter/numeric) /17	F/13	
PennMulch HVO	K	G	
ProScape SS	K	G	
Spyker	4 ½	3	
Vicon	24	21	
These settings were calibrated and field tested.			
However, age and condition of spreader, speed of			
operation and evenness of terrain may require slightly			
different settings for desired coverage.			

