


Lebanon Pro



BRAND

FERTILIZER WITH Prodiamine 0.29%

PREEMERGENCE WEED CONTROL

18-0-4




CONTAINS **SCU**

GUARANTEED ANALYSIS

Total Nitrogen (N) 18.0%
 18.0% Urea Nitrogen*
 Soluble Potash (K₂O) 4.0%
 Sulfur (S) 2.0%
 2.0% Free Sulfur (S)

Derived From: Urea, Sulfur Coated Urea, Muriate of Potash.

Chlorine (Cl) not more than 4.0%
 *4.5% Slowly available Nitrogen from Sulfur Coated Urea.



F699

22-40180

Information regarding the contents and levels of metals in this product is available on the Internet at:
<http://www.regulatory-info-lebsea.com>

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 lb. Nitrogen per 1,000 sq. ft.

SUGGESTED SPREADER SETTINGS		12/06/11	
Spreader	5.2#/1,000 sq. ft.	Spreader	5.2#/1,000 sq. ft.
LebanonTurf	5	Lesco (letter/Numeric)	K/22
Andersons AccuPro	0	PennMulch HVO	N
Earthway Rotary	18	ProScape SS	N
Gandy	30	Spyker	5
Lely	6 1/2 II	Vicon (all models)	30

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

MESA®, PennMulch® and ProScape® are registered trademarks of Lebanon Seaboard Corporation.

For selective preemergence control of grass and broadleaf weeds in:

- established turfgrasses (excluding golf course putting greens) and lawns
- landscape ornamentals (including established perennials & wildflower plantings)

ACTIVE INGREDIENT:

Prodiamine: 0.29%
 OTHER INGREDIENTS: 99.71%
 TOTAL 100.00%

EPA Reg. No. 961-367

EPA Est. No. 961-PA-01:961-IL-01
 Superscript used is first letter of lot number

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.
 Before buying this product, read the Warranty Section elsewhere on this label.
 If terms are not acceptable, return the unopened container to the place of purchase at once.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact (888) 208-1368 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

If material is spilled: Contain and sweep up material of spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.



For technical assistance or more information about our products visit www.LebanonTurf.com

Manufactured by:

Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042

Made in the U.S.A.

www.LebanonTurf.com

Net Weight 50 LBS. (22.7 kg)

FERTILIZER WITH Prodiamine 0.29%

PREEMERGENCE WEED CONTROL

18-0-4



CONTAINS SCU

GUARANTEED ANALYSIS

Total Nitrogen (N) 18.0%
 18.0% Urea Nitrogen*
 Soluble Potash (K₂O) 4.0%
 Sulfur (S) 2.0%
 2.0% Free Sulfur (S)

Derived From: Urea, Sulfur Coated Urea, Muriate of Potash.

Chlorine (Cl) not more than 4.0%
 *4.5% Slowly available Nitrogen from Sulfur Coated Urea.



F699

22-40180

Information regarding the contents and levels of metals in this product is available on the Internet at:
<http://www.regulatory-info-lebsea.com>

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 lb. Nitrogen per 1,000 sq. ft.

SPREADER		SUGGESTED SPREADER SETTINGS		12/06/11
Spreader	5.2#/1,000 sq. ft.	Spreader	5.2#/1,000 sq. ft.	
LebanonTurf	5	Lesco (letter/Numeric)	K/22	
Andersons AccuPro	0	PennMulch HVO	N	
Earthway Rotary	18	ProScape SS	N	
Gandy	30	Spyker	5	
Lely	6½ II	Vicon (all models)	30	

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

MESA®, PennMulch® and ProScape® are registered trademarks of Lebanon Seaboard Corporation.

For selective preemergence control of grass and broadleaf weeds in:

- established turfgrasses (excluding golf course putting greens) and lawns
- landscape ornamentals (including established perennials & wildflower plantings)

ACTIVE INGREDIENT:

Prodiamine: 0.29%
 OTHER INGREDIENTS: 99.71%
 TOTAL..... 100.00%

EPA Reg. No. 961-367

(L) (D)
 EPA Est. No. 961-PA-01:961-IL-01
 Superscript used is first letter of lot number

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

Before buying this product, read the Warranty Section elsewhere on this label.

If terms are not acceptable, return the unopened container to the place of purchase at once.

KEEP OUT OF REACH OF CHILDREN CAUTION

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact (888) 208-1368 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

If material is spilled: Contain and sweep up material of spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.



For technical assistance or more information about our products visit www.LebanonTurf.com

Manufactured by:

Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042

Made in the U.S.A.

www.LebanonTurf.com

Net Weight 50 LBS. (22.7 kg)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turfgrasses (excluding golf course putting greens) and lawns, landscape ornamentals, established perennials and wildflower plantings.

This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climactic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Do not graze or feed livestock forage cut from areas treated with this product.

Do not apply this product through any type of irrigation system.

Do not apply aerially.

Do not apply to golf course putting greens.

APPLICATION DIRECTIONS

Apply uniformly with suitable, calibrated application equipment.

ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turfgrasses and lawns. The maximum amount of this product that may be applied per year is given for each turfgrass species in the **Maximum Annual Rates** section of this label. Most effective weed control in turfgrasses will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

CRABGRASS SEED GERMINATION DATES



Approximate Date



USE PRECAUTIONS

The following precautions apply to the use of this product in turfgrasses and lawns:

Application of this product may thin emerged annual bluegrass and newly overseeded grasses.

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

Do not apply to newly set sod until the following year.

Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.

Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.

Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES

This product is suggested for use on the turf-grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

MAXIMUM APPLICATION RATE/CALENDAR YEAR of Lebanon Pro Brand Fertilizer with Prodiamine 0.29% Preemergence Weed Control by Turfgrass Species¹

Turf Species:	Lbs. Product/Acre	Lbs. Product/1,000 Sq. Ft.	Lbs. AI/Acre
Bermudagrass ² , Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turf-type), Zoysia	518	11.9	1.5
Buffalograss, Kentucky Bluegrass, Perennial Ryegrass	345	7.9	1.0
Creeping Red Fescue	260	6.0	0.75
Creeping Bentgrass	225	5.2	0.65

¹) These are the maximum rates per calendar year by species limitations.

²) May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 179 lbs./Acre[4.1 Lbs./1000 Sq.Ft.]. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

• Do not apply more than 518 lbs. [11.9 Lbs./1000 Sq. Ft.] of this product per calendar year.

• Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the **Maximum Annual Rates Table**.

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:

Barnyardgrass	Crabgrass (large, smooth)	Johnsongrass (from seed)	Browntop)	Speedwell, Persian
Bluegrass, Annual (Poa annua)	Crowfootgrass	Junglerice	Pigweed	Sprangletop
Carpetweed	Cupgrass, Woolly	Knotweed	Purslane, Common	Spurge, Prostrate
Chickweed, Common	Foxtails, Annual	Kochia	Pusley, Florida	Witchgrass
Chickweed, Mouseear (from seed)	Goosegrass ¹	Lambsquarter, Common	Rescuegrass ³	Woodsorrel, Yellow (from seed)
	Henbit	Lovegrass	Shepherd's Purse ²	
	Itchgrass	Panicum (Texas, Fall,	Signalgrass, Broadleaf	

¹ In many areas a single application of 225 to 518 lbs./Acre [5.2 to 11.9 Lbs./1000 Sq. Ft.] of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 225 to 345 lbs./Acre 5.2 to 7.9 Lbs./1000 Sq. Ft.] followed after 60 to 90 days by a second application at doses that would not exceed those given in the **Maximum Annual Rate Table**. Do not exceed the maximum rate for turfgrass species listed in the **Maximum Annual Rate Table** above.

² Applications for this weed should be made in late summer, fall or winter prior to germination.

³ Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates listed for each turf species. All applications must be made prior to germination of the weed seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

LEBANON PRO BRAND FERTILIZER WITH PRODIAMINE 0.29%PREEMERGENCE WEED CONTROL

Product	Rate (lb./A)	Months Before Overseeding		
		North	Transition	South
172	.50	4	4	4
225	.65	5	4	4
260	.75	6	5	5
275	.80	---	6	6
345	1.00	---	7	7
393	1.14	---	---	9
448	1.30	---	---	10
518	1.50	---	---	12

LANDSCAPE ORNAMENTALS

(including Established Perennials and Wildflower Plantings)

This product may be applied for residual preemergence weed control in ornamentals.

USE RATES

Apply this product at 225 to 518 lbs./Acre [5.2 to 11.9 Lbs./1000 Sq. Ft.] in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 518 lbs./Acre [11.9 Lbs./1000 Sq. Ft.].

APPLICATION TIMING AND INFORMATION

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

ESTABLISHED LANDSCAPE ORNAMENTALS

This product may be applied at any time to established ornamentals as a broadcast, over the top or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

NEWLY TRANSPLANTED LANDSCAPE ORNAMENTALS

This product may be applied to newly transplanted ornamentals as a broadcast, over the top or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

TOLERANT ORNAMENTAL SPECIES

TOLERANT ORNAMENTAL SPECIES

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

Scientific Name	Common Name	Scientific Name	Common Name	Scientific Name	Common Name	Scientific Name	Common Name
<i>Abies spp.</i>	Fir species** (Balsam, Fraser, Noble, etc.)	<i>C. Dammeri</i>	Bearberry Cotoneaster	<i>J. conferta</i>	Shore Juniper; Blue Pacific	<i>P. tobira</i>	Japanese Pittosporum
<i>Abelia grandiflora</i>	Abelia: Sherwood	<i>C. microphyllus</i>	Rockspray Cotoneaster	<i>J. davurica</i>	Parsoni	<i>Podocarpus macrophyllus</i>	Japanese Yew
<i>Acer palmatum</i>	Japanese Maple	<i>Cretaegus spp.</i>	Hawthorne	<i>J. horizontalis</i>	Creeping Juniper	<i>Prunus Laurocerasus</i>	English Laurel
<i>A. platanoides</i>	Norway Maple	<i>Crococsmia sp.</i>	Lucifer	<i>Juglans sp.</i>	Walnut*	<i>Prunus sp.</i>	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Achillea sp.</i>	Yarrow: King Edward	<i>Cupressus sempervirens</i>	Italian Cypress	<i>Justicia brandegeana</i>	Shrimp Plant	<i>Pseudotsuga Menziesii</i>	Douglas Fir**
<i>Actinidia chinensis</i>	Kiwi*	<i>Delosperma alba</i>	White Trailing Ice Plant	<i>Lagerstromia indica</i>	Crape Myrtle	<i>Pyracantha coccinea</i>	Firethorn (Scarlet)
<i>Agapanthus africanus</i>	Lily-of-the-Nile (African Lily)	<i>Delosperma sp.</i>	Cooperi Pink	<i>Lagerstomia indica X Fauriei</i>	Crape Myrtle; Tuscarora	<i>P. Fortuneana</i>	Firethorn
<i>Agapanthus orientalis</i>		<i>Delphinium sp.</i>	Larkspur: Blue Elf	<i>Lantana montevidensis</i>	Weeping Lantana	<i>P. Koidzumii</i>	Firethorn
<i>Akebia quinata</i>	Five-Leaf or Chocolate Vine	<i>Dianthus deltooides</i>	Dianthus; Maiden Pinks "Zing"	<i>Lavender sp.</i>	Lavender; Munstead	<i>Pyrus sp.</i>	Bradford Pear sp.
<i>Allium cernuum</i>	Lady's Leek, Nodding Onion	<i>D. gratianopolitanus</i>	Cheddar Pink	<i>Leontopodium alpinum</i>	Edelweiss	<i>Quercus Shumardii</i>	Oak, Shumard's Red
<i>Anemone hybrida</i>	Japanese Anemone	<i>Dodonea viscosa</i>	Lady's Leek, Nodding Onion	<i>Ligustrum sinense</i>	Hop Bush	<i>Quercus spp.</i>	Oak species
<i>Aquilegia sp.</i>	Aquilegia: Red and Gold	<i>Echinacea pupurea</i>	Coneflower, Purple; Magnus	<i>Ligustrum amurense</i>	Amur Privet	<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita	<i>Elaeagnus pungens</i>	Silverberry	<i>L. japonicum</i>	Japanese Privet	<i>Raphiolepis umbellata</i>	Yedda Hawthorn
<i>Arctotheca calendula</i>	Cape Weed	<i>Euonymus Fortunei</i>	Wintercreeper	<i>L. lucidum</i>	Glossy Privet (Waxleaf)	<i>Rhododendron</i>	'Coral Bells'
<i>Artemisia sp.</i>	Wormwood: Silver Mound, Castle	<i>E. japonica</i>	Japanese Spindle Tree (Evergreen)	<i>Lillium sp.</i>	Lily: Jazz	<i>(including Azalea)</i>	'Delaware Valley White'
<i>Aster sp.</i>	Aster: Bonny Blue, Purple Dome	<i>E. kiautschovica</i>	Euonymus	<i>Liriope Muscari</i>	Big Blue Lily; Turf; Silver Mound;		'Flame Creeper'
<i>Aster X Frikartii</i>		<i>Fatsia japonica</i>	Spreading Euonymus		Evergreen Giant		'Formosa'
<i>Athrium Filiz-femina</i>	Lady Fern; Fern Lady	<i>Forsythia intermedia</i>	Japanese Aralia	<i>Liriope Muscari var. variegata</i>	Japanese Aralia		'Girard Crimson'
<i>Aucuba japonica</i>	Japanese Aucuba	<i>Forsythia suspensa</i>	Border Forsythia	<i>Liriope spicata</i>	Liriope, Creeping		'George L. Tabor'
<i>Begonia sp.</i>	Fibrous Begonia: Hardy Grandis	<i>Forsythia suspensa</i>	Weeping Forsythia	<i>Lobelia Cardinalis</i>	Cardinal Flower, Indian Pink		'Hino-crimson'
<i>Berberis gladwynensis</i>	Barberry	<i>F. viridissima</i>	Greenstem Forsythia	<i>Lonicera japonica</i>	Japanese Honeysuckle		'PJM'
<i>B. Julianae</i>	Wintergreen Barberry	<i>Gaillardia sp.</i>	Gaillardia, Blanket Flower: "Goblin"	<i>L. tatarica</i>	Tatarian Honeysuckle		'Roseum Elegans'
<i>B. mentorensis</i>	Mentor Barberry	<i>Gardenia jasminoides</i>	Gardenia, Cape-Jasmine, Radicans	<i>Loropetalum chinense</i>	Burgundy		'Wakeiebisu'
<i>B. Thunbergii</i>	Japanese Barberry	<i>Gaura sp.</i>		<i>Lythrum sp.</i>	Loosestrife; Modern Pink		'White Gumbo'
<i>B. verruculosa</i>	Warty Barberry	<i>Gentiana dahurica</i>	Gentian	<i>Magnolia spp.</i>	Magnolia species**	<i>Rosa Banksiae</i>	Lady Bank's Rose
<i>Bergenia cordifolia</i>		<i>Geranium cinereum</i>	Cranesbill	<i>Maleophora luteola</i>	Ice Plant	<i>Rosmarinus officinalis</i>	Rosemary*
<i>Boltonia asteroides</i>	Snowbank	<i>Gladiolus spp.</i>	Gladiolus species**	<i>Malus sp.</i>	Crabapple*	<i>Rudbeckia sp.</i>	Black-Eyed Susan; Goldstrum
<i>Buddleia davidii</i>	Butterfly-Bush (Dwarf Blue); Royal Red	<i>Gypsophila repens</i>	Baby's Breath	<i>Miscanthus sinensis</i>	Yaku Jima, Silberfeder	<i>Santolina virens</i>	Santolina
<i>Buxus microphylla</i>	Japanese Boxwood	<i>Hedera helix</i>	English Ivy	<i>Nandina domestica</i>	Heavenly Bamboo	<i>Saxifraga sp.</i>	Saxifrage; Purple Dome
<i>Callistemon viminalis</i>	Weeping Bottlebrush	<i>Hellanthemum sp.</i>	Sunrose	<i>Narcissus spp.</i>	Narcissus species**	<i>Scabiosa sp.</i>	Pincushion Flower
<i>Calluna vulgaris</i>	Scotch Heather	<i>Hemerocallis sp.</i>	Daylily: Aztec Gold, Stella De Oro, Tender Love	<i>Nerium oleander</i>	Oleander	<i>Sedum album</i>	Stonecrop
<i>Campanula carpatica</i>	Tussock Bellflower, (White Clips)	<i>Hibiscus</i>	Rose of Sharon**	<i>Olea europaea</i>	Olive*	<i>Sedum cautilcola</i>	Stonecrop; Lidakense
<i>Campis X Tagliabuana</i>	Trumpet Creeper, Trumpet Flower, Madame Galen	<i>Hibiscus sp.</i>	Mallow: Disco Belle White	<i>Ophiopogon japonicus</i>	Mondo Grass	<i>Sedum dasyphyllum</i>	Stonecrop
<i>Carpobrotus edulis</i>	Hottentot Fig (Ice Plant)	<i>Hibiscus Rosa-sinensis</i>	Chinese Hibiscus	<i>Osmanthus heterophyllus</i>	Osmanthus (False Holly); Gulf Tide	<i>Sedum spurium</i>	Stonecrop; Dragon's Blood
<i>Cassia artemisoides</i>	Feathery Cassia	<i>Hosta plantaginea</i>	Hosta, Plantain Lily (Fragrant)	<i>Osteospermum fruticosum</i>	Trailing African Daisy	<i>Spiraea Bumalda</i>	Spirea: Anthony Waterer
<i>Ceanothus rigidus</i>	Wild Lilac	<i>H. Sieboldiana</i>	Hosta, "Searsucker"	<i>Oxydendrum arboreum</i>	Sourwood	<i>Syzygium paniculatum</i>	Australian Brushcherry
<i>Ceratostigma plumbagonoides</i>	False Cypress	<i>Houtunymia cordata var. variegata</i>		<i>Paeonia suffruticosa</i>	Tree Peony	<i>Taxus cuspidata</i>	Japanese Yew
<i>Chamaecyparis pisifera</i>		<i>Hydrangea macrophylla</i>	Bigleaf Hydrangea	<i>Pennisetum setaceum</i>	Fountain Grass (Dwarf)	<i>T. media</i>	Yew
<i>Chrysanthemum nipponicum</i>		<i>Ilex Bernyi</i>	Holly	<i>Perovskia atriplicifolia</i>		<i>Teucrium sp.</i>	Germander
<i>Cleyera japonica</i>	Cleyera	<i>Ilex cornuta</i>	Chinese Holly	<i>Persea americana</i>	Avocado*	<i>Thalictrum dipterocarpum</i>	Meadow Rue
<i>Citrus spp.</i>	Citrus species*	<i>I. crenata</i>	Japanese Holly; Helli	<i>Photinia Fraseri</i>	Fraser's Photinia (Redtip)	<i>Thuja occidentalis</i>	American Arborvitae
<i>Coreopsis sp.</i>	Coreopsis (Calliopsis): Early Sunrise, Moonbeam	<i>I. opaca</i>	American Holly	<i>Physotegia Virginiana</i>	Dragonhead, False; Vivid	<i>Trachelospermum asiaticum</i>	Star Jasmine
<i>Cornus florida</i>	Flowering Dogwood	<i>I. vomitoria</i>	Yaupon Holly; Schillings	<i>Picea spp.</i>	Spruce species** (Colorado Blue, Norway, etc.)	<i>Tsuga canadensis</i>	Canada Hemlock
<i>C. stolonifera</i>	American Dogwood	<i>Inula ensifolia</i>		<i>Pieris japonica</i>	Lily-of-the-Valley Bush	<i>Tulipa spp.</i>	Tulip species
<i>Cortaderia selloana</i>	Pampas Grass	<i>Iris ensata</i>	Sword-Leaved Iris; Jodlesong	<i>Pinus brutia</i>	Calabrian Pine	<i>Veronica sp.</i>	Veronica, Speedwell; Sunny Border
<i>Cotoneaster apiculatus</i>	Cranberry Cotoneaster	<i>Iris siberica</i>	Siberian Iris; Cabernet	<i>P. canariensis</i>	Canary Island Pine	<i>Viburnum japonicum</i>	Japanese Viburnum
<i>C. buxifolius</i>	Cotoneaster	<i>Iris spp.</i>	Iris species**	<i>P. elliotii</i>	Slash Pine	<i>V. odoratissimum</i>	Sweet Viburnum
		<i>Jasminium nudiflorum</i>	Winter Jasmine	<i>P. halepensis</i>	Aleppo Pine	<i>V. plicatum</i>	Japanese Snowball
		<i>Juniperus chinensis</i>	Chinese Juniper; Nick's Compact, Parsonii	<i>P. nigra</i>	Austrian Pine	<i>V. rigidum</i>	Canary Island Viburnum
				<i>P. palustris</i>	Longleaf Pine	<i>V. suspensum</i>	Arrowwood Viburnum
				<i>P. radiata</i>	Monterey Pine	<i>V. tinus</i>	Laurustinus
				<i>P. Thunbergiana</i>	Japanese Black Pine	<i>V. trilobium</i>	Cranberry Bush
				<i>P. strobus</i>	Eastern White Pine	<i>V. wrightii</i>	Leatherleaf Viburnum
				<i>P. syvestris</i>	Scotch Pine	<i>Vinca major</i>	Greater Periwinkle, Vinca
				<i>P. taeda</i>	Loblolly Pine	<i>Vinca minor</i>	Common Periwinkle, Vinca
				<i>P. virginiana</i>	Virginia Pine	<i>Vitis sp.</i>	Grape*
				<i>Pistachio sp.</i>	Pistachio*	<i>Weigela florida</i>	Old Fashioned Weigela
				<i>Pittosporum rhombifolium</i>	Queensland Pittosporum	<i>Yucca aloifolia</i>	Spanish Bayonet, Yucca
						<i>Y. filamentosa</i>	Adam's Needle, Yucca

* Do not use on food producing trees, vines, or plants.
** Not for use on container grown plants.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store product in its original container in a cool, dry, locked place out of the reach of children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

WARRANTY IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Lebanon Seaboard Corporation. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, LEBANON SEABOARD CORPORATION MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Lebanon Seaboard Corporation is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Lebanon Seaboard Corporation DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT LEBANON SEABOARD CORPORATION'S ELECTION, THE REPLACEMENT OF PRODUCT.