


# Lebanon Pro



## BRAND

Fertilizer with

TEAM<sup>®</sup> PRO 0.86%

PREEMERGENCE CRABGRASS CONTROL

22-0-6

CONTAINS SCU

GUARANTEED ANALYSIS

Total Nitrogen

22.0% Urea Nitrogen\*

Soluble Potash (K<sub>2</sub>O)

Sulfur (S)

1.2% Free Sulfur (S)

Derived from: Sulfur Coated Urea, Urea, Muriate of Potash.

Chlorine (Cl) not more than

\*Contains 8.8% Slowly Available Nitrogen from Sulfur Coated Urea

22%

6%

1.2%

6.0%

SIZE GUIDE NUMBER

SGN 210

0 88685 41436 7

22-41436

Information regarding the contents and levels of metals in this product is available on the Internet at:  
(CA) <https://apps4.cdfa.ca.gov/FertilizerProducts/>  
(OR) [http://oda.state.or.us/dbs/heavy\\_metal/search.lasso](http://oda.state.or.us/dbs/heavy_metal/search.lasso)  
(WA) <http://agr.wa.gov/pestfert/fertilizers/productdatabase.aspx>

SUGGESTED SPREADER SETTINGS			
Spreader	4# / 1,000 Sq. Ft.	Spreader	4# / 1,000 Sq. Ft.
LebanonTurf	4½	Lesco (letter/Numeric)	J / 21
Andersons AccuPro	M	PennHuich HVO	M
Earthway Rotary	16	ProScope SS	M
Gandy	28	Spyker	4½
Lely	6 II	Vicon (all models)	26

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established turfgrass and landscape ornamentals

#### ACTIVE INGREDIENTS:

benefin: *N*-butyl-*N*-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-*p*-toluidine . . . 0.43%  
trifluralin:  $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-*N,N*-dipropyl-*p*-toluidine . . . 0.43%  
OTHER INGREDIENTS . . . . . 99.14%  
TOTAL . . . . . 100.00%

EPA Reg. No. 62719-289-961  
Team<sup>®</sup> is a Trademark of Dow AgroSciences LLC

(L) (D)  
EPA Est. 961-PA-01:961-IL-01  
Superscript used is first letter of lot number.

KEEP OUT OF REACH OF CHILDREN  
WARNING

### Precautionary Statements Hazards to Humans and Domestic Animals

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In some Individuals

Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Refer to back panel for additional precautionary information including Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.



Distributed by:  
**Lebanon Seaboard Corporation**  
1600 E. Cumberland St. • Lebanon, PA 17042  
Made in the U.S.A.  
[www.LebanonTurf.com](http://www.LebanonTurf.com)

NET WT. 50 LBS. (22.7 kg)

# Fertilizer with TEAM<sup>®</sup> PRO 0.86%

## PREEMERGENCE CRABGRASS CONTROL

# 22-0-6



## CONTAINS SCU

### GUARANTEED ANALYSIS

Total Nitrogen ..... 22%

22.0% Urea Nitrogen\*

Soluble Potash (K<sub>2</sub>O) ..... 6%

Sulfur (S) ..... 1.2%

1.2% Free Sulfur (S)

Derived from: Sulfur Coated Urea, Urea, Muriate of Potash.

Chlorine (Cl) not more than ..... 6.0%

\*Contains 8.8% Slowly Available Nitrogen from Sulfur Coated Urea

F699



## 22-41436

Information regarding the contents and levels of metals in this product is available on the Internet at:

(CA) <https://apps4.cdfa.ca.gov/FertilizerProducts/>

(OR) [http://oda.state.or.us/dbs/heavy\\_metal/search.lasso](http://oda.state.or.us/dbs/heavy_metal/search.lasso)

(WA) <http://agr.wa.gov/pestfert/fertilizers/productdatabase.aspx>

### SUGGESTED SPREADER SETTINGS

Spreader	4# / 1,000 Sq. Ft.	Spreader	4# / 1,000 Sq. Ft.
LebanonTurf	4½	Lesco (letter/Numeric)	J / 21
Andersons AccuPro	M	PennMulch HVO	M
Earthway Rotary	16	ProScape SS	M
Gandy	28	Spyker	4½
Lely	6 II	Vicon (all models)	26

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established turfgrass and landscape ornamentals

**ACTIVE INGREDIENTS:**

benefin: *N*-butyl-*N*-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-*p*-toluidine . . 0.43%  
trifluralin:  $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-*N,N*-dipropyl-*p*-toluidine . . . 0.43%

**OTHER INGREDIENTS . . . . . 99.14%**  
**TOTAL . . . . . 100.00%**

EPA Reg. No. 62719-289-961  
Team® is a Trademark of Dow AgroSciences LLC

(L) (D)  
EPA Est. 961-PA-01:961-IL-01  
Superscript used is first letter of lot number.

**KEEP OUT OF REACH OF CHILDREN**  
**WARNING**

**Precautionary Statements**  
**Hazards to Humans and Domestic Animals**

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In some Individuals

Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Refer to back panel for additional precautionary information including Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.



**Distributed by:**

**Lebanon Seaboard Corporation**

**1600 E. Cumberland St. • Lebanon, PA 17042**

**Made in the U.S.A.**

**[www.LebanonTurf.com](http://www.LebanonTurf.com)**

**NET WT. 50 LBS. (22.7 kg)**





# Fertilizer with TEAM<sup>®</sup> PRO 0.86% PREEMERGENCE CRABGRASS CONTROL

## Personal Protective Equipment (PPE)

All loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material
- Goggles, faceshield or safety glasses

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.  
**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

## Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

## Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.  
Read all directions carefully before applying this product.

**Application Restrictions:** Do not apply this product in a way that will contact workers or other persons or pets, either directly or through drift. Only protected handlers may be in the area during application.

**Entry Restrictions:** Do not enter or allow people or pets to enter the treated area until dusts have settled. If soil incorporation is required after application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

**Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.**

## Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.  
**Pesticide Storage:** Store in original container. In case of leak or spill, contain material and dispose as waste.  
**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.  
**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

## General Information

This product is a granular fertilizer containing Team®, a preemergence herbicide for application to established turfgrass and landscape ornamentals (trees, shrubs, groundcovers, and flowers) for control of annual grasses and broadleaf weeds.

## General Use Precautions and Restrictions

- Do not apply this product to golf course putting greens.
- Do not apply to dichondra.
- Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.

**Reseeding:** Do not reseed treated areas for at least 8 weeks after a single application of 174 lb of product per acre, or 12 to 16 weeks after a single application of 349 lb of product per acre unless injury can be tolerated. For split applications, do not reseed treated areas for 8 to 10 weeks after the last 174 lb of product per acre rate is applied, or 10 to 12 weeks after the last 233 lb of product per acre rate is applied unless injury can be tolerated.

## This product is recommended for use on the following turfgrass species:

### Established Cool Season Turfgrass

bentgrass fairways  
fescues  
Kentucky bluegrass  
perennial ryegrass

### Established Warm Season Turfgrass

bahiagrass  
Bermudagrass  
centipedegrass  
fescue, tall  
St. Augustinegrass  
zoysiagrass

**This product may be used for annual weed control in landscape settings on the following established ornamental plant species:**

Ornamental Trees

Common Name
almond
apple, crabapple
apricot
ash, white
bald cypress
birch, European white
blackgum
cherry
chestnut, Chinese
cottonwood
dogwood, flowering
dogwood, kousa
douglas fir
fir, balsam
hemlock, Canada
honeylocust
larch, Japanese
locust, black
maple, Norway
maple, red
maple, silver
maple, sugar
oak, pin
oak, scarlet
peach
pine, Austrian
pine, eastern white
pine, Japanese black
pine, loblolly
pine, red
pine, scotch
planetree, London
plum
redbud, eastern
spruce, Colorado
spruce, Norway
spruce, white
sweetgum
sycamore
tuliptree
walnut, black

Scientific Name
<i>Prunus dulcis</i>
<i>Malus</i> spp.
<i>Prunus armeniaca</i>
<i>Fraxinus americana</i>
<i>Taxodium distichum</i>
<i>Betula pendula</i>
<i>Nyssa sylvatica</i>
<i>Prunus</i> spp.
<i>Castanea mollissima</i>
<i>Populus deltoides</i>
<i>Cornus florida</i>
<i>Cornus kousa</i>
<i>Pseudotsuga menziesii</i>
<i>Abies balsamea</i>
<i>Tsuga canadensis</i>
<i>Gleditsia triacanthos</i>
<i>Larix kaempferi</i>
<i>Robinia pseudoacacia</i>
<i>Acer platanoides</i>
<i>Acer rubrum</i>
<i>Acer saccharinum</i>
<i>Acer saccharum</i>
<i>Quercus palustris</i>
<i>Quercus coccinea</i>
<i>Prunus persica</i>
<i>Pinus nigra</i>
<i>Pinus strobus</i>
<i>Pinus thunbergiana</i>
<i>Pinus taeda</i>
<i>Pinus resinosa</i>
<i>Pinus sylvestris</i>
<i>Platanus acerifolia</i>
<i>Prunus</i> spp.
<i>Cercis canadensis</i>
<i>Picea pungens</i>
<i>Picea abies</i>
<i>Picea glauca</i>
<i>Liquidambar styraciflua</i>
<i>Platanus occidentalis</i>
<i>Liriodendron tulipifera</i>
<i>Juglans nigra</i>

Ornamental Shrubs

Common Name
andromeda, Japanese
arborvitae, American
azalea
barberry, Japanese
barberry, mento
boxwood, common
boxwood, littleleaf
camellia, Japanese
camellia, sasanqua
cherrylaurel, American
cinquefoil
cleyera, Japanese
cotoneaster, cranberry
cotoneaster, zabel
deutzia
elaeagnus, silverberry
euonymus, spreading
euonymus, winged
euonymus, wintercreeper
firethorn
forsythia
guava, pineapple
holly
honeysuckle
India hawthorn
juniper
laurel, mountain
lilac, common
mockorange
pittosporum, Japanese
privet
redcedar, eastern
rhododendron
spiraea, vanhoutte
viburnum
weigela
willow
yew, anglojap
yew, Japanese
yewpine

Scientific Name
<i>Pieris japonica</i>
<i>Thuja occidentalis</i>
<i>Rhododendron</i> spp.
<i>Berberis thunbergii</i>
<i>Berberis mentorensis</i>
<i>Buxus sempervirens</i>
<i>Buxus microphylla</i>
<i>Camellia japonica</i>
<i>Camellia sasanqua</i>
<i>Prunus caroliniana</i>
<i>Potentilla</i> spp.
<i>Cleyera japonica</i>
<i>Cotoneaster apiculata</i>
<i>Cotoneaster zabelii</i>
<i>Deutzia</i> spp.
<i>Elaeagnus pungens</i>
<i>Euonymus kiautschovica</i>
<i>Euonymus alatus</i>
<i>Euonymus fortunei</i>
<i>Pyracantha</i> spp.
<i>Forsythia</i> spp.
<i>Feijoa sellowiana</i>
<i>Ilex</i> spp.
<i>Lonicera</i> spp.
<i>Raphiolepis indica</i>
<i>Juniperus</i> spp.
<i>Kalmia latifolia</i>
<i>Syringa vulgaris</i>
<i>Philadelphus</i> spp.
<i>Pittosporum tobira</i>
<i>Ligustrum</i> spp.
<i>Juniperus virginiana</i>
<i>Rhododendron</i> spp.
<i>Spiraea vanhouttei</i>
<i>Viburnum</i> spp.
<i>Weigela</i> spp.
<i>Salix</i> spp.
<i>Taxus media</i>
<i>Taxus cuspidata</i>
<i>Podocarpus macrophyllus</i>

Ornamental Groundcover Plantings

Common Name
aaronsbeard
bellflower, adriatic
bellflower, poscharsky
ceanothus
coreopsis
cotoneaster
coyote brush
crown vetch
daisy trailing African
fern, asparagus
gazania
germander
ice plant, largeleaf
ivy, Algerian
ivy, English
lily-of-the-nile
lilyturf, bigblue
marigold
myoporum
plumbago, dwarf
rockrose
rosemary
rupturewort
snow-in-summer
speedwell
St. Johnswort
stonecrop (sedum)
strawberry, beach
thrift
verbena
wirevine, creeping
yarrow, wooly

Scientific Name
<i>Hypericum calycinum</i>
<i>Campanula elatines</i>
<i>Campanula poscharskyana</i>
<i>Ceanothus</i> spp.
<i>Coreopsis</i> spp.
<i>Cotoneaster</i> spp.
<i>Baccharis pilularis</i>
<i>Coronilla vana</i>
<i>Osteospermum fruticosum</i>
<i>Asparagus densiflorus</i>
<i>Gazania</i> spp.
<i>Teucrium chamaedrys</i>
<i>Carpobrotus edulis</i>
<i>Hedera canariensis</i>
<i>Hedera helix</i>
<i>Agapanthus</i> spp.
<i>Liriope muscari</i>
<i>Tagetes</i> spp.
<i>Myoporum laetum</i>
<i>Ceratostigma umbaginoides</i>
<i>Cistus</i> spp.
<i>Rosmarinus officinalis</i>
<i>Herniaria glabra</i>
<i>Cerastium tomentosum</i>
<i>Veronica</i> spp.
<i>Hypericum coris</i>
<i>Sedum</i> spp.
<i>Fragaria chiloensis</i>
<i>Armeria maritima</i>
<i>Verbena</i> spp.
<i>Muehlenbeckia axillaris</i>
<i>Achillea tomentosa</i>

Roses and Other Established Flowers

Common Name	Common Name
African daisy	marigold
aster (perennial)	marigold, cape
balsam	morningglory
blackeyed susan	nasturtium
calendula	nicotiana
carnation	petunia
centaurea, velvet	phlox
chrysanthemum	pincushion flower
coreopsis	poppy, California
cornflower	portulaca
cosmos	rose
dahlia	salvia
dianthus	shasta daisy
dusty miller	snapdragon
floss flower	snow-on-the-mountain
forget-me-not	stock
four o'clock	sunflower
gaillardia	sweet alyssum
gladiolus	sweet pea
golden glow	sweet sultan
impatiens	sweet william
ixora	vinca
lobelia	yarrow
lupine	zinnia

Weeds Controlled

Weeds controlled when applied prior to weed germination:

Annual Grasses

Common Name
annual bluegrass
barnyardgrass (watergrass)
crabgrass, large or hairy
crabgrass, smooth
goosegrass (silver crabgrass or crowfoot)
green foxtail
yellow foxtail

Scientific Name

<i>Poa annua</i>
<i>Echinochloa crus-galli</i>
<i>Digitaria sanguinalis</i>
<i>Digitaria ischaemum</i>
<i>Eleusine indica</i>
<i>Setaria viridis</i>
<i>Setaria glauca</i>

Broadleaf Weeds

Common Name
chickweed, common
knotweed, prostrate
lambsquarters, common
pigweed
purslane, common

Scientific Name

<i>Stellaria media</i>
<i>Polygonum aviculare</i>
<i>Chenopodium album</i>
<i>Amaranthus</i> spp.
<i>Portulaca oleracea</i>

Weeds Suppressed

Weeds suppressed when applied prior to weed germination:

Common Name

creeping spurge
creeping woodsorrel
panicum, Texas
prostrate spurge
pusley, Florida
spotted spurge
woodsorrel

Scientific Name

<i>Euphorbia serpens</i>
<i>Oxalis corniculata</i>
<i>Panicum texanum</i>
<i>Euphorbia humistrata</i>
<i>Richardia scabra</i>
<i>Euphorbia maculata</i>
<i>Oxalis</i> spp.

Application Directions

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

Uses

Established Turfgrass

Application Timing

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application. Apply this product 1 to 2 weeks prior to the germination of annual grass weeds. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor.

In areas where germination of the annual weed grass species continues for extended period of time, make a second application at the recommended rate 8 to 10 weeks after the first application on cool season turfgrass and 10 to 12 weeks after the first application on warm season turfgrass.

Broadcast Application Rates	Cool Season Turfgrass	Warm Season Turfgrass
Summer annual grass weeds (spring application)	174 to 349 lb/acre (4 to 8 lb/1000 sq ft)	349 lb/acre (8 lb/1000 sq ft)
Winter annual grass weeds (fall application)	174 to 349 lb/acre (4 to 8 lb/1000 sq ft)	349 lb/acre (8 lb/1000 sq ft)

This product may also be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time. For a split application, apply this product at 174 lb plus 174 lb per acre for cool season turfgrass and in the northern half of the warm season turfgrass area, or 233 lb plus 233 lb per acre in the southern half of the warm season turfgrass area. Make the second application 8 to 10 weeks following the 174 lb per acre rate and 10 to 12 weeks following the 233 lb per acre rate.

Specific Use Restrictions

- Do not make more than 2 applications of this product per year on turfgrass.
- **Maximum Single Application Rate:** 349 lb of Lebanon Pro Brand fertilizer with Team® Pro 0.86% preemergence crabgrass control per acre.
- **Minimum Re-Treatment Interval:** Cool season turf – 56 to 70 days; warm season turf – 70 to 84 days.
- This product must be watered in as soon as possible after application.

Established Ornamental Plants in Landscape Settings

Apply to ornamental trees, shrubs, groundcovers and flowers, and to non-bearing fruit and nut trees and non-bearing vineyards, which are defined as plants that will not bear fruit for at least 1 year after treatment.

Application Timing

Apply 1 to 2 weeks prior to the germination of weed seeds or clean cultivate and hoe to remove existing weeds before applying this product. Apply to the smooth soil surface (no large clods) to ensure uniform distribution around established plants or on ornamental beds.

Broadcast Application Rate

Apply this product at a rate of 4 to 8 lb per 1000 sq ft as a spring application or 4 to 8 lb per 1000 sq ft as a fall application. In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application. Use the higher rate in the rate range where heavy weed infestation is anticipated. Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours after application.

Specific Use Restrictions

- Do not make more than 2 applications of this product per year on all ornamental plants.
- **Non-bearing fruit and nut trees, non-bearing vineyards:**
  - **Maximum Single Application Rate:** 8 lb of Lebanon Pro Brand fertilizer with Team® Pro 0.86% preemergence crabgrass control per 1000 sq. ft.
  - Applications to plants that will bear fruits, berries, or nuts within 12 months are prohibited.
- **Landscape ornamentals:**
  - **Maximum Single Application Rate:** 8 lb of Lebanon Pro Brand fertilizer with Team® Pro 0.86% preemergence crabgrass control per 1000 sq. ft.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT PERMITTED BY LAW, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.