

# Lebanon Pro



## FERTILIZER with DIMENSION® 0.13% PRE & POSTEMERGENCE ANNUAL GRASSY WEED CONTROL

# 0-0-7

### GUARANTEED ANALYSIS

Soluble Potash (K<sub>2</sub>O) ..... 7%

Derived From: Muriate of Potash.

Chlorine (Cl) not more than ..... 7.0%



Information regarding the contents and levels of metals in this product is available on the Internet at:  
<http://www.regulatory-info-lebsea.com>

SUGGESTED SPREADER SETTINGS					
Spreader	3.2# K <sub>2</sub> O/1,000 sq. ft.	6.4#K <sub>2</sub> O/1,000 sq. ft.	Spreader	3.2#K <sub>2</sub> O/1,000 sq. ft.	6.4#K <sub>2</sub> O/1,000 sq. ft.
LebanonTurf	4 1/4	5 1/2	Lesco (letter/Numeric)	I/17	L/24
Andersons AccuPro	K	P	PennMulch HVO	L	P
Earthway Rotary	15	20	ProScape SS	L	P
Gandy	26	32	Spyker	4 1/4	5 1/2
Lely	5 II	7 II	Vicon (all models)	24	31

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

MESA®, PennMulch® and ProScape® are registered trademarks of Lebanon Seaboard Corporation.

Dimension® is a selective herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes.

### ACTIVE INGREDIENT:

\*Dithiopyr..... 0.13%

OTHER INGREDIENTS:..... 99.87%

TOTAL ..... 100.00%

\*This product contains 0.065 lb. of the active ingredient, dithiopyr, per 50 lb. bag  
Product protected by U.S. Patent No. 4,692,184. Other patents pending.

EPA Reg. No. 961-360  
Dimension is a registered trademark of Dow AgroSciences LLC.

Cas No. 97886-45-8  
(L) (D)

EPA Est. No. 961-PA-01:961-IL-01  
Superscript used is first letter of lot number.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT.  
USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

Before buying this product, read the Warranty Section elsewhere on this label. If terms are not acceptable, return the unopened container to the place of purchase at once.

In New York State, this product may only be used by Professional Applicators at no more than 8.8 lbs. per 1000 sq. ft. (0.5 lb. active ingredient per acre) per year; and is prohibited for use in Nassau and Suffolk Counties.

## KEEP OUT OF REACH OF CHILDREN CAUTION

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

### FIRST AID

- IF IN EYES:**
- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
  - Call a poison control center or doctor for treatment advice.
- IF INHALED:**
- Move person to fresh air.
  - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
  - Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Emergency Medical Information call (888) 208-1368. For any other emergency involving this product, call day or night (800) 424-9300.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

**NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION, OR FOR RESEARCH PURPOSES.**

### PRODUCT INFORMATION

This product is a selective herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds as listed in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

### USE PRECAUTIONS

#### Turfgrass Safety

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury.

Apply this product only to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

#### Cool-Season Grasses

Bentgrass, creeping\* (*Agrostis palustris*)  
Bluegrass, Kentucky (*Poa pratensis*)  
Fescue, fine\*\* (*Festuca rubra*)  
Fescue, tall (*Festuca arundinacea*)  
Ryegrass, perennial (*Lolium perenne*)

#### Warm-Season Grasses

Bahiagrass (*Paspalum notatum*)  
Bermudagrass (*Cynodon dactylon*)  
Buffalograss\*\*\* (*Buchloe dactyloides*)  
Carpentgrass (*Axonopus affinis*)  
Centipedegrass (*Eremochloa ophiuroides*)  
Kikuyugrass (*Pennisetum clandestinum*)  
St. Augustinegrass (*Stenotaphrum secundatum*)  
Zoysiagrass (*Zoysia japonica*)

\* Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested.

**DO NOT** apply this product to Colonial bentgrass (*Agrostis tenuis*) varieties.

\*\* Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf', and 'Waldina'. Not all varieties of fine fescue have been tested.

\*\*\* **DO NOT** use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

NOTE: It is not possible to evaluate all turfgrass varieties for tolerance to Dimension. It is suggested that the user evaluate the response of turfgrass varieties to this product on a limited area, at specified use rates, prior to initiating large-scale use.

Manufactured by:  
**Lebanon Seaboard Corporation**  
1600 E. Cumberland St. • Lebanon, PA 17042  
Made in the U.S.A.  
800-233-0628 • (717) 273-1685

# Net Weight 50 LBS. (22.7 kg)

**RESEEDING, OVERSEEDING, OR SPRIGGING**

Reseeding, overseeding, or sprigging of treated areas with this product is to be delayed until 12 weeks from the time of application. Reseeding, overseeding or sprigging before 12 weeks after application may prevent establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization are to be followed. If overseeding with perennial ryegrass in the fall, overseed 8 weeks or later after a single application. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

**OTHER USE PRECAUTIONS**

The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller. For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, are to be done before application of this product. **DO NOT** apply this product until the grass has recovered from these cultural practices.

- **DO NOT** use clippings from treated turf for mulching around vegetables or fruit trees.
- Keep people and pets off treated areas until dust has settled.

**APPLICATION DIRECTIONS  
Control of Crabgrass**

**Preemergence and Early Postemergence Control**

This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs. It can also provide "early postemergence" control of crabgrass. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first see the crabgrass plants in the lawn or turf. The practical benefit of this product's early postemergence activity is that it can give the user a 2-4 week longer period of time to make applications to control crabgrass.

**Application Frequency and Timing**

This product may be applied as a single application or as a split application for crabgrass control in the spring, summer, or fall.

**Spring Applications**

For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the control programs listed in **Table 1** below, depending on the user's location, the turfgrass mowing height, and whether the use is preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

- Use Program 1 (given in Table 1, below) for preemergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.
- \*Use Program 2 (given in Table 1, below) for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control of crabgrass.
- Where split fertilizer applications are suggested, a rate in Table 1 may be applied in two applications made 6-10 weeks apart and prior to crabgrass emergence.

**Table 1: Application Use Rates  
for Crabgrass\*  
(Lbs. of this Product per 1,000 Sq. Ft.)**

PROGRAM	1	2
<b>USE</b> (turfgrass cut)	<b>PREEMERGENCE</b> (high)	<b>PREEMERGENCE</b> (low)
		<b>POSTEMERGENCE</b>
<b>REGION</b>		
<b>North</b>	2.2	4.4
<b>Transition</b>	3.2	6.4
<b>South</b>	4.4	8.8
<b>Coastal South</b>	6.4	8.8**

**Table 2: Application Use Rates  
For Fall Applications  
(Lbs. of Product per 1,000 Sq. Ft.)**

REGION	PREEMERGENCE
<b>North</b>	6.4
<b>Transition</b>	8.8*
<b>South</b>	8.8*
<b>Coastal South</b>	8.8*

**Fall Applications**

This product can also be applied in the late summer or early fall (late August through November) at the use rates listed in **Table 2** to provide control of crabgrass through the summer.

The fall application should be followed by an appropriate spring application to provide season-long control.

\*DO NOT apply more than 8.8 lbs. of this product per 1000 sq. ft. per application, and no more than 26.4 lbs. of this product per 1000 sq. ft. per year.

In New York State, this product may only be used by Professional Applicators at no more than 8.8 lbs. per 1000 sq. ft. (0.5 lb. active ingredient per acre) per year; and is prohibited for use in Nassau and Suffolk Counties.

Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below.

**North:** All other areas.

**Transition:** DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, and coastal areas of CT, NY, and RI.

**South:** AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, and TX.

**Coastal South:** HI, FL, and southern coastal areas of AL, GA, LA, MS, NC, SC, and TX.

\*\* Use split or sequential applications for full control.

**Tips for Improved Control**

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result.

Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

**Control of Other Grass and Broadleaf Weeds**

**Spring Applications**

Used as directed for crabgrass control in the spring (Table 1), this product will also control or suppress the following weeds when applied prior to their emergence:

Barnyardgrass	Goosegrass	Oxalis (buttercup, creeping & yellow woodsorrel)	Smutgrass
Bluegrass (annual, roughstalk)	Kikuyugrass	Purslane (common)	Speedwell (corn)
Crowfootgrass	Lespedeza (common)	Ryegrass (annual & perennial)	Spurge (prostrate & spotted)
Foxtail (yellow & green)*			

\* Also controlled at the Program 2 rates.

**Fall Applications**

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring (Table 2), this product will also control or suppress the following weeds when applied prior to their emergence:

Bluegrass (annual, roughstalk), Bittercress, Chickweed, Geranium (Carolina), Henbit, Parsley-piert, Pineappleweed, Shepherdspurse

**APPLICATION EQUIPMENT INSTRUCTIONS**

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate listed above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

**STORAGE AND DISPOSAL**

**Do not contaminate water, food, or feed by storage or disposal.**

**PESTICIDE STORAGE:** Store product in its original container in a cool, dry, locked place out of the reach of children and animals.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

**WARRANTY**

**NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.**