SLEBANON SCALLE.

Eco-Blend with Iron

15-2-5





Iron (Fe) 2.0%

Derived From: Ammonium Sulfate, Biosolids, Methylene Ureas, Urea, Sulfate of Potash, Ferric Oxide, Ferrous Sulfate.

*12.2% Slowly available Nitrogen from Methylene Ureas and Biosolids.

12.2 /6 Slowly available withogen from methylene oreas and biosofius

GUARANTEED	MINIMUM	ANALYSIS
------------	---------	-----------------

lotal nitrogen (n)	15%
1.8% Ammoniacal Nitrogen	
5.8% Water Insoluble Nitrogen*	
1.0% Urea Nitrogen	
6.4% Other Water Soluble Nitrogen*	
Available Phosphate (P205)	2%
Soluble Potash (K ₂ 0)	5%
Sulfur (S)	3.8%
3.8% Combined Sulfur (S)	
Iron (Fe) actual	2.0%
0.2% Water Soluble Iron (Fe)	
Chlorine (CI) maximum	2.0%

Derived From: Ammonium Sulfate, Biosolids, Methylene Ureas, Urea, Sulfate of Potash, Ferric Oxide, Ferrous Sulfate.

*12.2% Slowly available Nitrogen from Methylene Ureas and Biosolids.

Canada

Application Rates:

To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 6.0 lbs per 1,000 sq. ft. or 261 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 3.3 lbs per 1,000 sq. ft. or 144 lbs. per acre.

0.2% Water Soluble Iron (Fe)

SUGGESTED SPREADER SETTINGS			
Spreader 0.9#N	/1,000 sq. ft.	0.5#N /1,000 sq. ft.	
Lebanon Turf	41/2	31/2	
Andersons AccuPro	M	H	
Earthway Rotary	16	14	
Gandy	28	24	
Lely	6 II	4 II	
Lesco (letter/numeric)	J/21	H/15	
PennMulch HVO	M	J	
ProScape SS	M	J	
Spyker	41/2	31/2	
Vicon	26	22	
These settings were calibrated and field tested			

These settings were calibrated and field tested.
However, age and condition of spreader, speed of
operation and evenness of terrain may require slightly
different settings for desired coverage.

Application Rates and Frequencies in Canada:
Follow your professional landscape management
plan for your local course or area. A suggested
application rate is 1 to 2.8 Kg for 100 m2, applied
every 4 to 8 weeks during the local growing season.

U.S.A.

NOTICE: This product contains the secondary nutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing.

Do not wash off with water.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Application Rates:

To feed at the rate of 0.43 kg Nitrogen (N) Equivalent to 4.7 kg of product per 100 sq. meters.

To feed at the rate of 0.24 kg Nitrogen (N) Equivalent to 1.3kg of product per 100 sq. meters.

Equivalent to 1.5kg of product per 100 sq. meters.				
SUGGESTED SPREADER SETTINGS				
Spreader	0.43 kg N/100 m ²	0.24 kg# N/100 m ²		
Lebanon Turf	41/2	31/2		
Andersons Accu	ıPro M	Н		
Earthway Rotar	y 16	14		
Gandy	28	24		
Lely	6 II	4 II		
Lesco (letter/nu	meric) J/21	H/15		
PennMulch HVO	M	J		
ProScape SS	M	J		
Spyker	41/2	31/2		
Vicon	26	22		

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Hazard Statements

May be harmful if inhaled repeatedly over long periods. Keep out of reach of children.

Precautionary Statements

Avoid Breathing dust.
Seek medical advice or attention if you feel unwell.

Dispose in accordance with all federal, state and local regulations.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

Apply only to actively growing turf. Do not exceed 3.2 lbs. of total nitrogen per 1,000 sq. ft. annually.

In Maryland, New York and Virginia: This fertilizer contains phosphorus and may not be used for lawn maintenance on turf in the states of Maryland, New York or Virginia except when

1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory* performed no more than 3 years before the application;

2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) Re-establishing or repairing a turf area.

*In Maryland, the laboratory must be identified under Section 8-803.7 of the Agriculture Article, Annotated Code of Maryland.

In New Hampshire: This fertilizer contains phosphate. In New Hampshire, phosphate fertilizers may only be used for turf maintenance in golf courses, parks, athletic fields, and sod farms.



Manufactured by:
Lebanon Seaboard Corporation
1600 E. Cumberland St. • Lebanon, PA 17042
Made in the U.S.A.



Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com

NET WEIGHT



LBS. (18.14 Kg)