PRODIAMINE GROUP 3 HERBICIDE



terrain may require slightly different settings for desired coverage.

For selective preemergence control of grass and broadleaf weeds as listed in:

- established turfgrasses (excluding golf course putting greens) and lawns
- landscape ornamentals (including established perennials & wildflower plantings) ACTIVE INGREDIENT:

Prodiamine		0.43%
OTHER INGREDIENTS:	ç	99.57%
TOTAL:		0.00%
	(1)	

EPA Reg. No. 961-362

EPA Est. No. 961-PA-01:961-IL-01 Superscript used is first letter of lot number.

Barricade is a registered trademark of a Syngenta Group Company.

ProScape® and MESA® are registered trademarks of Lebanon Seaboard Corporation.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

Before buying this product, read the Warranty Section elsewhere on this label.

If terms are not acceptable, return the unopened container to the place of purchase at once.

KEEP OUT OF REACH OF CHILDREN

CAUTION PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

	0	-		
к	S	L	А	IC

IF IN EYES:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
	Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	Take off contaminated clothing.
	Rinse skin immediately with plenty of water for 15-20 minutes.
	Call a poison control center or doctor for treatment advice.
IF INHALED:	Move person to fresh air.
	• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
	Call a poison control center or doctor for further treatment advice.
Have the product container of	r label with you when calling a poison control center or doctor, or going for treatment.
You may also contact 888-20	8-1368 for emergency medical treatment information

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. **DO NOT** apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. **DO NOT** contaminate water when disposing of equipment washwaters. If material is spilled: Contain and sweep up material of spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill.

Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants not listed on this label and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE

This product is a selective preemergence herbicide that provides residual control of many listed grass and broadleaf weeds in established turfgrasses (excluding golf course putting greens) and lawns, landscape ornamentals, established perennials and wildflower plantings. This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

DO NOT graze or feed livestock forage cut from areas treated with this product.

DO NOT apply this product through any type of irrigation system.

DO NOT apply aerially.

DO NOT apply to golf course putting greens.

Manufactured by: Lebanon Seaboard Corporation 1600 E. Cumberland St. • Lebanon, PA 17042 800-233-0628 • (717) 273-1685

www.LebanonTurf.com

NET WEIGHT: 50 LBS. (22.7 KG)

WEED RESISTANCE MANAGEMENT

For resistance management, Lebanon ProScape Fertilizer with Barricade 0.43% Preemergence Weed Control is a Group 3 herbicides. The resistant biotypes may dominate the weed population may contain certain or develop plants naturally resistant to Lebanon ProScape Fertilizer with Barricade 0.43% Preemergence Weed Control and other Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps: Rotate the use Lebanon ProScape Fertilizer with Barricade 0.43% Preemergence Weed Control or other Group 3 herbicides within a growing season with different herbicide groups that control the same weeds.

Adopt an integrated weed management program for herbicide use that includes scouting and uses historical information related to herbicide use, and that considers cultural, biological varieties and other turf management practices.

Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.

If weed population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.

Contact your sales representative, pest control advisors, or local extension specialist for additional pesticide resistance management and/or integrated weed management recommendations for specific types of plants and weed biotypes.

APPLICATION DIRECTIONS

Apply uniformly with suitable, calibrated application equipment.

ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turfgrasses and lawns. The maximum amount of this product that may be applied per year is given for each turfgrass species in the **Maximum Annual Rates** section of this label. Most effective weed control in turfgrasses will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.



USE PRECAUTIONS

The following precautions apply to the use of this product in turfgrasses and lawns:

Application of this product may thin emerged annual bluegrass and newly overseeded grasses.

DO NOT apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil. **DO NOT** apply to newly set sod until the following year.

Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.

Disturbing the herbicide barrier with cultural practices such as discing may result in reduced weed control.

DO NOT apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (Poa annua) are desirable species.

RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Maximum use rate selection must be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES

This product is for use on the turfgrass species listed in the following table. DO NOT exceed the maximum yearly rate as given in the following table:

MAXIMUM APPLICATION RATE/CALENDAR YEAR of LEBANON PROSCAPE FERTILIZER WITH BARRICADE 0.43% PREEMERGENCE WEED CONTROL by Turfgrass Species¹⁾

Turf Species:	Lbs. Product/Acre	Lbs. Product/1,000 Sq. Ft.	Lbs. Al/Acre
Bermudagrass ²⁾ , Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turf-type), Zoysia	349	8.0	1.5
Buffalograss, Kentucky Bluegrass, Perennial Ryegrass	233	5.3	1.0
Creeping Red Fescue	175	4.0	0.75
Creeping Bentgrass	150	3.4	0.65

¹⁾ These are the maximum rates per calendar year by species limitations.

²⁾ May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 121 lbs./Acre. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescue grass due to reduced product rates used in sprigging situations.
 • DO NOT apply more than 349 lbs. of this product per calendar year.

• Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but DO NOT exceed the maximum application rates specified in the Maximum Annual Rates Table.

In the states of MD, NJ, VA and WA apply this product at no more than 5.3 lbs. of product per 1,000 sq. ft.

When used in accordance with this labe	I this product will provide control of the following we	eds:					
Barnyardgrass	Chickweed, Mouseear (from seed)	Foxtails, Annual	Johnsongrass (from seed)	Lambsquarters, Common	Purslane, Common	Speedwell, Persian	
Bluegrass, Annual (Poa annua)	Crabgrass (large, smooth)	Goosegrass ¹¹	Junglerice	Lovegrass	Pusley, Florida	Sprangletop	
Carpetweed	Crowfootgrass	Henbit	Knotweed	Panicum (Texas, Fall, Browntop)	Rescuegrass ³³	Spurge, Prostrate	
Chickweed, Common	Cupgrass, Woolly	Itchgrass	Kochia	Pigweed	Shepherd's Purse ²²	Witchgrass	
					Signalgrass Broadleaf	Woodsorrel Yellow (from seed)	

¹⁾ In many areas a single application of 150 to 349 lbs./Acre of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 150 to 233 lbs./Acre of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 150 to 233 lbs./Acre of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 150 to 233 lbs./Acre of the maximum annual Rate Table. DO NOT exceed the maximum annual Rate Table above.

WEEDS CONTROLLED

²⁾ Apply this product in late summer, fall, or winter prior to germination.

³⁾ Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates listed for each turf species. All applications must be made prior to germination of the weed seeds

WHEN TO APPLY AFTER OVERSEEDING TURF

DO NOT apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

L

LEBANON PROSCAPE FERTILIZE PREEMERGENCE WEED CON			Months Before	Overseeding
Product	Á	North	Transition	South
116	.50	4	4	4
150	.66	5	4	4
175	.74	6	5	5
185	.80		6	6
233	1.00		7	7
265	1.14			9
302	1.30			10
349	1.50			12

LANDSCAPE ORNAMENTALS (including Established Perennials and Wildflower Plantings)

This product may be applied for residual preemergence weed control in ornamentals.

USE RATES

Apply this product at 150 to 349 lbs./Acre in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications are to be made so long as the total amount of product applied does not exceed the maximum annual application rate of 349 lbs./Acre.

APPLICATION TIMING AND INFORMATION

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following

application

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

ESTABLISHED LANDSCAPE ORNAMENTALS

Apply this product in the fall and/or spring to established ornamentals as a broadcast, over the top or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

NEWLY TRANSPLANTED LANDSCAPE ORNAMENTALS

Apply this product in the fall and/or spring to newly transplanted ornamentals as a broadcast, over the top or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

TOLERANT ORNAMENTAL SPECIES

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. Apply this product over the top of these species.

product over the top of these species. Scientific Name	Common Name	Scientific Name	Common Name	Scientific Name	Common Name	Scientific Name	Common Name
Abies spp.	Fir species**	Cupressus sempervirens	Italian Cypress	Justicia brandegeeana	Shrimp Plant	P. tobira	Japanese Pittosporum
Ables spp.	(Balsam, Fraser, Noble, etc.)	Delosperma alba	White Trailing Ice Plant	Lagerstromia indica	Crepe Myrtle	Podocarpus macrophyllus	Japanese Yew
Abelia grandiflora	Abelia: Sherwood	Delosperma sp.	Cooperi Pink	Lagerstomia indica X Fauriei	Crepe Myrtle; Tuscarora	Prunus laurocerasus	English Laurel
0 5	Japanese Maple	Delphinium sp.	Larkspur: Blue Elf	Lantana montevidensis	Weeping Lantana	Prunus sp.	Almond, Apricot, Nectarine, Peach,
	Norway Maple	Dianthus deltoides	Dianthus; Maiden Pinks "Zing"	Lavender sp.	Lavender; Munstead	1 · · · · · · · · · · · · · · · · · · ·	Plum and Prune*
	Yarrow: King Edward	D. gratianopolitanus	Cheddar Pink	Leontopodium alpinum	Edelweiss	Pseudotsuga menziesii	Douglas Fir**
	Kiwi*	D. grananoponianus Dodonea viscosa	Hop Bush	Ligustrum sinense	Chinese Privet; Variegata	Pyracantha coccinea	Firethorn (Scarlet)
Agapanthus africanus	NIWI	Echinacea pupurea	Coneflower, Purple; Magnus	Ligustrum amurense	Amur Privet	P. Fortuneana	Chinese Firethorn
0 I U	Lily-of-the-Nile	Elaeagnus commutata	Silverberry	L. japonicum	Japanese Privet	P. Koidzumii	Firethorn: Formosa; Taiwan
~ ~ ~ ~	Five-Leaf or Chocolate Vine	Elaeagnus pungens	Silverthorn	L. lucidum	Glossy Privet (Waxleaf)	Pyrus sp.	Bradford Pear sp.
	Lady's Leek, Nodding Onion	Euonymus fortunei	Wintercreeper	Lillium sp.	Lily: Jazz	Quercus shumardii	Oak, Shumard's Red
	Japanese Anemone	E. japonicus	Japanese Spindle Tree	Liriope Muscari	Big Blue Lily; Turf; Silver Mound;	Quercus spp.	Oak species
Aquilegia sp.	Aquilegia: Red and Gold	E. jupomens	(Evergreen Euonymus)	Entope museuri	Evergreen Giant	\tilde{R} haphiolepis indica	Indian Hawthorne
	Vine Hill Manzanita	E. kiautschovica	Creeping Strawberry Bush;	Liriope Muscari var. variegata	Liriope, Variegated	Rhaphiolepis umbellata	Yedda Hawthorn
1 U U	Cape Weed	E. Rumsenovieu	Manhattan Euonymus	Liriope spicata	Liriope, Creeping	Rhododendron	'Coral Bells'
	Wormwood; Silver Mound, Castle	Fatsia japonica	Japanese Aralia	Lobelia Cardinalis	Cardinal Flower, Indian Pink	(including Azalea)	'Delaware Valley White'
Aster sp.	Aster: Bonny Blue, Purple Dome	Forsythia x intermedia	Border Forsythia	Lonicera japonica	Japanese Honeysuckle	Rosa Banksiae	'Lady Bank's Rose
	Frikarti's aster	Forysythis suspensa	Weeping Forsythia	L. tatarica	Tatarian Honeysuckle	Rosmarinus officinalis	Rosemary*
	Lady Fern; Fern Lady	F. viridissima	Greenstem Forsythia	Loropetalum chinense	Burgundy	Rudbeckia sp.	Black-Eyed Susan; Goldstrum
	Japanese Aucuba	Gaillardia spp.	Gaillardia, Blanket Flower:	Lythrum sp.	Loosestrife; Modern Pink	Santolina virens	Santolina
	Fibrous Begonia: Hardy Grandis	Gumarana spp.	"Goblin"	Magnolia spp.	Magnolia species**	Saxifraga sp.	Saxifrage; Purple Dome
	William Penn Barberry	Gardenia jasminoides	Gardenia, Cape-Jasmine,	Maleophora luteola	Rocky Point Ice Plant	Scabiosa sp.	Pincushion Flower
	Wintergreen Barberry	Surdenia Jushino aes	Radicans	Malus sp.	Crabapple*	Sedum album	White Stonecrop
	Mentor Barberry	Gaura sp.	Tradicans	Miscanthus sinensis	Yaku Jima, Silberfeder	Sedum cauticola	Cliff Stonecrop; Lidakense
	Japanese Barberry	Gentiana dahurica	Gentian	Nandina domestica	Heavenly Bamboo	Sedum dasyphyllum	Corsican Stonecrop
	Warty Barberry	Geranium cinereum	Ashy Cranesbill	Narcissus spp.	Narcissus species**	Sedum spurium	Stonecrop; Dragon's Blood
Bergenia filix-femina	Pigsqueak	Gladiolus spp.	Gladiolus species**	Nerium oleander	Oleander	Senna artemisoides	Feathery Cassia
Boltonia asteroides	Snowbank,; False Aster	Gypsophila repens	Baby's Breath	Olea europaea	Olive*	Spiraea Bumalda	Spirea: Anthony Waterer
	Butterfly-Bush (Dwarf Blue);	Hedera helix	English Ivy	Ophiopogon japonicus	Mondo Grass	Syzygium paniculatum	Australian Brushcherry
	Royal Red	Hellanthemum sp.	Sunrose	Osmanthus heterophyllus	Osmanthus (False Holly): Gulf Tide	Taxus cuspidata	Japanese Yew
	Japanese Boxwood	Hemerocallis sp.	Daylily: Aztec Gold, Stella De	Osteospermum fruticosum	Trailing African Daisy	T. media	Yew
	Weeping Bottlebrush		Oro, Tender Love	Oxydendrum arboreum	Sourwood	Teucrium sp.	Germander
	Scotch Heather	Hibiscus syriacus	Rose of Sharon**	Paeonia suffruticosa	Tree Peony; Moutan Peony	Thalictrum dipterocarpum	Meadow Rue
Campanula carpatica	Tussock Bellflower, (White Clips)	Hibiscus sp.	Mallow: Disco Belle White	Pennisetum setaceumi	Fountain Grass (Dwarf)	Thuja occidentalis	American Arborvitae
	Trumpet Creeper, Trumpet	Hibiscus rosa-sinensis	Chinese Hibiscus	Perovskia atriplicifolia	Russian Sage	Trachelospermum asiatum	Star Jasmine
	Flower, Madame Galen	Hosta plantaginea	Hosta, Plantain Lily (Fragrant)	Persea americana	Avocado*	Tsuga canadensis	Canada Hemlock
	Hottentot Fig (Ice Plant)	H. Sieboldiana	Hosta, "Seersucker"	Photinia Fraseri	Frasier's Photinia (Redtip)	Tulipa spp.	Tulip species
	Wild Lilac	Houttuynia cordata var.	Chameleon Plant	Physotegia Viginiana	Dragonhead, False; Vivid	Veroniac sp.	Veronic, Speedwell; Sunny Border
Ceratostigma plumbagonoides		Hydrangea macrophylla	Bigleaf Hydrangea	Picea spp.	Spruce species**	Viburnum japonicum	Japanese Viburnum
· · ·	False Cypress	Ilex Bernyi	Perny Holly	11	(Colorado Blue, Norway, etc.)	V. denatum	Arrowhead Viburnum
Chrysanthemum nipponicum		Ilex cornuta	Chinese Holly	Pieris japonica	Lily-of-the-Valley Bush	V. odoratissimum	Sweet Viburnum
Cleyera japonica	Cleyera	I. crenata	Japanese Holly; Helleri	Pinus brutia	Calabrian Pine	V. plicatum	Japanese Snowball
Citrus spp.	Citrus species*	I. opaca	American Holly	P. canariensis	Canary Island Pine	V. rhytidophyllum	Leatherleaf Viburnum
Coreopsis spp.	Coreopsis (Calliopsis): Early	I. vomitoria	Yaupon Holly; Schillings	P. elliottii	Slash Pine	V. rigidum	Laurustinus
	Sunrise, Moonbeam	Inula ensifolia	Swordleaf Inula	P. halepensis	Aleppo Pine	V. Suspensum	Sandankwa
Cornus florida	Flowering Dogwood American	Iris ensata	Sword-Leaved Iris; Jodlesong	P. nigra	Austrian Pine	V. dentatum	Canary Island Viburnum
5	Dogwood	Iris siberica	Siberian Iris; Cabernet	P. palustrus	Longleaf Pine	V. tinus	Cranberry Bush
C. stolonifera	Red Osier Dogwood	Iris spp.	Iris species**	P. radiata	Monterey Pine	V. trilobium	Tulip species
Cortaderia selloana	Pampas Grass	Jasminium nudiflorum	Winter Jasmine	P. thunbergii	Japanese Black Pine	V. wrightii	White Viburnum
	Cranberry Cotoneaster	Juniperus chinensis	Chinese Juniper; Nick's	P. strobus	Eastern White Pine	V. rhytidophyllum	Leatherleaf Viburnum
C. buxifolius	Cotoneaster		Compact, Parsonii	P. syvestris	Scotch Pine	Vinca major	Greater Periwinkle, Vinca
	Bearberry Cotoneaster	J. conferta	Shore Juniper; Blue Pacific	P. taeda	Loblolly Pine	Vinca minor	Common Periwinkle, Vinca
C. Dammeri					Virginia Pine	Vitis sp.	
C. Dammeri C. microphyllus	Littleleaf Cotoneaster	J. davurica	Parson's Juniper	P. virginiana	Virginia i no	viiis sp.	Grape
C. microphyllus	Littleleaf Cotoneaster Hawthorne	J. davurica J. horizontalis	Parson's Juniper Creeping Juniper	P. virginiana Pistachio sp.	Pistachio*	Weigela florida	Grape* Old Fashioned Weigela
C. microphyllus Cretaegus spp.	Littleleaf Cotoneaster Hawthorne Lucifer	J. davurica J. horizontalis Juglans sp.	Parson's Juniper Creeping Juniper Walnut*	P. virginiana Pistachio sp. Pittosporum rhombifolium	-		Old Fashioned Weigela Spanish Bayonet, Yucca

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store product in its original container in a cool, dry, locked place out of the reach of children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. CONTAINER HANDLING: Nonrefillable container. DO NOT reuse or refill this container. Offer for recycling if available. Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed be available.

by State and local authorities, by burning. If burned stay out of smoke.

WARRANTY NOTE: To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions. * Do not use on food producing trees, vines, or plants. ** Not for use on container grown plants.