LebanonTurf

Country Club Maximum Dispersion

- Utilizes a unique, highly dispersing granule technology
- Contains North Atlantic Ocean sea kelp
- Features METH-EX slow release nitrogen
- SGN 125 is ideal for low-cut turf on tees and fairways

FERTILIZER

GUARANTEED ANALYSIS Total Nitrogen (N) 12.0% Ammoniacal Nitrogen 1.0% Water Insoluble Nitrogen*	16%
2.2% Urea Nitrogen 1.8% Other Water Soluble Nitrogen * Available Phosphate (P ₂ O ₅) Soluble Potash (K ₂ O) Magnesium (Mg)	8%
0.8% Water Soluble Magnesium (Mg) Sulfur (S)	
0.2% Water Soluble Iron (Fe) Manganese (Mn) 0.5% Water Soluble Manganese (Mn)	
Derived From: Ammonium Sulfate, Ammonium Phosphate, Methylene Ureas, Ure Sulfate of Potash, Kelp Meal, Sulfate of Potash Magnesia, Ferric Oxide, Ferrous Manganese Sulfate.	
*2.8% Slowly Available Nitrogen from Methylene Urea. Chlorine (CI) not more than	2.0%
SOIL/PLANT AMENDING INGREDIENTS Active Ingredients: Humic Acids (derived from Leonardite)	
Inert Ingredients (as soil amending)	U.S.A lity.

GUARANTEED MINIMUM ANALYSIS Total Nitrogen (N)	. 16%
1.8% Other Water Soluble Nitrogen * Available Phosphate (P ₂ 0 ₅) Soluble Potash (K ₂ 0) Magnesium (Mg)	. 8 %
0.8% Water Soluble Magnesium (Mg) Sulfur (S)	
0.2% Water Soluble Iron (Fe) Manganese (Mn) (actual)	
Derived From: Ammonium Sulfate, Ammonium Phosphate, Methylene Ureas, Ur Sulfate of Potash, Kelp Meal, Sulfate of Potash Magnesia, Ferric Oxide, Ferrous Manganese Sulfate.	ea,
*2.8% Slowly Available Nitrogen from Methylene Urea.	
ALSO CONTAINS NON PLANT FOOD INGREDIENTS SOIL/PLANT AMENDING INGREDIENTS Active Ingredients:	
Humic Acids (derived from Leonardite)	CANADA ility.

NOTICE: This product contains the secondary micronutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Application Rates:

To feed at the rate of 0.9 lb.Nitrogen (N) per 1,000 sq. ft., apply this product at 5.6 lbs per 1,000 sq. ft. or 244 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 3.1 lbs per

1,000 Sq. It. 01 133 Ibs. per acre.					
SUGGESTED SPREADER SETTINGS					
Spreader 0.	5#N/1,000 sq. ft.	0.9#N/1,000 sq. ft.			
Lebanon Turf	41/2	51/4			
Andersons AccuPr	o K	0			
Earthway Rotary	15	19			
Gandy	26	31			
Lely	5 II	6½ II			
Lesco (letter/nume	eric) i/17	K/23			
PennMulch HVO	L	0			
ProScape SS	L	0			
Spyker	41/2	5 ½			
Vicon	24	30			

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Apply only to actively growing turf. Do not exceed 3.2 lbs. of total nitrogen per 1,000 sq. ft. annually.

In Maryland, New Jersey, New York and Virginia: This fertilizer contains phosphorus and may not be used for lawr maintenance on turf in the states of Maryland, New Jersey, New York or Virginia except when

1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory* performed no more than 3 years before the application;

2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established or 3) Re-establishing or repairing a turf area.

*In Maryland, the laboratory must be identified under Section 8-803.7 of the Agriculture Article, Annotated Code of Maryland.

In New Hampshire: This fertilizer contains phosphate. In New Hampshire, phosphate fertilizers may only be used for turf maintenance in golf courses, parks, athletic fields, and sod farms.

Warning

Hazard Statements Harmful if swallowed May be harmful if swallo Causes skin irritation. Causes eye irritation. May be harmful if inhaled

May cause respiratory irritation.

Wash hands and exposed skin thoroughly after handling. Do not eat, drink or smoke when using this product.

Wear protective gloves/protective clothing/eye protection/face protection. If in eyes, rinse with water and seek medical attention if symptoms persist. Rinse mouth.

IF SWALLOWED: Call a POISON CENTER or doctor/physician if you feel

IF ON SKIN: Wash with plenty of soap and water.

IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell If skin irritation occurs: Get medical advice/ attention. If eye irritation persists: Get medical advice/attention.

Store locked up. Store in a well-ventilated place.

Keep container tightly closed.

Dispose of contents/container properly as required by law for waste pesticides.

Keep out of reach of children

Dispose in accordance with all federal, state and local regulations. Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact

with skin or eyes, flush with plenty of water; for eyes get medical attention. institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommenda-

tions in your approved nutrient management plan Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

Do not apply to frozen turf.

WARRANTY NOTE To the extent consistent with applicable law, Buyer assumes all

responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:

To feed at the rate of 0.43 kg Nitrogen (N) Equivalent to 2.4 kg of product per 100 sq. meters.

To feed at the rate of 0.22 kg Nitrogen (N)

Equivalent to 1.3 kg of product per 100 sg. meters.

- 1	SUGGESTED SPREADER SETTINGS				
	Spreader 0.22	2 kg N/100 m².	0.43 kg N/100 m		
	Lebanon Turf	41/2	51/4		
	Andersons AccuPro	K	0		
	Earthway Rotary	15	19		
	Gandy	26	31		
	Lely	5 II	6½ II		
	Lesco (letter/numerio	c) i/17	K/23		
	PennMulch HVO	Ĺ	0		
	ProScape SS	L	0		
	Spyker	41/2	51/2		

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



21-29089

this product is available on the Internet at: http://www.regulatory-info-lebsea.com

For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation 1600 E. Cumberland St • Lebanon, PA 17042

800-233-0628 • (717-273-1685)