LebanonTurf

Country Club Bruce Visibility

- Increased granule visibility for easy application
- Utilizes unique Composite Technology®
- Features **METH**·**EX**[®] slow-release nitrogen
- SGN 80 granules are ideal for putting greens

FERTILIZER **17-0-1**7

GUARANTEED ANALYSIS

Total Nitrogen (N)17%
5.4% Ammoniacal Nitrogen
2.7% Water Insoluble Nitrogen*
4.5% Urea Nitrogen
4.4% Other Water Soluble Nitrogen*
Soluble Potash (K,O)17%
Iron (Fe) 1.0%
0.1% Water Soluble Iron (Fe)
Manganese (Mn) 0.5%
0.5% Water Soluble Manganese
Derived from: ammonium sulfate, methylene ureas, urea, sulfate of potash, iron sulfate, manganese sulfate. *7.1% slowly available nitrogen from methylene ureas.

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)17%
5.4% Ammoniacal Nitrogen
2.7% Water Insoluble Nitrogen*
4.5% Urea Nitrogen
4.4% Other Water Soluble Nitrogen*
Soluble Potash (K ₂ O)17%
Iron (Fe) (actual)1.0%
0.10% Water Soluble Iron (Fe)
Manganese (Mn) (actual)0.5%
0.5% Water Soluble Manganese
*7.1% slowly available nitrogen from methylene ureas.
Derived from: ammonium sulfate, methylene ureas, urea, sulfate of potash, iron sulfate, manganese sulfate.

NOTICE: This product contains the micronutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Application Rates: To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 5.0 lbs per 1,000 sq. ft. or 218	Warning Hazard Statements May be harmful if swallowed.
lbs. per acre. To feed at the rate of ½ lb. Nitrogen (N) per	Causes skin irritation.
1,000 sq. ft., apply this product at 2.8 lbs per 1,000 sq. ft. or 122 lbs. per acre.	May be harmful if inhaled.
SUGGESTED SPREADER SETTINGS	May cause respiratory irritation.
Spreader 0.9# N/1,000 sq. ft. 0.5 # N/1,000 sq. ft.	Causes serious eye irritation. Causes eye irritation.
	Precautionary Statements for handling: Wash hands and exposed skin thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection. If in eyes, rinse with water and seek medical attention if symptoms persist. Take off contaminated clothing and wash before reuse. IF ON SKIN: Wash with plenty of soap and water.
These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.	IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If skin irritation occurs: Get medical advice/ attention. If eye irritation persists: Get medical advice/attention. Store locked up.
	Store in a well-ventilated place. Keep container tightly closed. Keep out of reach of children

Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your turf, and sweep any product that lands on impervious surfaces back into your turf.

Do not apply to frozen turf.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m applied every 4 to 8 weeks during the local growing season.

Application Rates:

To feed at the rate of 0.41 kg Nitrogen (N) apply at 2.48 kg of product per 100 sq. meters To feed at the rate of 0.23 kg Nitrogen (N) apply at 1.37 kg of product per 100 sq. meters

SUGGESTED SPREADER SETTINGS				
Spreader		0.23 kg N/100 m ²		
Andersons Accul	Pro G	E		
Earthway Rotary	12	10		
Gandy	23	20		
Lely	3½	2		
Lesco (letter/nur	neric)F/16	D/14		
Spyker	2 ½	1½		
Vicon	18	15		

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com

For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation 1600 E. Cumberland St • Lebanon, PA 17042

800-233-0628 • 717-273-1685

40 Lbs. (18.14 kg) Net Weight

U.S.A