LebanonTurf

Country Club IV

- Increased granule visibility for easy application
- Utilizes unique Composite Technology®
- Features METH-EX® slow-release nitrogen
- SGN 80 granules are ideal for putting greens

GUARANTEED ANALYSIS Total Nitrogen (N) 2.0% Ammoniacal Nitrogen 3.9% Water Insoluble Nitrogen' 4.1% Urea Nitrogen 8.0% Other Water Soluble Nitrogen* Available Phosphate (P2O5) Soluble Potash (K₂O).....18% Magnesium (Mg).... 0.5% Water Soluble Magnesium (Mg) Sulfur (S) ... 6.2% Combined Sulfur (S) Iron (Fe). 0.05% Water Soluble Iron (Fe) Manganese (Mn) 0.5% Water Soluble Manganese erived from: urea, methylene ureas, ammonium phosphate, sulfate of potash, sulfate of potash magnesia, ferrous sulfate, manganese sulfate. *11.9% slowly available nitrogen from methylene ureas.

dominative minimon analisi	
Total Nitrogen (N)	18%
2.0% Ammoniacal Nitrogen	
3.9% Water Insoluble Nitrogen*	
4.1% Urea Nitrogen	
8.0% Other Water Soluble Nitrogen*	
Available Phosphate (P ₂ O ₅)	9%
Soluble Potash (K ₂ O)	18%
Magnesium (Mg) (actual)	0.5%
0.5% Water Soluble Magnesium (Mg)	
Sulfur (S)	6.2%
6.2% Combined Sulfur (S)	
Iron (Fe) (actual)	0.5%
0.05% Water Soluble Iron (Fe)	
Manganese (Mn) (actual)	0.5%
0.5% Water Soluble Manganese	
Chlorine (CI) (maximum)	2.0%
11.9% slowly available nitrogen from methylene ureas.	
Derived from: urea, methylene ureas, ammonium phosphate, sulfate	
of potash, sulfate of potash magnesia, ferrous sulfate, manganese sulfate.	CANADA

GUARANTEED MINIMUM ANALYSIS

U.S.A Chlorine (CI) not more than .. NOTICE: This product contains the micronutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Application Rates:

To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 5.0 lbs per 1,000 sq. ft. or 218 lbs. per acre. To feed at the rate of $\frac{1}{2}$ lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.8 lbs per 1,000 sq. ft. or 122 lbs. per acre.

SUGGESTED SPREADER SETTINGS			
Spreader	0.9# N/1,000 sq. ft.	0.5 # N/1,000 sq. f	
Andersons AccuPro G		E	
Earthway Rota	ary 12	10	
Gandy	23	20	
Lely	31/2 II	2 II	
Lesco (letter/numeric)F/16		D/14	
Spyker	2 ½	11/2	
Vicon	18	15	

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Apply only to actively growing turf. Do not exceed 3.2 lbs. of total nitrogen per 1,000 sq. ft. annually

In Maryland: This product is to to be used as a seed

In Maryland, New Jersey, New York and

Virginia: This fertilizer contains phosphorus and may not be used for lawn maintenance on turf in the states of Maryland, New Jersey, New York or Virginia except when

1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory* performed no more than 3 years before the application

2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State

3) Re-establishing or repairing a turf area.

*In Maryland, the soil testing laboratory must be identified under Section 8-803.7 of the Agriculture Article, Annotated Code of Maryland.

In New Hampshire: This fertilizer contains phosphate. In New Hampshire, phosphate fertilizers may only be used for turf maintenance in golf courses, parks, athletic fields, and sod farms.



May be harmful if swallowed. Causes skin irritation

May be harmful if inhaled. May cause respiratory irritation. Causes serious eye irritation.

Hazard Statements

Causes eve irritation. Precautionary Statements for handling:

Wash hands and exposed skin thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection. If in eyes, rinse with water and seek medical attention if symptoms persist.

Take off contaminated clothing and wash before reuse. IF ON SKIN: Wash with plenty of soap and water.

IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

If skin irritation occurs: Get medical advice/ attention. If eve irritation persists: Get medical advice/attention.

Store locked up.

Store in a well-ventilated place. Keep container tightly closed.

Keep out of reach of children

Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your turf, and sweep any product that lands on impervious surfaces back into your turf.

Do not apply to frozen turf.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m applied every 4 to § weeks during the local growing season.

Application Rates:

To feed at the rate of 0.41 kg Nitrogen (N) apply at 2.48 kg of product per 100 sq. meters To feed at the rate of 0.23 kg Nitrogen (N) apply at 1.37 kg of product per 100 sq. meters

SUGGESTED SPREADER SETTINGS				
Spreader	0.41 kg N/100 m ²	0.23 kg N/100 m ²		
Andersons AccuPro G		E		
Earthway Rot	tary 12	10		
Gandy	23	20		
Lely	3½ II	2 II		
Lesco (letter/numeric)F/16		D/14		
Spyker	2 ½	1½		
Vicon	18	15		

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com

For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation

1600 E. Cumberland St • Lebanon, PA 17042 800-233-0628 • 717-273-1685