# LebanonTurf

# Country Club® Blended Fertilizer

21-0-20

## **MESA**.

### **GUARANTEED ANALYSIS** 5.1% Ammoniacal Nitrogen

3.2% Water Insoluble Nitrogen\* 7.3% Urea Nitrogen 5.4% Other Water Soluble Nitrogen\*

**Application Rates:** 

Lebanon Turf

Gandy

Lely

Andersons AccuPro

Lesco (letter/numeric) i/19

**Earthway Rotary** 

PennMulch HVO

**ProScape SS** Spyker

Sulfur (S) ...... 5.8% Combined Sulfur (S)

To feed at the rate of 0.9lb. Nitrogen (N) per 1,000

sq. ft., apply this product at 3.7 lbs per 1,000 sq. ft. or 160.8 lbs. per acre. To feed at the rate of 0.5 lb.

Nitrogen (N) per 1,000 sq. ft., apply this product at 2.1 lbs per 1,000 sq. ft. or 90.6 lbs. per acre.

15

41/2

These settings were calibrated and field tested.

different settings for desired coverage.

However, age and condition of spreader, speed of

Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com

operation and evenness of terrain may require slightly

SUGGESTED SPREADER SETTINGS 0.9#N/1,000 sq. ft. 0.5#N/1,000 sq. ft.

Iron (Fe)..... 0.15% Water Soluble Iron Derived From: Methylene Ureas, Ammonium Sulfate, Urea, Muriate of Potash, Ferric Oxide

Ferrous Sulfate. \*8.5% Slowly Available Nitrogen from Methylene Ureas.

Chlorine (CI). Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are

determined by the professional applicator based on local soils, regulations, and climate.

removed from these areas promptly by sweeping or blowing. Do not wash off with water.

4 II

G/15

### **GUARANTEED MINIMUM ANALYSIS**

5.1% Ammoniacal Nitrogen 3.2% Water Insoluble Nitrogen'

7.3% Urea Nitrogen 5.4% Other Water Soluble Nitrogen\*

Soluble Potash (K<sub>2</sub>0)..... Sulfur (S) ... 5.8% Combined Sulfur (S) Iron (Fe) Actual.....

0.15% Water Soluble Iron Derived From: Methylene Ureas, Ammonium Sulfate, Urea, Muriate of Potash, Ferric Oxide,

Ferrous Sulfate.. \*8.5% Slowly Available Nitrogen from Methylene Ureas.

NOTICE: This product contains the micronutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be

Hazard Statements: Causes eye irritation

and local regulations.

Precautionary Statements for handling:

Wash hands and exposed skin thoroughly after handling. If in eyes, rinse with water and seek medical attention if symptoms persist. If eye irritation persists: Get medical advice/attention.

Keep out of reach of children Precautionary Statements for disposal - Dispose in accordance with all federal, state

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with

skin or eyes, flush with plenty of water; for eyes get medical attention. Use in accordance with recommendations of a qualified individual or institution. such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved

nutrient management plan. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

Do not apply to frozen turf.

NOTE To the extent consistent with applicable law. Buver assumes all responsibility for safety and performance if this product is not used according to the directions

WARRANTY

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m<sup>2</sup>, applied every 4 to 8 weeks during the local growing season.

**Application Rates:** 

To apply at the rate of 0.43 kg N/100 sq. meters. Equivalent to 1.8 kg of product per 100 sq. meters.

To apply at the rate of 0.22 kg N/100 sq. meters. Equivalent to 1.0 kg of product per 100 sq. meters.

SI	JGGESTED SPREADER	R SETTINGS
Spreader	0.43 kg N/100 m <sup>2</sup> .	0.22 kg N/100 m <sup>2</sup> .
Lebanon Turf	41/2	4
Andersons Acc	uPro L	H
Earthway Rota	ry 15	14
Gandy	27	24
Lely	51∕2 II	4 II
Lesco (letter/n	umeric) i/19	G/15
PennMulch HV	0 M	J
ProScape SS	M	J
Spyker	41/2	4
Vicon	25	22

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage



MESA®, PennMulch® and ProScape® are registered trademarks of Lebanon Seaboard Corporation.

**LebanonTurf** 

For technical assistance or more information about our products visit www.LebanonTurf.com

**Lebanon Seaboard Corporation** 

1600 E. Cumberland St • Lebanon, PA 17042 800-233-0628 • (717) 273-1685

50 Lbs. (22.7 kg) Net Weight

# 21-0-20 MESA

#### **GUARANTEED ANALYSIS**

UUAIIAN I EED ANAEI 313	
Total Nitrogen	21%
5.1% Ammoniacal Nitrogen	
3.2% Water Insoluble Nitrogen*	
7.3% Urea Nitrogen	
5.4% Other Water Soluble Nitrogen*	
Soluble Potash (K <sub>2</sub> 0)	20%
Sulfur (S)	5.8%
5.8% Combined Sulfur (S)	
Iron (Fe)	1.5%
0.15% Water Soluble Iron	
Derived From: Methylene Ureas, Ammonium Sulfate, Urea, Muriate of Potash,	Ferric Oxide,
Ferrous Sulfate.	
*8.5% Slowly Available Nitrogen from Methylene Ureas.	
Chlorine (CI)	17.0%
	U.S.A

GUARANTEED MINIMUM ANALYSIS	040/
Total Nitrogen 5.1% Ammoniacal Nitrogen	21%
3.2% Water Insoluble Nitrogen*	
7.3% Urea Nitrogen	
5.4% Other Water Soluble Nitrogen*	
Soluble Potash (K <sub>2</sub> 0)	20%
Sulfur (S)	
5.8% Combined Sulfur (S)	
Iron (Fe) Actual	. 1.5%
0.15% Water Soluble Iron	
Chlorine (CI)	. 17.0%
Derived From: Methylene Ureas, Ammonium Sulfate, Urea, Muriate of Potash,	Ferric Oxide,
Ferrous Sulfate	
*8.5% Slowly Available Nitrogen from Methylene Ureas.	CANADA

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

NOTICE: This product contains the micronutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.

#### WARNING

#### Hazard Statements:

Causes eye irritation.

#### Precautionary Statements for handling: Wash hands and exposed skin thoroughly after handling.

Wash marks and exposed soft micrography after hardning. If in eyes, rinse with water and seek medical attention if symptoms persist. If eye irritation persists: Get medical advice/attention.

Keep out of reach of children

Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

Do not apply to frozen turf.

#### WARRANTY

**NOTE** To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100  $\text{m}^2,$  applied every 4 to 8 weeks during the local growing season.

CANADA

#### Application Rates:

To apply at the rate of 0.43 kg N/100 sq. meters. Equivalent to 1.8 kg of product per 100 sq. meters.

To apply at the rate of 0.22 kg N/100 sq. meters. Equivalent to 1.0 kg of product per 100 sq. meters.

SUGGESTED SPREADER SETTINGS			
Spreader 0.43	kg N/100 m <sup>2</sup> .	0.22 kg N/100 m <sup>2</sup> .	
Lebanon Turf	41/2	4	
Andersons AccuPro	L	Н	
Earthway Rotary	15	14	
Gandy	27	24	
Lely	5½ II	4 II	
Lesco (letter/numerio	i) i/19	G/15	
PennMulch HVO	M	J	
ProScape SS	M	J	
Spyker	41/2	4	
Vicon	25	22	

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



F699

22-29163

Application Rates:

To feed at the rate of 0.9lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 3.7 lbs per 1,000 sq. ft. or 160.8 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.1 lbs per 1,000 sq. ft. or 90.6 lbs. per acre.

SUGGESTED SPREADER SETTINGS				
Spreader 0.9#	N/1,000 sq. ft.	0.5#N/1,000 sq. ft.		
Lebanon Turf	41/2	4		
Andersons AccuPro	L	H		
Earthway Rotary	15	14		
Gandy	27	24		
Lely	5½ II	4 II		
Lesco (letter/numerio	c) i/19	G/15		
PennMulch HVO	M	J		
ProScape SS	M	J		
Spyker	41/2	4		
Vicon	25	22		
These settings were calibrated and field tested				

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com



For technical assistance or more information about our products visit www.LebanonTurf.com

## Lebanon Seaboard Corporation

1600 E. Cumberland St • Lebanon, PA 17042 800-233-0628 • (717) 273-1685

50 Lbs. (22.7 kg) Net Weight