

LebanonTurf



Country Club®

Blended Fertilizer

FERTILIZER



21-0-10 MESA

P A T E N T E D

GUARANTEED ANALYSIS

Total Nitrogen (N)	21%
4.0% Ammoniacal Nitrogen	
4.5% Water Insoluble Nitrogen*	
4.9% Urea Nitrogen	
7.6% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ O)	10%
Sulfur (S)	8%
8.0% Combined Sulfur (S)	

Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Sulfate of Potash.

*12.1% Slowly available Nitrogen from Methylene Ureas.

Chlorine (Cl) not more than	2.0%
	U.S.A

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)	21%
4.0% Ammoniacal Nitrogen	
4.5% Water Insoluble Nitrogen*	
4.9% Urea Nitrogen	
7.6% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ O)	10%
Sulfur (S)	8%
8.0% Combined Sulfur (S)	

Chlorine (Cl) (maximum).....	2.0%
------------------------------	------

Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Sulfate of Potash.

*12.1% Slowly available Nitrogen from Methylene Ureas.

CANADA

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Warning



Hazard Statements

Causes mild skin irritation.
May cause eye irritation on contact.
May be harmful if inhaled repeatedly over long time periods.
Suspected of causing cancer by prolonged/repeated inhalation.

Precautionary Statements

Avoid breathing dust.
Use appropriate personal protective equipment as required to avoid breathing dust.
If exposed or concerned, seek medical advice.
If skin irritation occurs, wash with plenty of soap and water.
Wash hands and exposed skin thoroughly after handling.
If in eyes, rinse cautiously with water for several minutes.
If eye irritation persists: seek medical attention.
Dispose in accordance with all federal, state and local regulations.

KEEP OUT OF REACH OF CHILDREN.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Information regarding the contents and levels of metals in this product is available on the Internet at:
<http://www.regulatory-info-lebsea.com>

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:
To feed at the rate of 0.9lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 4.2 lbs per 1,000 sq. ft. or 174 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.4 lbs per 1,000 sq. ft. or 120 lbs. per acre.


SUGGESTED SPREADER SETTINGS			
Spreader	0.9#N/1,000 sq. ft.	0.5#N/1,000 sq. ft.*	
Lebanon Turf	4½	4	
Andersons AccuPro	M	H	
Earthway Rotary	18	15	
Gandy	30	24	
Lely	6 II	4 II	
Lesco (letter/numeric)	K/22	H/16	
PennMulch HVO	N	J	
ProScape SS	N	J	
Spyker	4½	4	
Vicon	30	23	

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Application Rates:
To apply at the rate of 0.43 kg N/100 sq. meters. Equivalent to 2.0 kg of product per 100 sq. meters.
To apply at the rate of 0.22 kg N/100 sq. meters. Equivalent to 1.1 kg of product per 100 sq. meters.

SUGGESTED SPREADER SETTINGS			
Spreader	0.43 kg N/100 m ² .	0.22 kg N/100 m ² .	
Lebanon Turf	4½	4	
Andersons AccuPro	M	H	
Earthway Rotary	18	15	
Gandy	30	24	
Lely	6 II	4 II	
Lesco (letter/numeric)	K/22	H/16	
PennMulch HVO	N	J	
ProScape SS	N	J	
Spyker	4½	4	
Vicon	30	23	

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



F699 **22-35756** j/16

MESA®, PennMulch® and ProScape® are registered trademarks of Lebanon Seaboard Corporation.

LebanonTurf

For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation
1600 E. Cumberland St • Lebanon, PA 17042
800-233-0628 • (717) 273-1685

50 Lbs. (22.7 kg) Net Weight



FERTILIZER

21-0-10

MESA

P A T E N T E D

GUARANTEED ANALYSIS

Total Nitrogen (N)	21%
4.0% Ammoniacal Nitrogen	
4.5% Water Insoluble Nitrogen*	
4.9% Urea Nitrogen	
7.6% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ O)	10%
Sulfur (S)	8%
8.0% Combined Sulfur (S)	

Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Sulfate of Potash.

*12.1% Slowly available Nitrogen from Methylene Ureas.

Chlorine (Cl) not more than 2.0%
U.S.A

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)	21%
4.0% Ammoniacal Nitrogen	
4.5% Water Insoluble Nitrogen*	
4.9% Urea Nitrogen	
7.6% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ O)	10%
Sulfur (S)	8%
8.0% Combined Sulfur (S)	

Chlorine (Cl) (maximum)..... 2.0%

Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Sulfate of Potash.

*12.1% Slowly available Nitrogen from Methylene Ureas.

CANADA

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Warning



Hazard Statements

- Causes mild skin irritation.
- May cause eye irritation on contact.
- May be harmful if inhaled repeatedly over long time periods.
- Suspected of causing cancer by prolonged/repeated inhalation.

Precautionary Statements

- Avoid breathing dust.
- Use appropriate personal protective equipment as required to avoid breathing dust.
- If exposed or concerned, seek medical advice.
- If skin irritation occurs, wash with plenty of soap and water.
- Wash hands and exposed skin thoroughly after handling.
- If in eyes, rinse cautiously with water for several minutes.
- If eye irritation persists: seek medical attention.
- Dispose in accordance with all federal, state and local regulations.

KEEP OUT OF REACH OF CHILDREN.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Information regarding the contents and levels of metals in this product is available on the Internet at:
<http://www.regulatory-info-lebsea.com>

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:

To feed at the rate of 0.9lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 4.2 lbs per 1,000 sq. ft. or 174 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.4 lbs per 1,000 sq. ft. or 120 lbs. per acre.

SUGGESTED SPREADER SETTINGS

Spreader	0.9#N/1,000 sq. ft.	0.5#N/1,000 sq. ft.
Lebanon Turf	4½	4
Andersons AccuPro	M	H
Earthway Rotary	18	15
Gandy	30	24
Lely	6 II	4 II
Lesco (letter/numeric)	K/22	H/16
PennMulch HVO	N	J
ProScape SS	N	J
Spyker	4½	4
Vicon	30	23

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Application Rates:


To apply at the rate of 0.43 kg N/100 sq. meters. Equivalent to 2.0 kg of product per 100 sq. meters.

To apply at the rate of 0.22 kg N/100 sq. meters. Equivalent to 1.1 kg of product per 100 sq. meters.

SUGGESTED SPREADER SETTINGS

Spreader	0.43 kg N/100 m ² .	0.22 kg N/100 m ² .
Lebanon Turf	4½	4
Andersons AccuPro	M	H
Earthway Rotary	18	15
Gandy	30	24
Lely	6 II	4 II
Lesco (letter/numeric)	K/22	H/16
PennMulch HVO	N	J
ProScape SS	N	J
Spyker	4½	4
Vicon	30	23

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



F699

22-35756

j/16

FERTILIZER

29-0-10

MESA

P A T E N T E D



GUARANTEED ANALYSIS

Total Nitrogen (N)	29%
3.5% Ammoniacal Nitrogen	
2.2% Water Insoluble Nitrogen*	
19.3% Urea Nitrogen	
4.0% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ O).....	12%
Sulfur (S)	4.2%
5.7% Combined Sulfur (S)	

Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash.

*6.2% Slowly Available Nitrogen from Methylene Ureas.

Chlorine (Cl) not more than

U.S.A

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)	29%
3.5% Ammoniacal Nitrogen	
2.2% Water Insoluble Nitrogen*	
19.3% Urea Nitrogen	
4.0% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ O).....	12%
Sulfur (S)	4.2%
5.7% Combined Sulfur (S)	

Chlorine (Cl) maximum

9.0%

Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash.

*6.2% Slowly Available Nitrogen from Methylene Ureas.

CANADA

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.



Warning

Hazard Statements

May be harmful if inhaled repeatedly over prolonged periods.
Suspected of causing cancer by prolonged/repeated inhalation.
Causes mild skin irritation.

May cause eye irritation on contact

Precautionary Statements

Avoid breathing dust.
Use appropriate personal protective equipment as required to avoid breathing dust.

If skin irritation occurs: Wash with soap and water.

Wash hands and exposed skin thoroughly after handling.

If in eyes, rinse cautiously with water for several minutes. If eye irritation persists: seek medical attention.

Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

KEEP OUT OF REACH OF CHILDREN.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:

To feed at the rate of 0.43 kg Nitrogen (N)
Equivalent to 1.5 kg of product per 100 sq. meters.

To feed at the rate of 0.22 kg Nitrogen (N)
Equivalent to 0.9 kg of product per 100 sq. meters.

SUGGESTED SPREADER SETTINGS

Spreader	0.43# N/100 m ²	0.22# N/100 m ²
Lebanon Turf	4½	3½
Andersons AccuPro	M	H
Earthway Rotary	15	14
Gandy	30	24
Lely	6½ II	4 II
Lesco (letter/numeric)	J/21	G/15
PennMulch HVO	L	i
ProScape SS	L	i
Spyker	4½	3½
Vicon	26	22

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Application Rates:

To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 3.1 lbs per 1,000 sq. ft. or 135 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 1.7 lbs per 1,000 sq. ft. or 74 lbs. per acre.

SUGGESTED SPREADER SETTINGS

Spreader	0.9# /1,000 sq. ft.	0.5# /1,000 sq. ft.
Lebanon Turf	4½	3½
Andersons AccuPro	M	H
Earthway Rotary	15	14
Gandy	30	24
Lely	6½ II	4 II
Lesco (letter/numeric)	J/21	G/15
PennMulch HVO	L	i
ProScape SS	L	i
Spyker	4½	3½
Vicon	26	22

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



F699

22-29084

i/16

Information regarding the contents and levels of metals in this product is available on the Internet at:
<http://www.regulatory-info-lebsea.com>