

LebanonTurf

Country Club[®]

Blended Fertilizer

20-0-10



WITH **expo**[®] WITH **MESA**
 POTASSIUM SULFATE

GUARANTEED ANALYSIS

Total Nitrogen (N)	20%
2.3% Ammoniacal Nitrogen	
2.5% Urea Nitrogen	
5.6% Water Insoluble Nitrogen*	
9.6% Other Water Soluble Nitrogen*	
Soluble Potash (K₂O)	10%
Sulfur (S)	3.0%
3.0% Combined Sulfur (S)	

Derived From: Methylene Urea, potassium sulfate, ammonium sulfate.
 *15.2% Slowly available Nitrogen from Methylene Ureas

U.S.A

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)	20%
2.3% Ammoniacal Nitrogen	
2.5% Urea Nitrogen	
5.6% Water Insoluble Nitrogen*	
9.6% Other Water Soluble Nitrogen*	
Soluble Potash (K₂O)	10%
Sulfur (S)	3.0%
3.0% Combined Sulfur (S)	

Derived From: Methylene Urea, potassium sulfate, ammonium sulfate.
 *15.2% Slowly available Nitrogen from Methylene Ureas

CANADA

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Warning



Hazard Statements

Causes skin irritation.
 May be harmful if inhaled.
 Causes serious eye irritation.
 Causes eye irritation.

Precautionary Statements

Wash hands and exposed skin thoroughly after handling.
 Wear protective gloves/protective clothing/eye protection/face protection.
 If in eyes, rinse with water and seek medical attention if symptoms persist.
 Take off contaminated clothing and wash before reuse.
 IF ON SKIN: Wash with plenty of soap and water.
 IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell.
 IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
 If skin irritation occurs: Get medical advice/ attention.
 If eye irritation persists: Get medical advice/attention.
 Keep out of reach of children
 Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

Do not apply to frozen ground.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:
 To feed at the rate of 0.43 kg Nitrogen (N)
 Equivalent to 2.2 kg of product per 100 sq. meters.

To feed at the rate of 0.22 kg Nitrogen (N)
 Equivalent to 1.2 kg of product per 100 sq. meters.

SUGGESTED SPREADER SETTINGS		
Spreader	0.43 kg N/100 m ² .	0.22 kg N/100 m ² .
Lebanon Turf	4 1/4	3
Andersons AccuPro	J	G
Earthway Rotary	15	13
Gandy	25	23
Lely	5 1/2 II	3 1/2 II
Lesco (letter/numeric)	G/16	F/12
PennMulch HVO	K	G
ProScape SS	K	G
Spyker	4 1/2	3
Vicon	23	21

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Application Rates:
 To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 4.5 lbs per 1,000 sq. ft. or 197.6 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.5 lbs per 1,000 sq. ft. or 108.7 lbs. per acre.

SUGGESTED SPREADER SETTINGS		
Spreader	0.9# N/1,000 sq. ft.	0.5# N/1,000 sq. ft.
Lebanon Turf	4 1/4	3
Andersons AccuPro	J	G
Earthway Rotary	15	13
Gandy	25	23
Lely	5 1/2 II	3 1/2 II
Lesco (letter/numeric)	G/16	F/12
PennMulch HVO	K	G
ProScape SS	K	G
Spyker	4 1/2	3
Vicon	23	21

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



0 88685 37116 5
 F699 **22-37116** L/21

Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.regulatory-info-lebsea.com>

LebanonTurf

For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation
 1600 E. Cumberland St • Lebanon, PA 17042
 800-233-0628 • (717) 273-1685

50 Lbs. (22.7 kg) Net Weight

20-0-10



WITH **expo**®
POTASSIUM
SULFATE

WITH **MESA**®

GUARANTEED ANALYSIS

Total Nitrogen (N)	20%
2.3% Ammoniacal Nitrogen	
2.5% Urea Nitrogen	
5.6% Water Insoluble Nitrogen*	
9.6% Other Water Soluble Nitrogen*	
Soluble Potash (K₂O)	10%
Sulfur (S)	3.0%
3.0% Combined Sulfur (S)	

Derived From: Methylene Urea, potassium sulfate, ammonium sulfate.
*15.2% Slowly available Nitrogen from Methylene Ureas

U.S.A

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)	20%
2.3% Ammoniacal Nitrogen	
2.5% Urea Nitrogen	
5.6% Water Insoluble Nitrogen*	
9.6% Other Water Soluble Nitrogen*	
Soluble Potash (K₂O)	10%
Sulfur (S)	3.0%
3.0% Combined Sulfur (S)	

Derived From: Methylene Urea, potassium sulfate, ammonium sulfate.
*15.2% Slowly available Nitrogen from Methylene Ureas

CANADA

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Warning



Hazard Statements

Causes skin irritation.
May be harmful if inhaled.
Causes serious eye irritation.
Causes eye irritation.

Precautionary Statements

Wash hands and exposed skin thoroughly after handling.
Wear protective gloves/protective clothing/eye protection/face protection.
If in eyes, rinse with water and seek medical attention if symptoms persist.
Take off contaminated clothing and wash before reuse.
IF ON SKIN: Wash with plenty of soap and water.
IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
If skin irritation occurs: Get medical advice/ attention.
If eye irritation persists: Get medical advice/attention.
Keep out of reach of children
Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

Do not apply to frozen ground.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:

To feed at the rate of 0.43 kg Nitrogen (N)
Equivalent to 2.2 kg of product per 100 sq. meters.

To feed at the rate of 0.22 kg Nitrogen (N)
Equivalent to 1.2 kg of product per 100 sq. meters.

SUGGESTED SPREADER SETTINGS

Spreader	0.43 kg N /100 m ² .	0.22 kg N /100 m ² .
Lebanon Turf	4¼	3
Andersons AccuPro	J	G
Earthway Rotary	15	13
Gandy	25	23
Lely	5½ II	3½½ II
Lesco (letter/numeric)	G/16	F/12
PennMulch HVO	K	G
ProScape SS	K	G
Spyker	4½	3
Vicon	23	21

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Application Rates:

To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 4.5 lbs per 1,000 sq. ft. or 197.6 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.5 lbs per 1,000 sq. ft. or 108.7 lbs. per acre.

SUGGESTED SPREADER SETTINGS

Spreader	0.9# N /1,000 sq. ft.	0.5# N /1,000 sq. ft.
Lebanon Turf	4¼	3
Andersons AccuPro	J	G
Earthway Rotary	15	13
Gandy	25	23
Lely	5½ II	3½½ II
Lesco (letter/numeric)	G/16	F/12
PennMulch HVO	K	G
ProScape SS	K	G
Spyker	4½	3
Vicon	23	21

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



F699

22-37116

L/21

Information regarding the contents and levels of metals in this product is available on the Internet at:
<http://www.regulatory-info-lebsea.com>