LebanonTurf

Country Club® Blended Fertilizer

4-0-14 IMIC * AACCA



GUARANTEED ANALYSIS

1.9% Ammoniacal Nitrogen 2.5% Water Insoluble Nitrogen* 5.2% Urea Nitrogen 4.4% Other Water Soluble Nitrogen * Soluble Potash (K₂0) 14% Sulfur (S) 6.7% Combined Sulfur (S) Derived From: methylene ureas, ammonium sulfate, sulfate of potash, urea. *6.9% Slowly Available Nitrogen from Methylene Ureas. NON PLANT FOOD/OTHER INGREDIENTS

Humic acids (derived from Leonardite). PLANT FOOD INGREDIENTS Plant Food Fertilizer...

Product contains humic acids which may aid in nutrient availability.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Lbs. K//1,000 Sq. Ft. Lbs Humic Acids//1,000 Sq. Ft.

Lbs N/1,000 Sq. Ft. Lbs Fert//1,000 Sq. Ft. Lbs Required for One Acre Sq. Ft. Coverage per bag 7.800

Application Rates:

To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq ft., apply this product at 6.4 lbs per 1,000 sq. ft. or 279 lbs. per acre. To feed at the rate of $lac{1}{2}$ lb. Nitrogen (N) pe 1,000 sq. ft., apply this product at 3.6 lbs per 1,000 sq.

rt. or 157 ibs. per acre.			
SUGGESTED SPREADER SETTINGS			
Spreader 0.	9# N/1,000 sq. ft.	0.5 # N/1,000 sq. ft.	
Lebanon Turf	43/4	4	
Andersons AccuPro M		i	
Earthway Rotary	16	141/2	
Gandy	28	24	
Lely	61/2 II	4½ II	
Lesco (letter/numeric) J/21		H/15	
PennMulch HVO	M	J	
ProScape SS	M	J	
Spyker	43/4	4	
Vicon	26	22	
These settings were calibrated and field tested.			
However, age and condition of spreader, speed of			

operation and evenness of terrain may require slightly

different settings for desired coverage.

Hazard Statements Causes skin irritation

May be harmful if inhaled



Causes eye irritation **Precautionary Statements**

Wash hands and exposed skin thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection. If in eyes, rinse with water and seek medical attention if symptoms persist. Take off contaminated clothing and wash before reuse. IF ON SKIN: Wash with plenty of soap and water.

IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell. If skin irritation occurs: Get medical advice/ attention. If eye irritation persists: Get medical advice/attention. Keep out of reach of children

Precautionary Statements for disposal Dispose in accordance with all federal, state and local regulations

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommenda

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn. Do not apply to frozen ground.

WARRANTY

tions in your approved nutrient management plan.

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

PennMulch® and ProScape® are registered trademarks of Lebanon Seaboard Corporation. Country Club and the Lebanon Diamond are registered trademarks of Lebanon Seaboard Corporation.

GUARANTEED MINIMUM ANALYSIS

1.9% Ammoniacal Nitrogen

2.5% Water Insoluble Nitrogen* 5.2% Urea Nitrogen

4.4% Other Water Soluble Nitrogen *

Soluble Potash (K₂0) 14%

6.7% Combined Sulfur (S)

Derived From: methylene ureas, ammonium sulfate, sulfate of potash, urea.

*6.9% Slowly Available Nitrogen from Methylene Ureas.

NON PLANT FOOD/OTHER INGREDIENTS Humic acids (derived from Leonardite)..

PLANT FOOD INGREDIENTS

Plant Food Fertilizer.... 92.2% CANADA

Application Rates:

Application Rates and Frequencies in Canada: Follow your or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Equivalent to 3 kg of product per 100 sq. meters.

To feed at the rate of 0.43 kg Nitrogen (N)

To feed at the rate of 0.22 kg Nitrogen (N)

Equivalent to 1.8 kg of product per 100 sq. meters.

_qa::a::a::a::a::a::a::a::a::a::a::a::a::	rio ng er premaer pr	
SU	IGGESTED SPREADE	R SETTINGS
Spreader	0.43 kg N /100 m ²	0.22 kg N /100 m ²
Lebanon Turf	43/4	4
Andersons AccuPro M		i
Earthway Rotar	y 16	141/2
Gandy	28	24
Lely	6½ II	4½ II
Lesco (letter/numeric)J/21		H/15
PennMulch HVC	M	J
ProScape SS	M	J
Spyker	43/4	4
Vicon	26	22
These settings	were calibrated a	nd field tested.

However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage



Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com 800-233-0628 · (717) 273-1685



For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation

1600 E. Cumberland St • Lebanon, PA 17042 800-233-0628 • (717) 273-1685

50 Lbs. (22.7 kg) Net Weight