LebanonTurf

Country Club® Blended Fertilizer



22-0-12 (SON 15)



CHADANTEED ANALYSIS

Total Nitrogen (N)	
1.0% Ammoniacal Nitrogen	
8.7% Water Insoluble Nitrogen*	
7.3% Urea Nitrogen	
5.0% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ 0)	
Sulfur (S)	
5.5% Combined Sulfur (S)	

Derived From: Methylene Ureas, Urea, Ammonium Sulfate, Sulfate of Potash.

*13.7% Slowly available nitrogen from Methylene Ureas.

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)	22%
1.0% Ammoniacal Nitrogen	
8.7% Water Insoluble Nitrogen*	
7.3% Urea Nitrogen	
5.0% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ 0)	12%
Sulfur (S)	5.5%
5.5% Combined Sulfur (S)	

Derived From: Methylene Ureas, Urea, Ammonium Sulfate, Sulfate of Potash.

*13.7% Slowly available nitrogen from Methylene Ureas.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Application Rates:

To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 4.1 lbs per 1,000 sq. ft. or 178 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.3 lbs per 1,000 sq. ft. or 100 lbs. per acre.

ins her 1,000 sq. it. or 100 ins. her acie.				
SUGGESTED SPREADER SETTINGS				
Spreader 0.9#N	/1,000 sq. ft.	0.5#N/1,000 sq. f		
Lebanon Turf	41/2	4		
Andersons AccuPro	L	Н		
Earthway Rotary	15	14		
Gandy	27	24		
Lely	51∕2 II	4 II		
Lesco (letter/numeric)	I/19	G/15		
PennMulch HVO	M	J		
ProScape SS	M	J		
Spyker	41/2	4		
Vicon	25	22		
These settings were	calibrated a	nd field tested		

In Pennsylvania: Do not exceed 0.9 lbs of total nitrogen per 1,000 sq. ft. per application.

different settings for desired coverage.

However, age and condition of spreader, speed of operation and evenness of terrain may require slightly

Information regarding the contents and levels of metals in

this product is available on the Internet at: http://www.regulatory-info-lebsea.com

Hazard Statements

Causes skin irritation May be harmful if inhaled. Causes serious eve irritation. Causes eye irritation.

U.S.A

Precautionary Statements

Wash hands and exposed skin thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection. If in eyes, rinse with water and seek medical attention if symptoms persist. Take off contaminated clothing and wash before reuse IF ON SKIN: Wash with plenty of soap and water. IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If skin irritation occurs: Get medical advice/ attention If eye irritation persists: Get medical advice/attention.

Keep out of reach of children Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

Do not apply to frozen ground

WARRANTY

NOTE: To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

CANADA

Application Rates:

To feed at the rate of 0.41 kg Nitrogen (N) per 1,000 sq. ft., apply this product at 1.86 kg per 1,000 m² or $59\ kg$ per hectare. To feed at the rate of 0.22 kg Nitrogen (N) per 1,000 sq. ft., apply this product at 1.04 kg per 1,000 m² or 45 kg per hectare.

SUGGEST	ED SPREADE	R SETTINGS
Spreader 0.43 k	g #N/100 m².	0.22 kg #N/100 m ² .
Lebanon Turf	41/2	4
Andersons AccuPro	L	Н
Earthway Rotary	15	14
Gandy	27	24
Lely	5½ II	4 II
Lesco (letter/numeric)	I/19	G/15
PennMulch HVO	M	J
ProScape SS	M	J
Spyker	41/2	4
Vicon	25	22

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

LebanonTurf

For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation

1600 E. Cumberland St • Lebanon, PA 17042 800-233-0628 • (717) 273-1685

50 Lbs. (22.7 kg) Net Weight