

BENFLURALIN	GROUP 3	HERBICIDE
TRIFLURALIN	GROUP 3	HERBICIDE



FERTILIZER WITH TEAM[®] HERBICIDE

CRABGRASS CONTROL

15-0-5

CONTAINS SCU

GUARANTEED ANALYSIS

Total Nitrogen (N).....	15%
15.0% Urea Nitrogen*	
Soluble Potash (K ₂ O)	5%
Sulfur (S)	1.5%
1.5% Free Sulfur (S)	



Derived From: Sulfur Coated Urea, Urea, Muriate of Potash.

Chlorine (Cl) not more than 5.0%
*4.5% Slowly available Nitrogen from Sulfur Coated Urea.

SUGGESTED SPREADER SETTINGS					
Spreader	4 lbs./1,000 sq. ft.	6 lbs./1,000 sq. ft.	Spreader	4lbs./1,000 sq. ft.	6lbs./1,000 sq. ft.
LebanonTurf	4	5½	Lesco (letter/Numeric)	J/21	L/24
Andersons AccuPro	M	O	PennMulch HVO	N	P
Earthway Rotary	16	20	ProScape SS	N	P
Gandy	28	30	Spyker	4	5½
Lely	6 II	7 II	Vicon (all models)	26	31

Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.regulatory-info-lebsea.com>

A Granular Fertilizer Containing TEAM for the Preemergence Control of Crabgrass, Foxtail, Goosegrass, *Poa Annua* and Other Annual Weed Grasses in Established Turfgrass.

ACTIVE INGREDIENTS:

Benfluralin	0.770%
Trifluralin	0.385%

OTHER INGREDIENTS:..... 98.845%

TOTAL: 100.000%

Benfluralin CAS. No. 1861-40-1 Trifluralin CAS. No. 1582-09-8 (L) (D)

EPA Reg. No. 961-346 EPA Est. No. 961-PA-01;961-IL-01

TEAM is a Trademark of Gowan Company, LLC. Superscript used is first letter of lot number.

KEEP OUT OF REACH OF CHILDREN

CAUTION

NOTICE: Read the entire label. Use only according to label directions. Before buying this product, read the Warranty Section elsewhere on this label. If terms are not acceptable, return the unopened container at once.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Socks plus shoes, and
- Chemical-resistant gloves made of any waterproof material.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS
<ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID	
IF ON SKIN: OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Emergency Medical Information, call (888) 208-1368.	

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. **DO NOT** apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize drift.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Application Restrictions: **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during applications.

Entry Restrictions: This product must be watered in as soon as possible after application. **DO NOT** enter or allow other people (or pets) to enter the treated area until dusts have settled. Except for those people involved in the watering-in, **DO NOT** enter or allow other people (or pets) to enter the treated areas until the watering-in is complete and the surface is dry.

WEED RESISTANCE MANAGEMENT

For resistance management, **Lebanon Pro Brand fertilizer with TEAM HERBICIDE Crabgrass Control** is a Group 3 herbicide. Any weed population may contain certain or develop plants naturally resistant to **Lebanon Pro Brand fertilizer with TEAM HERBICIDE Crabgrass Control** and other Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps: Rotate the use of **Lebanon Pro Brand fertilizer with TEAM HERBICIDE Crabgrass Control** or other Group 3 herbicides within a growing season with different herbicide groups that control the same weeds.

Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include:(1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resis-tance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment. If weed population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.

Contact your sales representative, pest control advisors, or local extension specialist for additional pesticide resistance management and/or integrated weed management recommendations for specific types of plants and weed biotypes.

PRODUCT INFORMATION

This turf fertilizer product contains TEAM, a preemergence herbicide which is applied to established turfgrasses for the control of the winter annual grass: Annual Bluegrass (*Poa annua*); and summer annual grasses: Smooth and Hairy Crabgrass, Goosegrass (silver crabgrass or crowfootgrass), Barnyardgrass (watergrass) and Green and Yellow Foxtail. This product also provides partial control of prostrate spurge (*Euphorbia humistrata*), spotted spurge (*Euphorbia maculata*), creeping spurge (*Euphorbia serpens*), creeping woodsorrel (*Oxalis corniculata*) and yellow woodsorrel (*Oxalis stricta*).

This Product Is For Use On The Following Established Warm and Cool Season Perennial Turfgrass Species:

Established Cool Season Turfgrasses - Bentgrass Fairways, Kentucky Bluegrass, Perennial Ryegrass, Fescues

Established Warm Season Turfgrasses - Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass, Zoysiagrass

This product should be applied one to two weeks prior to the anticipated germination of the target annual weeds. Apply using a properly calibrated granular fertilizer spreader which will uniformly distribute the product over the target area. Avoid skips and excessive overlaps as poor weed control or turfgrass injury may occur. Optimum weed control performance will be obtained if treated areas are irrigated soon after application. This product will not harm most nearby ornamentals.

In areas with extended growing seasons or where desired, a second application at the recommended rate can be made 8 to 10 weeks following the initial application on cool season turfgrasses and 10 to 12 weeks following the initial application on warm season turfgrasses.

PRECAUTIONS AND RESTRICTIONS

DO NOT Ship or Store with Food, Feeds, Drugs or Clothing.

DO NOT aerially apply this product.

Reseeding: Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. **DO NOT** reseed treated areas for a minimum of 8 weeks after a single application of 4 lbs. per 1,000 sq. ft. (174 lbs. per acre) or 12 to 16 weeks after a single application of 6 lbs. per 1,000 sq. ft. (260 lbs. per acre). For split applications, **DO NOT** reseed treated areas for 8 to 10 weeks after the last 3 lbs. per 1,000 sq. ft. rate (130 lbs. per acre) or 10 to 12 weeks after the last 4 lbs. per 1,000 sq. ft. rate (174 lbs. per acre).

Use caution to confine product to intended application area, as the active ingredients may stain concrete, plastic or other materials.

DO NOT apply this product to golf course putting greens, dichondra, or newly sprigged areas of Bermudagrass, St. Augustinegrass, Centipedegrass, or Zoysiagrass.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application.

APPLICATION INSTRUCTIONS		
Broadcast Application Rates:	Cool Season Turfgrasses	Warm Season Turfgrasses
Summer Annual Grass Weeds (Spring Application)	130 to 195 lbs./acre 3 to 4.5 lbs./1,000 sq. ft.	260 lbs./acre 6 lbs./1,000 sq. ft.
Winter Annual Grass Weeds (Fall Application)	130 to 174 lbs./acre 3 to 4 lbs./1,000 sq. ft.	260 lbs./acre 6 lbs./1,000 sq. ft.

Split Applications for Extended Control and Lawn Care Operator Programs:

As an alternative to single application programs, this product may also be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time.

Apply this product at 3 lbs. plus 3 lbs. per 1,000 sq. ft. (130 lbs. plus 130 lbs. per acre) for Cool Season Turfgrasses and in the northern half of the Warm Season Turfgrass area of the U.S. or 4 lbs. plus 4 lbs. per 1,000 sq. ft. (174 lbs. plus 174 lbs. per acre) in the southern half of the Warm Season Turfgrasses area of the U.S. The second application should be made 8 to 10 weeks following the 3 lbs. per 1,000 sq. ft. rate (130 lbs. per acre) and 10 to 12 weeks following the 4 lbs. per 1,000 sq. ft. rate (174 lbs. per acre).

Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

For the active ingredient benfluralin, the following rate restrictions apply:

Cool Season Turf:

- Maximum of 1.5 lbs. a.i. per acre per application.
- Maximum of 2 applications per year.
- Maximum of 3 lbs. a.i. per acre per year.

Warm Season Turf:

- Maximum of 3 lbs. a.i. per acre per application.
- Maximum of 2 applications per year.
- Maximum of 6 lbs. a.i. per acre per year.

This product must be soil-incorporated within 24 hours of application either by at least 0.5 acre inch of water (rainfall, irrigation) or mechanical incorporation. If a second incorporation is needed it should be delayed 3 to 5 days after the first incorporation and be completed prior to planting.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage and disposal.

Pesticide Storage: Store product in its original container in a cool, dry, locked place out of the reach of children and animals.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

WARRANTY

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Lebanon Seaboard Corporation. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, LEBANON SEABOARD CORPORATION MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of LEBANON SEABOARD CORPORATION is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, LEBANON SEABOARD CORPORATION DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT LEBANON SEABOARD CORPORATION'S ELECTION, THE REPLACEMENT OF PRODUCT.

MANUFACTURED BY:

LEBANON SEABOARD CORPORATION

1600 EAST CUMBERLAND STREET

LEBANON, PA 17042

rev: EH/e23

For technical assistance or more information about our products visit: www.LebanonTurf.com

Net Weight: 50 LBS. (22.7 kg)