


# Lebanon Pro



**BRAND**

Fertilizer with  
**TEAM<sup>®</sup> PRO 0.86%**  
PREEMERGENCE CRABGRASS CONTROL  
**22-0-6**  
CONTAINS **SCU**



0 88685 40417 7

22-40417

**GUARANTEED ANALYSIS**

Total Nitrogen ..... 22%

22.0% Urea Nitrogen\*

Soluble Potash (K<sub>2</sub>O) ..... 6%

Sulfur (S) ..... 1.2%

1.2% Free Sulfur (S)

Derived from: Sulfur Coated Urea, Urea, Muriate of Potash.

Chlorine (Cl) not more than ..... 6.0%

\*Contains 8.8% Slowly Available Nitrogen from Sulfur Coated Urea . F699

Information regarding the contents and levels of metals in this product is available on the Internet at:  
(CA) <https://apps4.cdfa.ca.gov/FertilizerProducts/>  
(OR) [http://oda.state.or.us/dbs/heavy\\_metal/search.lasso](http://oda.state.or.us/dbs/heavy_metal/search.lasso)  
(WA) <http://agr.wa.gov/pestfert/fertilizers/productdatabase.aspx>

Spreader Settings as a Guide for Calibration				
Brand and Model of Spreader	174#/acre	233#/acre	Spreader	174#/acre 233#/acre
LebanonTurf	4%	5%	Lesco (letter/Numeric)	J/21 K/22
Andersons AccuPro	M	0	PennMulch HVO	M
Earthway Rotary	16	19	ProScape SS	M
Gandy	28	31	Spyker	4% 5%
Lely	6 II	6½ II	Vicon (all models)	26 30

**A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, Poa annua and other annual grasses and broadleaf weeds in established turfgrass and landscape ornamentals.**

**ACTIVE INGREDIENTS:**  
benefin: *N*-butyl-*N*-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-*p*-toluidine . . 0.43%  
trifluralin:  $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-*N,N*-dipropyl-*p*-toluidine . . . 0.43%  
**OTHER INGREDIENTS** . . . . . 99.14%  
**TOTAL** . . . . . 100.00%

EPA Reg. No. 10163-346-961  
Team<sup>®</sup> is a registered trademark of Gowan Company, LLC

(L) (D)  
EPA Est. 961-PA-01:961-IL-01  
Superscript used is first letter of lot number.

**KEEP OUT OF REACH OF CHILDREN**  
**WARNING**  
**Precautionary Statements**  
**Hazards to Humans and Domestic Animals**

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In some Individuals.

Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Refer to back panel for additional precautionary information including Directions for Use.

**Notice: Read the entire label.** Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-424-9300.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

**Distributed by:**  
**Lebanon Seaboard Corporation**  
**1600 E. Cumberland St. • Lebanon, PA 17042**  
**800-233-0628 • (717) 273-1685**

**NET WT. 50 LBS. (22.7 kg)**

# Fertilizer with TEAM<sup>®</sup> PRO 0.86%

## PREEMERGENCE CRABGRASS CONTROL

# 22-0-6



## CONTAINS SCU

### GUARANTEED ANALYSIS

Total Nitrogen ..... 22%

22.0% Urea Nitrogen\*

Soluble Potash (K<sub>2</sub>O) ..... 6%

Sulfur (S) ..... 1.2%

1.2% Free Sulfur (S)

Derived from: Sulfur Coated Urea, Urea, Muriate of Potash.

Chlorine (Cl) not more than ..... 6.0%

\*Contains 8.8% Slowly Available Nitrogen from Sulfur Coated Urea.

F699



**22-40417**

Information regarding the contents and levels of metals in this product is available on the Internet at:

(CA) <https://apps4.cdfa.ca.gov/FertilizerProducts/>

(OR) [http://oda.state.or.us/dbs/heavy\\_metal/search.lasso](http://oda.state.or.us/dbs/heavy_metal/search.lasso)

(WA) <http://agr.wa.gov/pestfert/fertilizers/productdatabase.aspx>

#### Spreader Settings as a Guide for Calibration

Brand and Model of Spreader	174#/acre	233#/acre	Spreader	174#/acre	233#/acre
LebanonTurf	4¾	5¼	Lesco (letter/Numeric)	J/21	K/22
Andersons AccuPro	M	0	PennMulch HVO	M	0
Earthway Rotary	16	19	ProScape SS	M	0
Gandy	28	31	Spyker	4¾	5¼
Lely	6 II	6½ II	Vicon (all models)	26	30

**A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, Poa annua and other annual grasses and broadleaf weeds in established turfgrass and landscape ornamentals.**

### ACTIVE INGREDIENTS:

benefin: *N*-butyl-*N*-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-*p*-toluidine . . 0.43%

trifluralin:  $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-*N,N*-dipropyl-*p*-toluidine . . . 0.43%

**OTHER INGREDIENTS** . . . . . 99.14%

**TOTAL** . . . . . 100.00%

**KEEP OUT OF REACH OF CHILDREN**  
**WARNING**  
**Precautionary Statements**  
**Hazards to Humans and Domestic Animals**

**Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In some Individuals.**

**Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.**

**Refer to back panel for additional precautionary information including Directions for Use.**

**Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-424-9300.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.



**Distributed by:**

**Lebanon Seaboard Corporation**  
**1600 E. Cumberland St. • Lebanon, PA 17042**  
**800-233-0628 • (717) 273-1685**

**NET WT. 50 LBS. (22.7 kg)**

**Personal Protective Equipment (PPE)**  
**All loaders, applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material
- Goggles, faceshield or safety glasses

Follow manufacturer’s instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**  
Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**First Aid**

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.  
**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.  
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-888-478-0798 for emergency medical treatment information.

**Environmental Hazards**

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

**Directions for Use**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.  
Read all directions carefully before applying this product.

**Application Restrictions:** Do not apply this product in a way that will contact workers or other persons or pets, either directly or through drift. Only protected handlers may be in the area during application.

**Entry Restrictions:** Do not enter or allow people or pets to enter the treated area until dusts have settled. If soil incorporation is required after application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

**Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.**

**Storage and Disposal**

Do not contaminate water, food or feed by storage or disposal.  
**Pesticide Storage:** Store in original container only. In case of leak or spill, contain material and dispose as waste.  
**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.  
**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**General Information**

This product is a granular fertilizer containing Team®, a preemergence herbicide for application to established turfgrass and landscape ornamentals (trees, shrubs, groundcovers, and flowers) for control of annual grasses and broadleaf weeds.

**General Use Precautions and Restrictions**

- Do not apply this product to golf course putting greens.
- Do not apply to dichondra.
- Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.

**Re seeding:** Do not reseed treated areas for at least 8 weeks after a single application of 174 lb of product per acre, or 12 to 16 weeks after a single application of 349 lb of product per acre unless injury can be tolerated. For split applications, do not reseed treated areas for 8 to 10 weeks after the last 174 lb of product per acre rate is applied, or 10 to 12 weeks after the last 233 lb of product per acre rate is applied unless injury can be tolerated.

**This product is recommended for use on the following turfgrass species:**

Established Cool Season Turfgrass	Established Warm Season Turfgrass
bentgrass fairways	bahiagrass
fescues	Bermudagrass
Kentucky bluegrass	centipedegrass
perennial ryegrass	fescue, tall
	St. Augustinegrass
	zoysiagrass

This product may be used for annual weed control in landscape settings on the following established ornamental plant species:

Ornamental Trees		Ornamental Shrubs		Ornamental Groundcover Plantings		Roses and Other Established Flowers	
Common Name	Scientific Name	Common Name	Scientific Name	Common Name	Scientific Name	Common Name	Common Name
almond	Prunus dulcis	andromeda, Japanese	Pieris japonica	aaronsbeard	Hypericum calycinum	African daisy	marigold
apple, crabapple	Malus spp.	arborvitae, American	Thuja occidentalis	bellflower, adriatic	Campanula elatines	aster (perennial)	marigold, cape
apricot	Prunus armeniaca	azalea	Rhododendron spp.	bellflower, poscharsky	Campanula poscharskyana	balsam	morningglory
ash, white	Fraxinus americana	barberry, Japanese	Berberis thunbergii	ceanothus	Ceanothus spp.	blackeyed susan	nasturtium
bald cypress	Taxodium distichum	barberry, mento	Berberis mentorensis	coreopsis	Coreopsis spp.	calendula	nicotiana
birch, European white	Betula pendula	boxwood, common	Buxus sempervirens	cotoneaster	Cotoneaster spp.	carnation	petunia
blackgum	Nyssa sylvatica	boxwood, littleleaf	Buxus microphylla	coyote brush	Baccharis pilularis	centaurea, velvet	phlox
cherry	Prunus spp.	camellia, Japanese	Camellia japonica	crown vetch	Coronilla vana	chrysanthemum	pincushion flower
chestnut, Chinese	Castanea mollissima	camellia, sasanqua	Camellia sasanqua	daisy trailing African	Osteospermum fruticosum	coreopsis	poppy, California
cottonwood	Populus deltoides	cherrylaurel, American	Prunus caroliniana	fern, asparagus	Asparagus densiflorus	cornflower	portulaca
dogwood, flowering	Cornus florida	cinquefoil	Potentilla spp.	gazania	Gazania spp.	cosmos	rose
dogwood, kousa	Cornus kousa	cleyera, Japanese	Cleyera japonica	germander	Teucrium chamaedrys	dahlia	salvia
douglas fir	Pseudotsuga menziesii	cotoneaster, cranberry	Cotoneaster apiculata	ice plant, largeleaf	Carpobrotus edulis	dianthus	shasta daisy
fir, balsam	Abies balsamea	cotoneaster, zabel	Cotoneaster zabelii	ivy, Algerian	Hedera canariensis	dusty miller	snapdragon
hemlock, Canada	Tsuga canadensis	deutzia	Deutzia spp.	ivy, English	Hedera helix	floss flower	snow-on-the-mountain
honeylocust	Gleditsia triacanthos	elaeagnus, silverberry	Elaeagnus pungens	lily-of-the-nile	Agapanthus spp.	forget-me-not	stock
larch, Japanese	Larix kaempferi	euonymus, spreading	Euonymus kiautschovica	lilyturf, bigblue	Liriope muscari	four o'clock	sunflower
locust, black	Robinia pseudoacacia	euonymus, winged	Euonymus alatus	marigold	Tagetes spp.	gaillardia	sweet alyssum
maple, Norway	Acer platanoides	euonymus, wintercreeper	Euonymus fortunei	myoporum	Myoporum laetum	gladiolus	sweet pea
maple, red	Acer rubrum	firethorn	Pyracantha spp.	plumbago, dwarf	Ceratostigma umbaginoides	golden glow	sweet sultan
maple, silver	Acer saccharinum	forsythia	Forsythia spp.	rockrose	Cistus spp.	impatiens	sweet william
maple, sugar	Acer saccharum	guava, pineapple	Feijoa sellowiana	rosemary	Rosmarinus officinalis	ixora	vinca
oak, pin	Quercus palustris	holly	Ilex spp.	rupturewort	Herniaria glabra	lobelia	yarrow
oak, scarlet	Quercus coccinea	honeysuckle	Lonicera spp.	snow-in-summer	Cerastium tomentosum	lupine	zinnia
peach	Prunus persica	India hawthorn	Raphiolepis indica	speedwell	Veronica spp.		
pine, Austrian	Pinus nigra	juniper	Juniperus spp.	St. Johnswort	Hypericum coris		
pine, eastern white	Pinus strobus	laurel, mountain	Kalmia latifolia	stonecrop (sedum)	Sedum spp.		
pine, Japanese black	Pinus thunbergiana	lilac, common	Syringa vulgaris	strawberry, beach	Fragaria chiloensis		
pine, loblolly	Pinus taeda	mockorange	Philadelphus spp.	thrift	Armeria maritima		
pine, red	Pinus resinosa	pittosporum, Japanese	Pittosporum tobira	verbena	Verbena spp.		
pine, scotch	Pinus sylvestris	privet	Ligustrum spp.	wirevine, creeping	Muehlenbeckia axillaris		
planetree, London	Platanus acerifolia	redcedar, eastern	Juniperus virginiana	yarrow, wooly	Achillea tomentosa		
plum	Prunus spp.	rhododendron	Rhododendron spp.				
redbud, eastern	Cercis canadensis	spiraea, vanhouttei	Spiraea vanhouttei				
spruce, Colorado	Picea pungens	viburnum	Viburnum spp.				
spruce, Norway	Picea abies	weigela	Weigela spp.				
spruce, white	Picea glauca	willow	Salix spp.				
sweetgum	Liquidambar styraciflua	yew, anglojap	Taxus media				
sycamore	Platanus occidentalis	yew, Japanese	Taxus cuspidata				
tuliptree	Liriodendron tulipifera	yewpine	Podocarpus macrophyllus				
walnut, black	Juglans nigra						

Weeds Controlled				Weeds Suppressed			
Weeds controlled when applied prior to weed germination:				Weeds suppressed when applied prior to weed germination:			
Annual Grasses		Broadleaf Weeds					
Common Name	Scientific Name	Common Name	Scientific Name	Common Name	Scientific Name	Common Name	Scientific Name
annual bluegrass	Poa annua	chickweed, common	Stellaria media	creeping spurge	Euphorbia serpens		
barnyardgrass (watergrass)	Echinochloa crus-galli	knotweed, prostrate	Polygonum aviculare	creeping woodsorrel	Oxalis corniculata		
crabgrass, large or hairy	Digitaria sanguinalis	lambsquartars, common	Chenopodium album	panicum, Texas	Panicum texanum		
crabgrass, smooth	Digitaria ischaemum	pigweed	Amaranthus spp.	prostrate spurge	Euphorbia humistrata		
goosegrass (silver crabgrass or crowfoot)	Eleusine indica	purslane, common	Portulaca oleracea	pusley, Florida	Richardia scabra		
green foxtail	Setaria viridis			spotted spurge	Euphorbia maculata		
yellow foxtail	Setaria glauca			woodsorrel	Oxalis spp.		

Application Directions

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer’s directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

maintaining half in water and at a right angle to the first. A splashboard mounted below the release point or drop type spreaders can improve the uniformity of product distribution.

**Uses**  
**Established Turfgrass**

**Application Timing**

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application. Apply this product 1 to 2 weeks prior to the germination of annual grass weeds. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor.

In areas where germination of the annual weed grass species continues for extended period of time, make a second application at the recommended rate 8 to 10 weeks after the first application on cool season turfgrass and 10 to 12 weeks after the first application on warm season turfgrass.

Broadcast Application Rates	Cool Season Turfgrass	Warm Season Turfgrass
Summer annual grass weeds (spring application)	174 to 349 lb/acre (4 to 8 lb/1000 sq ft)	349 lb/acre (8 lb/1000 sq ft)
Winter annual grass weeds (fall application)	174 to 349 lb/acre (4 to 8 lb/1000 sq ft)	349 lb/acre (8 lb/1000 sq ft)

This product may also be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time. For a split application, apply this product at 174 lb plus 174 lb per acre for cool season turfgrass and in the northern half of the warm season turfgrass area, or 233 lb plus 233 lb per acre in the southern half of the warm season turfgrass area. Make the second application 8 to 10 weeks following the 174 lb per acre rate and 10 to 12 weeks following the 233 lb per acre rate.

**Specific Use Restrictions**

- Do not make more than 2 applications of this product per year on turfgrass.
- **Maximum Single Application Rate:** 349 lb of Turf Fertilizer – Contains Team Pro 0.86% per acre.
- **Minimum Re-Treatment Interval:** Cool season turf – 56 to 70 days; warm season turf – 70 to 84 days.
- This product must be watered in as soon as possible after application.

**Established Ornamental Plants in Landscape Settings**

Apply to ornamental trees, shrubs, groundcovers and flowers, and to non-bearing fruit and nut trees and non-bearing vineyards, which are defined as plants that will not bear fruit for at least 1 year after treatment.

**Application Timing**

Apply 1 to 2 weeks prior to the germination of weed seeds or clean cultivate and hoe to remove existing weeds before applying this product. Apply to the smooth soil surface (no large clods) to ensure uniform distribution around established plants or on ornamental beds.

**Broadcast Application Rate**

Apply this product at a rate of 4 to 8 lb per 1000 sq ft as a spring application or 4 to 8 lb per 1000 sq ft as a fall application. In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application. Use the higher rate in the rate range where heavy weed infestation is anticipated. Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours after application.

**Specific Use Restrictions**

- Do not make more than 2 applications of this product per year on all ornamental plants.
- **Non-bearing fruit and nut trees, non-bearing vineyards:**
  - **Maximum Single Application Rate:** 8 lb of this product – Contains Team Pro 0.86% per 1000 sq ft.
  - Applications to plants that will bear fruits, berries, or nuts within 12 months are prohibited.
- **Landscape ornamentals:**
  - **Maximum Single Application Rate:** 8 lb of this product – Contains Team Pro 0.86% per 1000 sq ft.

**Terms and Conditions of Use**

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

**Warranty Disclaimer**

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT PERMITTED BY LAW, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

**Inherent Risks of Use**

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

**Limitation of Remedies**

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.