

KEEP OUT OF REACH OF CHILDREN CAUTION

In the state of New York, the product must only be used by commercial applicators at no more than 294 lbs of Dimension 0.17% FG per acre per year (0.5 lb dithiopyr/acre/year). In Nassau and Suffolk Counties of New York, do not exceed 147 lb of Dimension 0.17% FG per acre per year (0.25 lb dithiopyr/acre/year)

> It is a violation of law in Pennsylvania, Maryland, New Hampshire, and New Jersey to apply this product to turf at a rate that results in greater than 0.7 lb per 1,000 sq.ft. of readily available nitrogen or more than 0.9 lb per 1,000 sq.ft. of total nitrogen with a least 20% of which shall consist of slow release nitrogen per application. Additional restrictions apply to total nitrogen applied to turf per season. Other state and local restrictions may apply.

Florida Applications: Do not exceed 6 applications per year on bermudagrass in the south. Do not apply more than 294 lb (0.5 lb active ingredient) per acre (6.8 lb per 1000 sq ft) per application or more than 882 lb (1.5 lb active ingredient) per acre (20.4 lb per 1000 sq ft) per year if using split or sequential applications.

Do not apply near water storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back onto your lawn and garden. [Maryland Applications]

No more than 0.7 lb per 1,000 sq.ft. of readily available nitrogen during any given 30 day period. No more than 0.9 lb (cool season grasses) or 1.0 lb (warm season grasses) per 1,000 sq.ft. of total nitrogen with a least 20% of which shall consist of slow release nitrogen during any given 30 day period.

Lebanon[®] Pro

FERTILIZER with Dimension® 0.17% FG

Turf and Ornamental Herbicide

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)

17.0% Urea Nitrogen*

Derived From: Urea, polymer coated urea.

*4.25% slowly available nitrogen from polymer coated urea.



Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.cdfa.ca.gov/egov/is/fert/

http://www.oda.state.or.us/dbs/heavy_metal/search.lasso http://www.agr.wa.gov/PestFert/Fertilizers/ProductDatabase.aspx

Calibration Settings at Specified Rate					
Spreader	3.4 lbs. /1,000 FT ²	5.1 lbs. /1,000 FT ²	6.8 lbs. /1,000 FT ²		
LebanonTurf	41/2	51/4	6		
Andersons AccuPro	L	0	R		
Earthway Rotary	151/2	18	22		
Gandy	27	30	34		
Lely	5½ II	6½ II	7½ II		
Lesco (letter/Numeric)	i/17	J/21	L/24		
PennMulch HVO	M	0	R		
ProScape SS	M	0	R		
Spyker .	41/2	51/4	6		
Vicon (all models)	25	29	33		

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

For season-long control of crabgrass and control or suppression of listed annual grasses including golf course fairways, roughs and tee boxes.

> EPA Reg. No. 62719-485-961 EPA Est. No. 961-PA-01:961-IL-01 Superscript used is first letter of lot number. This product contains 0.085 pound of the active ingredient dithiopyr per 50 pound bag

Dimension is a ®Trademarks of Corteva Agriscience and its affiliated companies.

ACTIVE INGREDIENT:	
dithiopyr: 3,5- pyridinedicarbothioic acid,2-(difluoromethyl)- 4-(2-methylpropyl)-6-(trifluoromethyl)-S,	
S-dimethyl ester	0.179
OTHER INGREDIENTS:	
TOTAL	100.009

- . Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes
- For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes Eve Irritation • Harmful If Inhaled

Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Hot Line Number: Have the product container or label with you when calling a poison control center (1-800-222-1222) or doctor or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment or general product information

Environmental Hazards

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

Non-target Organism Advisory: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize drift.

and broadleaf weeds in ornamental landscape areas, established lawns and ornamental turf, Groundwater Advisory: This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

> Surface Water Advisory: This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground. This product is classified as having a high potential for reaching surface water via runoff for

Refer to back of bag for Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 888-208-1368.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

Pesticide Storage: Store this product only in its original container in a dry, cool, secured storage area.

Pesticide Disposal: Wastes resulting from the use of this product that cannot be used or chemically reprocessed must be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available, or dispose of empty bag in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Herbicide Resistance Management

For resistance management, this product is a Group 3 herbicide based upon the mode of action classification system of the Weed Science Society of America. Any weed population may contain or develop plants naturally resistant to this product and other Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay development of herbicide resistance, take one or more of the following steps.

- Alternate use of this product or other Group 3 herbicides with other products with different herbicide groups that control the same weeds.
- This product can be tank mixed or used sequentially with other approved products to broaden the spectrum of weed control, provide multiple modes of action, and control weeds that this product does not control.
- Use herbicides based on an IPM program.
- · Monitor treated areas and control escaped weeds.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product and switch to another management strategy or herbicide with a different mode of action, if available.
- For more information or to report suspected resistance, contact your extension specialist or contact a customer service representative at

Use Directions for Turf

This product provides season-long control of crabgrass and control or suppression of other listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. When used according to label directions, this product will not harm nearby established ornamentals identified as "Tolerant Ornamentals" elsewhere in this label.

Use Precautions for Turf

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. Initial application of this product should be made only after turfgrasses have developed a vigorous root system, a uniform stand, and have received at least two mowings following seeding, sodding, or sprigging. Use of this product on turfgrasses that are not well-established, or weakened by stress from weather, pests, disease, chemical, or mechanical influences, may increase the chances of turf injury.

Precautions for Reseeding, Overseeding, or Sprigging

- Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application program totaling
- 6.8 lb/1000 sq ft (0.5 lb active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses
- When reseeding or overseeding, follow established cultural practices for seedbed preparation, seeding, irrigation and fertilization.
- For best overseeding results, use mechanical or power seeding equipment (slit seeders) designed to provide good contact between soil and seed.

Precautions for Control of Crabgrass

Early postemergence weed control with this product is limited to crabgrass. Postemergence application effectively controls crabgrass, but only if application is made prior to the 3-leaf stage of growth or first tiller. For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product.

Use Restrictions

- Do not use clippings from treated turf for mulching around vegetables or fruit trees.
- Do not allow people or pets to enter treated area until dust has settled.
- Do not graze livestock or feed forage cut from areas treated with this product.
- Do not apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed:

Cool-Season Grasses		Warm-Season Grasses	
bentgrass, creeping †	Agrostis palustris	bahiagrass	Paspalum notatum
bluegrass, Kentucky	Poa pratensis	bermudagrass	Cynodon dactylon
fescue, fine ††	Festuca rubra	buffalograss †††	Buchloe dactyloides
fescue, tall	Festuca arundinacea	carpetgrass	Axonopus affinis
ryegrass, perennial	Lolium perenne	centipedegrass	Eremochloa ophiuroides
		kikuyugrass	Pennisetum clandestinum
	Ī	St. Augustinegrass	Stenotaphrum secundatum
		zoysiagrass	Zoysia japonica

† Suppression only

† Use of this product on certain varieties of creeping bentgrass, such as 'cohansey', 'carmen', seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. Do not apply this product to colonial bentgrass (Agrostis tenuis).

†† Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'atlanta', 'banner', 'beauty', 'bilgart', 'CF-2', 'enjoy', 'HF-93', 'highlight', 'ivalo', 'Jamestown', 'koket', 'majenta', 'many', pennlawn', 'tamara', 'tatjana', 'waldorf', and 'Waldina'. Not all varieties of fine fescue have been tested.

††† Do not use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Distributed by:

Lebanon Seaboard Corporation 1600 E. Cumberland St. • Lebanon, PA 17042 800-233-0628 • (717) 273-1685

NET WEIGHT ■ LBS.(22.7 KG)

Note: It is not possible to evaluate all turfgrass varieties for tolerance to this product. It is suggested that the user evaluate the response of turfgrass varieties not listed on this label on a limited area, at specified use rates, prior to initiating large-scale use.

Application Directions for Turf

Apply Dimension® 0.17% FG turf and ornamental herbicide with equipment that provides a uniform particle distribution. Calibrate application equipment prior to usage. Avoid streaking, skips, or excessoverlaps during

Preemergence Application Rates, Frequency and Timing

For preemergence grass and broadleaf weed control, apply Dimension® 0.17% FG turf and ornamental herbicide as single or sequential application at 147 to 294 pounds (0.25 to 0.5 lb active ingredient) per acre. Applicators may choose to make a single application or sequential applications at 147 to 294 pounds per acre at 5 to 10 week intervals based on one or more of the factors listed below. Avoid streaking, skips, or excess

- Length of residual weed control desired
 Height of turf (lower cut turf may require higher use rates)
- History and success of weed control at the application site (use higher listed application rates if herbicide treatment history is unknown or weed control was poor with previous applications)
- Exposure to high temperatures and heavy rainfall or irrigation (this will shorten the residual preemergence
- · On turf sites adjacent to hard surfaces including driveways, sidewalks and parking lots where residual activity may be reduced
- Some target weed species (such as but not limited to Poa annua, goosegrass and sandbur) will require higher use rates

Tips for Improved Weed Control

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the
- The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.

Maximum Use Rates (Applies to all Turfgrass Uses)

- Do not apply more than 294 lb (0.5 lb active ingredient) per acre (6.8 lb per 1000 sq ft) per application. Do not apply more than 6 times a year and do not apply more than 882 lb (1.5 lb active ingredient) per acre (20.4 lb per 1000 sq ft) per year if using split or sequential applications.
- To minimize the potential for turfgrass injury, do not make applications within five weeks of the previous
- In the state of New York, the product must only be used by commercial applicators at no more than 294 lbs of Dimension 0.17% FG per acre per year (0. 5 lb dithiopyr/acre/year). In Nassau and Suffolk Counties of New York, do not exceed 147 lb of Dimension 0.17% FG per acre per year (0.25 lb dithiopyr/acre/year)

Control of Craborass

Preemergence and Early Postemergence Applications from Late Winter through Summer

- This product provides "preemergence" control of crabgrass in established turfgrasses (including the large, smooth, and southern species) when applied prior to germination and emergence of crabgrass at use rates from 147 to 294 pounds (0.25 to 0.5 lb active ingredient) per acre.
- This product also provides "early postemergence" control of crabgrass if applied in late spring or early summer prior to the 3-leaf stage of growth or first tiller as described below. This stage of growth generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. The practical benefit of early postemergence crabgrass control is a 2 - 4-week longer period of time (window of application) during which crabgrass control can be achieved

Fall Applications

This product can also be applied in the late summer through early fall (late August through November) at use rates from 147 to 294 pounds per acre to provide control of crabgrass through the early part of the next spring. Fall application of this product may be followed by an appropriately timed application in the following spring for season-long control of crabgrass.

Weeds Controlled or Suppressed

Spring Applications

When applied preemergence (prior to germination) for control of crabgrass in the spring, this product will also control or suppress the following weeds

Grasses		Broadleaf Weeds			
barley	Hordeum spp.	bittercress †	Cardamine spp.		
barnyardgrass	Echinochloa crus-galli	carpetweed	Mollugo verticillata		
bluegrass, annual	Poa annua	chickweed †	Stellaria spp.		
brome	Bromus spp.	geranium, Carolina †	Geranium carolinianum		
crabgrass, large	Digitaria sanguinalis	henbit	Lamium spp.		
crabgrass, smooth	Digitaria ischaemum	knotweed, prostrate	Polygonum aviculare		
crabgrass, Southern	Digitaria ciliaris	lespedeza, common †	Lespedeza striata		
crowfootgrass †	Dactyloctenium aegyptium	marestail	Conyza canadensis		
dallisgrass (seedling)	Paspalum dilatatum	medic, black	Medicago lupulina		
goosegrass	Eleusine indica	mustard	Brassica spp.		
foxtail, green	Setaria verdi	oxalis, buttercup	Oxalis pes-caprae		
foxtail, yellow	Setaria faberi	pineappleweed †	Matricaria matricarioides		
kikuyugrass †	Pennisetum clandestinum	pigweed, redroot	Amaranthus retroflexus		
oats, wild	Avena fatua	parsley-piert †	Alchemilla arvensis		
ryegrass (annual & perennial)	Lolium spp.	purslane, common	Portulaca oleracea		
sandbur	Cenchrus spp.	rocket, London	Sisymbrium irio		
smutgrass	Sporobolus indicus	shepherdspurse	Capsella bursa-pastoris		
		speedwell, corn †	Veronica arvensis		
		spurge, garden	Euphorbia hirta		
		spurge, prostrate	Euphorbia humistrata		
		spurge, spotted	Euphorbia maculata		
		woodsorrel, creeping	Oxalis corniculata		
		woodsorrel, yellow	Oxalis stricta		

Fall Applications

When applied preemergence (prior to germination) in late summer through early fall (late August through November) this product will control or suppress the following weeds in addition to providing preemergence control of crabarass in the following spring.

bluegrass, annual	Poa annua
bittercress	Cardamine spp.
chickweed	Stellaria spp.
henbit	Lamium spp.
geranium, Carolina	Geranium carolinianum
parsley-piert	Alchemilla arvensis
pineappleweed	Matricaria matricarioides
shepherdspurse	Capsella bursa-pastoris

Application Instructions and Equipment

This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the specified application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference. Apply this product uniformly over the treatment area. More uniformity of application can usually be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Avoid streaking, skips, or overlaps during application.

Check equipment frequently to verify calibration and proper functioning.

Use Directions for Landscape Ornamentals

Dimension 0.17% FG provides preemergence control or suppression of listed annual grass and broadleaf weeds in plantings of ornamental plants listed on this label. This product may be applied in ornamental gardens, parks, golf courses and residential areas where ornamental plants are grown for aesthetic purposes. Apply this product with a properly calibrated spreader that will provide uniform particle distribution. Follow instructions in the "Application Instructions and Equipment" section above.

Use Rates

This product may be applied as a single application or split application. Split or sequential applications may be used to provide improved weed control or to provide extended weed control in areas with long growing seasons. To make a split application, divide the rate in the following table into separate applications made 5-10 weeks apart. Sequential applications may be made in the late summer following a spring application or in the spring after a fall application, provided maximum application rates per year given below are not exceeded.

		. ,	
294	6.8	0.68 (10.9 oz)	
Pounds of Product/Acre	Pounds of Product/1000 sq ft	Pounds of Product/100 sq ft †	

† Rate based on an area of 100 sq ft for treatment of small areas.

Maximum Use Rates

- Do not apply more than 294 lb (0.5 lb active ingredient) per acre (6.8 lb per 1000 sq ft) per application.
- Do not apply more than 6 times per year and do not apply more than 882 lb (1.5 lb active ingredient) per acre (20.4 lb per 1000 sq ft) per year if using split or sequential applications.
- To minimize the potential for plant injury, do not make additional applications within five weeks of the previous application.
- In the state of New York, the product must only be used by commercial applicators at no more than 294 lbs of Dimension 0.17% FG per acre per year (0.5 lb dithiopyr/acre/year). In Nassau and Suffolk Counties of New York, do not exceed 147 lb of Dimension 0.17% FG per acre per year (0.25 lb

Application Timing and Instructions

Dimension 0.17% FG is effective as a preemergence herbicide but will not control established weeds. Applications to mulched areas or bare ground must be made prior to weed seed germination. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Existing weedy vegetation may be controlled by hand weeding, cultivation or using postemergence herbicides prior to application.

Tips for Improved Weed Control

- The herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.
- Avoid disturbance of treated areas. Loss of weed control may result if the treated soil surface is disturbed by soil mixing or tillage.

• To avoid foliar damage to ornamentals, irrigate immediately after application to remove any herbicide granules adhering to foliage.

Use Restrictions

- Do not apply this product directly to bare roots of ornamental plants as injury may result.
- Do not incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots
- Do not apply to soil around ornamental plants that are under stress resulting from drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides, or damage due to insects, nematodes or disease.
- Do not apply under conditions that would affect uniformity of application or distribution on the soil surface. Uneven product distribution will result in uneven weed control. Application under windy conditions can result in uneven distribution or cause herbicide granules to drift from the intended treatment area.
- Apply this product only to established ornamentals.
- Do not use on food producing trees and ornamentals.

Weeds Controlled or Suppressed

Used as directed, Dimension 0.17% FG will control crabgrass and control or suppress other listed grass and broadleaf weeds when applied prior to their germination. Refer to "Use Directions for Turf" section fo a complete listing of weeds controlled or suppressed. This product will not control established broadleaf weeds or grasses, except for crabgrass in early stages of development. The area to be treated should be free of weeds prior to application.

Tolerant Ornamentals

The ornamentals listed below have shown tolerance to this product when applied according to directions in this label. However, this product has not been tested on all ornamental species, all cultivars of species on this label, or under all possible growing conditions. For species or cultivars not listed only treat a few plants in a limited area and observe for tolerance under local growing conditions prior to large-scale use.

Common Name	Botanical Name	Tolerant Cultivars
abelia, dwarf	Abelia X grandiflora	nana
ajuga	Ajuga reptans	bronze
	Ajuga genevensis	bronze beauty
almond, flowering	Prunus gladulosa	
apple †	Malus pumila	
arborvitae	Thuja occidentalis	nigra pyramidalis smaragh techny woodwardii
arborvitae, golden	Thuja orientalis	

ardenia

heather, twisted

nemlock, Canada

Common Name	Botanical Name	Tolerant Cultivars	Common Name	Botanical Name	Tolerant Cultivars	Common Name	Botanical Name	Tolerant Cultivars
aster, Chinese	Callistephus chinensis	dwarf queen	holly	llex spp.	blue boy	pine, mugo	Pinus mugho	. U.U. GIR GURUNGIO
ash, green	Fraxinus pennsylvanica		· ·		blue girl	pine, Scotch	Pinus sylvestris	
ash, mountain	Sorbus aucuparia			Ilex X meserveae	burfodii China girl	pine, slash pine, Swiss mt.	Pinus elliottii Pinus mugo	
ash, purple azalea	Fraxinus americana Rhododendron spp.	brilliant			compacta	pine, Virginia	Pinus virginiana	
azaioa	Timododonaron opp.	buccaneer			forsteri	pine, white	Pinus strobus	
		carror			hellerie Japanese northern	pineapple, guava †	Feijoa sellowiana Pieris taiwanensis	
		chimes (Belgian) Elsie Lee			beauty	pittosporum, Japan	i ielis taiwarierisis	
		exbury			needlepoint	potentilla	Potentilla nepalensis	abbotswood
		fashion		Ilex X attenuata	Nellie R. Stevens Savannah	privet	Potentilla fruticosa Ligustrum japonicum	golden vicary
		hardijzer beauty hershey red	holly, Chinese	llex comuta	Savarillari	priver	Ligusti di Ti Japoni cum	regal
		higasa	holly, Japanese	llex crenata				texanum
		hinocrimson Holland (hybrid)	holly, yaupon honeysuckle	Ilex vomitoria Lonicera japonica	claveyis dwarf			wax yellow tipped
		Marion Lee	Horicysuokic	Lonicora japonica	halliana	privet, glossy	Ligustrum lucidum	успом пррец
		northern lights			tatarian Canadian white	pyracantha	Pyracantha koldzumii	gnome
		orange cup orchid lights	hosta	Hosta spp.	zebelli red hosta albo marginata			lalandei victory
		snow	1100100	Hosta lancifolia	and manymata	queen palm	Arecastrum rammanzoffianum	
		southern charm	ice plant	Carpobrotus edulis		quince, Japanese †	Dhadadandran ann	allarina
azalea, flame azalea, kirishima	Rhododendron calendulaceum		impatiens iris	Impatiens spp. Iris spp.	dwarf blue	rhododendron	Rhododendron spp	album Cunningham white
bamboo, heavenly			ino	то орр.	wedgewood			fashion
barberry	Berberis thunbergii	aurea	ivy, English	Hedera helix	Bulgaria			hardy
		dwarf pigmy green	jasmine, Asian	Trachelospermum asiaticum	thorndale			pjm purple gem
		kobold	juniper	Juniperus spp.	arcadia			silvery pink
		pygmy red	, ,	I	armstrong	rhododendron, carolina	Rhododendron carolinianum	
barberry, purple	1	rose glow atropurpurea		Juniperus horizontalis	bar harbor blue chip	rhododendron, catawba rockcress	Rhododendron catawbiense Arabis spp.	snowcap
basket flower	Gaillardia grandiflora			Jamporao Horizontallo	blue pacific	rhodie max (rosebay)	Rhododendron maximum	
bearberry (common)	Arctostaphylos uva-ursi	Massachusetts			blue rug	rose †	Rosa banksiae	luta
bee balm begonia	Monarda didyma Begonia spp.				blue star broadmoor	rosemary † rosemary, bog	Rosmarinus officinalis Andromeda polifolia	nana
birch, river	Betula nigra				buffalo	salvia	Salvia farinacea	rhea
blackeyed Susan	Rudbeckia hirta	goldstrum			Calgary carpet	sedum	Sedum spurium	dragon blood red
blanket flower blueberry †	Gaillardia spp. Vaccinium spp.	bluecrop			emerald sea emerald spreader			red carpet yellow
bluebelly [vaccinium spp.	blue jay			endora compacta	snapdragon	Antirrhinum spp.	yonow
		jersey			fruitlandi	sourwood	Oxydendrum arboreum	fII
		north blue northland			green gold tip	spiraea spiraea	Astilbe X arendsii Spiraea spp.	fanall Anthony waterer red
bottlebrush	Callistemon citrinus	HOITHANA			hetzi	opiidod	ориава врр.	dolchicá
boxwood, Japanese		japonica			hughes			froebeli pink
boxwood, weller broom	Buxus sempervirens Cytisus spp.	moonlight			Manhattan blue parsoni			goldenflame red snowmound white
biodiii	Genista pilosa	Vancouver gold			pfitzeriana			Van Houtte white
bugle carpet					plumosa	spiraea, garland spruce, black hills	Spiraea spp. Picea glauca	
camellia	Camellia japonica	debutante mathotiana supreme			prince of Wales procumbens dwarf	spruce, Colorado blue	Picea pungens	
		chansonette			San Jose	spruce, norway	Picea abies	
	Camellia sasanqua				sargent blue	spruce, white sweetflag, grassy-leaved	Picea glauca Acorus gramineus	conica
candy tuft carex, variegated	lberis spp. Carex	snow white			sargent green scandia	sweetgum	Liquidambar styraciflua	
cedar, red	Juniperus virginiana				scopulorum moonglow	sycamore	Platanus occidentalis	
celosia	Celosia spp. Centaurea montana				scopulorum skyrocket spartan	tree fern (tiki fern) trumpet flower, evening	Asparagus virgatus Gelsemium sempervirens	
centaura cockscomb, plumosa	Celosia cristata	scarlet plumosa		Juniperus chinensis	tamariscifolia	tulip	Tulip, spp	apeldoorn
coleus	Coleus blumei	red kewpie		Juniperus sabina	weberi	verbena, shrub	Lantana sellowiana	Ot David
columbine copper leaf	Aguilegia spp. Acalypha wilkesiana				youngstown yukon belle	vervain viburnum	Verbena spp. Viburnum spp.	St. Paul American cranberry bush
coreopsis	Coreopsis spp.	moonbeam	king palm	Archontophoenix cunninghamiana	yakon bolio	The difficulty	Tiballian opp.	arrowwood
corn flower	Centaurea spp.		laurel, Australian	Pittosporum tobira				common snowball
cotoneaster coyotebrush	Cotoneaster apiculatus Baccharis pilularis		laurel, mountain leucothoe	Kalmia latifolia Leucothoe fontanesiana				European cranberry bush linden
cycads	Cycads revoluta		ligustrum, Japanese	Ligustrum japonicum				Mohican
cypress, bald	Taxodium distichum		lily, African	Agapanthus africanus	albus	vinca (periwinkle)	Vines miner	Wright
cypress, Italian cypress, Japanese false	Cupressus sempervirens Chamaecyparis obtusa	glauca gracilis	lily, African blue	1	Peter Pan	windmill palm	Vinca minor Trachycarpus fortunei	
cypress, leyland	Cupressocyparis leylandii	gracins	lily of the valley	Pieris japonica	Mt. Fire	xylosma	Xylosma congestum	
daffodil	Narcissus spp.	King Alfred	lilýturf	Liriopé muscari	evergreen giant	yarrow	Achillea spp.	dwarf
daylilly	Hemerocallis spp.	aztec gold bright vellow (hybrid)			lilac beauty majestic	yaupon yew	Ilex vomitoria Taxus cuspidata	dwarf denisiformis
		bright yellow (hybrid) single gold (evergreen) wilsonis yellow			monroe white	,	Taxus X media	
dianthus /	Dienthus co-	wilsonis yellow "	liriona graca	Liriona enicata	variegata	† Ornamental species only	: Do not use on food producing trees	and ornamentals.
dianthus (sweet william) delphinium	Dianthus spp. Delphinium spp.	magic fountain	liriope, green maple, Japanese	Liriope spicata Acer japonicum	+	Terms and Conditions of	Use	
dogwood	Cornus florida	· ·	maple, Norway	Acer platanoides			Warranty Disclaimer, Inherent Ris	sks of Use, and Limitation
dogwood, American	Cornus sericea	flavarimaea	maple, red †	Acer rubrum		Remedies are not accept	otable, return unopened package	at once to the seller for
douglas fir dusty miller	Pseudotsuga menziesii Senecio cineraria		maple, silver maple sugar †	Acer saccharinum Acer saccharum	-		To the extent permitted by law, us	
elm, drake	Ulmus parvifolia		marigold	Tagetes patula	honeycomb		of the terms under Warranty Discl	laimer, Inherent Risks o
euonymus	Euonymus fortunei	argenteo-variegata			variegata wheeleris dwarf	Limitation of Remedies.		
		auereo-marginata colorata	mock orange †	Philadelphus spp	golden	Warranty Disclaimer		
		emerald gaiety		1	snowflake double white		product conforms to the chemical	
		emerald en gold	mondo grass	Ophiopogon japonicus	euppyeide		poses stated on the label when u inherent risks set forth below. To	
		gold edge gold princess	moss rose myrtle, crape	Portulaca grandiflora Lagerstroemia indica	sunnyside faurei		THER EXPRESS OR IMPLIED W	
		silver king	, ,		langer		ARTICULAR PURPOSE OR ANY	
		tricolor			muskogee standard pink	WARRANTY.		
fan palm, European	Chamaerops humilis	vegetus	myrtle, wax	Myrica cerifera	sianuaru pink	Inherent Risks of Use		
fan palm, Mexican	Washingtonia robusta		nandina	Nandina domestica	compacta		ate all risks associated with use o	f this product. Plant ini
fern (various)	Asparagus spp.		narojegua	Marajeeue ann	nana	performance, or other un	nintended consequences may res	sult because of such fac
fescue fetterbush	Festuca glauca Leucothoe fontanesiana	rainbow	narcissus oak, laurel	Narcissus spp. Quercus laurifolia	 	the product contrary to la	abel instructions (including conditi	ions noted on the label,
ficus	Ficus retusa	nitidia	oak, pin	Quercus palustris			e, soil conditions, etc.), abnormal	
fir fraser	Abies fraseri	Amald -lt	oak, red	Quercus rubra			ricanes), presence of other materi	
forsythia	Forsythia X intermedia	Arnold dwarf bronxensis dwarf	oak, southern oak, willow	Quercus virginiana Quercus phellos	 		h are beyond the control of Seller.	. 10 the extent permitte
		lynwood gold	oleander	Nerium oleander	hardy red	such risks shall be assu	ilieu by Duyel.	
1	0	meadowlark			petité pink	Limitation of Remedies	hy law the evaluative remodules	neede or damagaa ras-
	Spring Glory	weeping		1.	Sister Agnes	TO the exterit permitted	by law, the exclusive remedy for I	osses oi uamages rest

ation of for a full refund ny other user s of Use and

abel and is nce with the d by law RCHANTABILITY OR IMPLIED

iniury, lack of factors as use of el. such as excessive rainfall, oplication, or itted by law, all ich risks shall be assumed by Buver.

Limitation of Remedies

Barbara karst

Osteospermum fruticosui

Pachysandra terminalis

Cortaderia selloana

/iola x wittrockiana

Petunia X hvbrida

Photinia X fraser

Pieris japonica

Pinus palustris

Pinus thunbergiana

Pinus nigra

Bougainvillea glabra Prunus persica Vinca minor

weeping

radicans

cockspur white

crimson cloud

Washington white Mediterranean pin

enchantress

Jack Evans

blue bird

Gardenia iasminoides

Eucalyptus citriodora Crataegus spp.

Erica cinerea

Tsuga canadensis Hibiscus spp.

palm, bangalov

pampas grass

hotinia red tin

pine, Australian

pine, longlea

pine, Japanese black pine, loblolly

pansy paper flower

petunia

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, tort, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following

- Refund of purchase price paid by buyer or user for product bought, or
- 2. Replacement of amount of product used.

To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.