FERTILIZER with

Dimension® 0.15% FG

Turf and Ornamental Herbicide

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	22 %
3.0% Ammoniacal Nitrogen	
2.0% Water Insoluble Nitrogen*	
13.8% Urea Nitrogen	
3.2% Other Water Soluble Nitrogen*	
Soluble Potash (K,0)	6%
Sulfur (S)	3.5%
3.5% Combined Sulfur (S)	

Derived From: Ammonium Sulfate, Methylene Urea, Urea, Muriate

*5.2% Slowly available nitrogen from methylene urea. Chlorine (CI) not more than 5.0%



Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.cdfa.ca.gov/egov/is/fert/

http://www.oda.state.or.us/dbs/heavy_metal/search.lasso http://www.agr.wa.gov/PestFert/Fertilizers/ProductDatabase.aspx

	Calib	ration Settings at	Specified Rate	
Spreader	2.9# /1,000 FT ²	3.8# /1,000 FT ²	5.8# /1,000 FT ²	7.7#/ 1,000 FT ²
LebanonTurf	41/2	43/4	5 ½	61/2
Andersons AccuPro	K	M	P	S
Earthway Rotary	15	16	20	24
Gandy	26	28	32	36
Lely	5 II	6 II	7 II	8 II
Lesco (letter/Numer	ic) i/17	J/21	L/24	N/28
PennMulch HVO	Ĺ	N	P	S
ProScape SS	L	N	P	S
Spyker	41/2	43/4	5 ½	61/2
Vicon (all models)	24	26	31	34
-				on of spreader, speed s for desired coverage

For season-long control of crabgrass and control or suppression of listed annual grasses and broadleaf weeds in ornamental landscape areas, established lawns and ornamental turf,

including golf course fairways, roughs and tee boxes. ACTIVE INGREDIENT: (L) (D) EPA Est. No. 961-PA-01:961-IL-01 dithiopyr: 3,5- pyridinedicarbothioic acid,2-(difluoromethyl)- 4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester... 0.15% OTHER INGREDIENTS: Superscript used is first letter of lot number.

This product contains 0.075 pound of the active ingredient dithiopyr per 50 pound bag. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes

For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns. Dimension is a ®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow.

In the state of New York, this product may be applied only by commercial applicators at a minimum of 167 lb (0.25 lb active ingredient) per acre (3.8 lb per 1000 sq ft) and no more than 333 lb (0.5 lb active ingredient) per acre (7.7 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

It is a violation of law in Maryland and New Jersey to apply this product to turf at a rate that results in greater than 0.7 lb per 1,000 sq.ft. of readily available nitrogen or more than 0.9 lb per 1,000 sq.ft. of total nitrogen with a least 20% of which shall consist of slow release nitrogen per application. Additional restrictions apply to total nitrogen applied to turf per season. Other state and local restrictions may apply. It is a violation of law in Maryland and New Jersey to apply this product to turf at a rate that results in greater than 0.7 lb per 1,000 sq.ft. of readily available nitrogen or more than 0.9 lb per 1,000 sq.ft. of total nitrogen with a least 20% of which shall consist of slow release nitrogen per application. Additional restrictions apply to total nitrogen applied to turf per season. Other state and local restrictions may apply. [Maryland or New Jersey Applications]

Do not apply near water storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back onto your lawn and garden. [Maryland Applications] No more than 0.7 lb per 1,000 sq.ft. of readily available nitrogen during any given 30 day period. No more than 0.9 lb (cool season grasses) or 1.0 lb (warm season grasses) per 1,000 sq.ft. of total nitrogen with a least 20% of which shall consist of slow release nitrogen during any given 30 day period. [Virginia Applications]

In Virginia and New Jersey this lawn/turf fertilizer contains phosphorous and is only for nonagricultural use on (i) turf during its first growing season, (ii) on turf areas being repaired or renovated, and (iii) on turf where a soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not for the routine maintenance of turf. [Virginia and New Jersey Applications which contain Phosphorus]

KEEP OUT OF REACH OF CHILDREN

CAUTION **FIRST AID**

in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Hot Line Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 888-208-1368 day or night, for emergency treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes Eye Irritation • Harmful If Inhaled

Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause

Environmental Hazards This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment

Refer to back of bag for Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once

In case of emergency endangering health or the environment involving this product, call 888-208-1368. Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Directions for Use

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal Pesticide Storage: Store this product only in its original container in a dry, cool, secured storage area.

Pesticide Disposal: Wastes resulting from the use of this product that cannot be used or chemically reprocessed must be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures. Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.



Distributed by:

Lebanon Seaboard Corporation 1600 E. Cumberland St. • Lebanon, PA 17042 800-233-0628 • (717) 273-1685

Use Directions for Turf

Lebanon ProScape Fertilizer with Dimension® 0.15% FG turf and ornamental herbicide provides season-long control of crabgrass and control or suppression of other listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. When used according to label directions, this product will not harm nearby established ornamentals identified as "Tolerant Ornamentals" elsewhere in this label.

Use Precautions for Turf

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. Initial application of this product should be made only after turfgrasses have developed a vigorous root system, a uniform stand, and have received at least two mowings following seeding, sodding, or sprigging. Use of this product on turfgrasses that are not well-established, or weakened by stress from weather, pests, disease, chemical, or mechanical influences, may increase the chances of turf injury. Sweep any product that lands on the driveway, sidewalk, or street back onto lawn and

• Florida Applications: Do not exceed 6 applications per year on bermudagrass in the south Do not apply more than 333 lb (0.5 lb active ingredient) per acre (7.7 lb per 1000 sq ft) per application or more than 1000 lb (1.5 lb active ingredient) per acre (23.1 lb per 1000 sq ft) per year if using split or sequential applications.

Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed:

Cool-Season Grasses		Warm-Season Grasses	
bentgrass, creeping †	Agrostis palustris	bahiagrass	Paspalum notatum
bluegrass, Kentucky	Poa pratensis	bermudagrass	Cynodon dactylon
fescue, fine ††	Festuca rubra	buffalograss †††	Buchloe dactyloides
fescue, tall	Festuca arundinacea	carpetgrass	Axonopus affinis
ryegrass, perennial	Lolium perenne	centipedegrass	Eremochloa ophiuroides
	•	kikuyugrass	Pennisetum clandestinum
		St. Augustinegrass	Stenotaphrum secundatum
		zoysiagrass	Zoysia japonica

† Use of this product on certain varieties of creeping bentgrass, such as 'cohansey', 'carmen', seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. Do not apply this product to colonial bentgrass (Agrostis tenuis).

"It'l Use of this product on certain varieties of fine fescue may result in undesirable turl injury. The following fine fescue varieties have been found to be sensitive to this product: 'atlanta', 'banner', 'beauty', 'bilgart', 'CF-2', 'enjoy', 'HF-93', 'highlight', 'ivalo', 'Jamestown', 'koket', 'majenta', 'mary', pennlawn', 'tamara', 'tatjana', 'waldorf', and 'Waldina'. Not all varieties of fine fescue have been tested.

††† Do not use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Precautions for Reseeding, Overseeding, or Sprigging

Precautions for Control of Crabgrass

Use Restrictions

For preemergence grass and broadleaf weed control, apply Dimension® 0.15% FG turf and ornamental herbicide as single or sequential application at 167 to 333 pounds (0.25 to 0.5 lb active ingredient) per acre. Applicators may choose to make a single application or sequential applications at 167 to 333 pounds per acre at 5 to 10 week intervals based on one or more of the factors listed below

- Length of the stocked control destined

 Height of the (lower cut furf may require higher use rates)

 History and success of weed control at the application site (use higher listed application rates if herbicide treatment history is unknown or weed control was poor with previous applications)
- Exposure to high temperatures and heavy rainfall or irrigation (this will shorten the residual preemergence performance)
 On turf sites adjacent to hard surfaces including driveways, sidewalks and parking lots where residual activity may be reduced
- Some target weed species (such as but not limited to Poa annua, goosegrass and sandbur) will require higher use rates

Tips for Improved Weed Control • For best results, apply this product within 2 to 3 days after mowing and delay mowing again for 2 to 3 days after the application. • The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.

- Do not apply more than 333 lb (0.5 lb active ingredient) per acre (7.7 lb per 1000 sq ft) per application
- Do not apply more than 6 times per year and do not apply more than 1000 lb (1.5 lb active ingredient) per acre (23.1 lb per 1000 sq ft) per year if using split or sequential applications.
 To minimize the potential for turfgrass injury, do not make applications within five weeks of the previous application.
 In the state of New York, this product may be applied only by commercial applicators at no more than 333 lb (0.5 lb active ingredient) per acre (7.7 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

Control of Craborass

Fall Applications

Weeds Controlled or Suppressed

Grasses		Broadleaf Weeds		
barley	Hordeum spp.	bittercress †	Cardamine spp.	
barnyardgrass	Echinochloa crus-galli	carpetweed	Mollugo verticillata	
bluegrass, annual	Poa annua	chickweed †	Stellaria spp.	
brome	Bromus spp.	geranium, Carolina †	Geranium carolinianum	
crabgrass, large	Digitaria sanguinalis	ĥenbit	Lamium spp.	
crabgrass, smooth	Digitaria ischaemum	knotweed, prostrate	Polygonum aviculare	
crabgrass, Southern	Digitaria ciliaris	lespedeza, common †	Lespedeza striata	
crowfootgrass †	Dactyloctenium aegyptium	marestail	Conyza canadensis	
dallisgrass (seedling)	Paspalum dilatatum	medic, black	Medicago lupulina	
goosegrass	Eleusine indica	mustard	Brassica spp.	
foxtail, green	Setaria verdi	oxalis, buttercup	Oxalis pes-caprae	
foxtail, yellow	Setaria faberi	pineappleweed †	Matricaria matricarioides	
kikuyugrass †	Pennisetum clandestinum	pigweed, redroot	Amaranthus retroflexus	
oats, wild	Avena fatua	parsley-piert †	Alchemilla arvensis	
ryegrass (annual & perennial)	Lolium spp.	purslane, common	Portulaca oleracea	
sandbur	Cenchrus spp.	rocket, London	Sisymbrium irio	
smutgrass	Sporobolus indicus	shepherdspurse	Capsella bursa-pastoris	
		speedwell, corn †	Veronica arvensis	
		spurge, garden	Euphorbia hirta	
		spurge, prostrate	Euphorbia humistrata	
		spurge, spotted	Euphorbia maculata	
		woodsorrel, creeping	Oxalis corniculata	
		woodsorrel, yellow	Oxalis stricta	

† Suppression only

Fall Applications

bluegrass, annual	Poa annua
bittercress	Cardamine spp.
chickweed	Stellaria spp.
henbit	Lamium spp.
geranium, Carolina	Geranium carolinianum
parsley-piert	Alchemilla arvensis
pineappleweed	Matricaria matricarioides
shepherdspurse	Capsella bursa-pastoris

Weed Resistance Management

Application Instructions and Equipment

growing seasons. To make a split application, divide the rate in the following table into separate applications made 5-10 weeks apart. Sequential applications may be made in the late summer following a spring application or in the spring after a fall application, provided maximum application rates per year given below are not exceeded. Pounds of Product/1000 sq ft Pounds of Product/100 sq ft † 0.77 (12.3 oz)

† Rate based on an area of 100 sq ft for treatment of small areas.

Pounds of Product/Acre

Maximum Use Rates

•Do not apply more than 333 lb (0.5 lb active ingredient) per acre (7.7 lb per 1000 sq ft) per application.

•Do not apply more than 6 times per year and do not apply more than 1000 lb (1.5 lb active ingredient) per acre (23.1 lb per 1000 sq ft) per year if using split or sequential applications.

•To minimize the potential for plant injury, do not make additional applications within five weeks of the previous application.

•In the state of New York, this product may be applied only by commercial applicators at no more than 333 lb (0.5 lb active ingredient) per acre (7.7 lb per 1000 sq ft) per year. Use of this product in Nassau

and Suffolk counties of New York is prohibited.

Application Timing and Restrictions

Lebanon ProScape Fertilizer with Dimension® 0.15% FG turf and ornamental herbicide is effective as a preemergence herbicide, but will not control established weeds. Applications to mulched areas or bare ground must be made prior to weed seed germination. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Existing weedy vegetation may be controlled by hand weeding, cultivation or using postemergence herbicides prior to application.

Tips for Improved Weed Control

• The herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application. Avoid disturbance of treated areas. Loss of weed control may result if the treated soil surface is disturbed by soil mixing or tillage.

Use Precautions • To avoid foliar damage to ornamentals, irrigate immediately after application to remove any herbicide granules adhering to foliage

Use Restrictions

Do not apply this product directly to bare roots of ornamental plants as injury may result.
Do not incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
Do not apply to soil around ornamental plants that are under stress resulting from drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides, or damage due to insects, nematodes or disease.
Do not apply under conditions that would affect uniformity of application or distribution on the soil surface. Uneven product distribution will result in uneven weed control. Application under windy conditions can result in uneven distribution or cause herbicide granules to drift from the intended treatment area.

Apply this product only to established ornamentals

Do not use on food producing trees and ornamentals

Weeds Controlled or Suppressed

Used as directed, Lebanon ProScape Fertilizer with Dimension® 0.15% FG turf and ornamental herbicide will control crabgrass and control or suppress other listed grass and broadleaf weeds when applied prior to their germination. Refer to "Use Directions for Turf" section for a complete listing of weeds controlled or suppressed. This product will not control established broadleaf weeds or grasses, except for crabgrass in early stages of development. The area to be treated should be free of weeds prior to application

Tolerant Ornamentals

The ornamentals listed below have shown tolerance to this product when applied according to directions in this label. However, this product has not been tested on all ornamental species, all

	ass varieties for tolerance to this produscale use.	**	- au Carianian		conditions prior to large-scale	el, or under all possible growing con use.	·	, ,		
• Reseeding, overseeding, or sprigging of a (0.5 lb active ingredient per acre) or more n	areas treated with this product within 10	or Reseeding, Overseeding 0 weeks after a single application of this pro		plication program totaling 7.7 lb/1000 sq ft	Common Name	Botanical Name	Tolerant Cultivars	Common Name	Botanical Name	Tolerant Cultivars
When reseeding or overseeding, follow es For best overseeding results, use mechan	stablished cultural practices for seedbe	ed preparation, seeding, irrigation and fertilized			abelia, dwarf ajuga	Abelia X grandiflora Ajuga reptans Ajuga genevensis	nana bronze bronze beauty		Juniperus horizontalis	blue rug blue star broadmoor
		autions for Control of Crab	•		almond, flowering apple †	Prunus gladulosa Malus pumila		l		buffalo Calgary carpet
Early postemergence weed control with this tiller. For best results, cultural practices that turfgrass has recovered from the mechanical turfgrass has recovered from the mechanical turfgrass.	at disturb the soil, such as core-, spike-	temergence application effectively controls on the controls of the control of	rabgrass, but only if application is made all be done before application of this pr	le prior to the 3-leaf stage of growth or first product. Do not apply this product until	arborvitae	Thuja occidentalis	nigra pyramidalis smaragh			emerald sea emerald spreader endora compacta
Do not use clippings from treated turf for n	,, , , , , , , , , , , , , , , , , , , ,	Use Restrictions					techny woodwardii			fruitlandi green
Do not allow people or pets to enter treate Do not graze livestock or feed forage cut from the content of the content	ed area until dust has settled.				arborvitae, golden aster, Chinese ash, green	Thuja orientalis Callistephus chinensis Fraxinus pennsylvanica	dwarf queen			gold tip hetzi hughes
nitrogen with a least 20% of which shall cor		a rate that results in greater than 0.7 lb per ication. Additional restrictions apply to total			ash, mountain ash, purple	Sorbus aucuparia Fraxinus americana				Manhattan blue parsoni
		rain is expected. Apply this product only to y	our lawn, and sweep any product that la	lands on the driveway, sidewalk, or street	azalea	Rhododendron spp.	brilliant buccaneer carror			pfitzeriana plumosa prince of Wales
back onto your lawn and garden. [Maryland	,	n 30 day period. No more than 0.9 lb (cool so	eason grasses) or 1.0 lb (warm season).	graceac) par 1 000 ca ft. of total nitrogen			chimes (belgian) elsie lee			procumbens dwarf San Jose
with a least 20% of which shall consist of sl	low release nitrogen during any given 3	30 day period. [Virginia Applications]					exbury fashion			sargent blue sargent green
In Virginia and New Jersey this lawn/turf ferti where a soil test performed within the last 3	years indicates a phosphorous deficiend	for nonagricultural use on (i) turf during its firm cy. This fertilizer is not for the routine mainter	nance of turf. [Virginia and New Jersey Ap	g repaired or renovated, and (iii) on turf Applications which contain Phosphorus]			hardijzer beauty hershey red higasa			scandia scopulorum moonglow scopulorum skyrocket
	mension® 0.15% FG turf and ornamen	pplication Directions for To natal herbicide with equipment that provides a		e application equipment prior to usage.			hinocrimson holland (hybrid)		Juniperus chinensis	spartan tamariscifolia
Avoid streaking, skips, or excess overlaps of	Preemergence	Application Rates, Freque					marion lee northern lights orange cup		Juniperus sabina	weberi youngstown yukon belle
		FG turf and ornamental herbicide as single of tions at 167 to 333 pounds per acre at 5 to 1					orchid lights snow	king palm laurel, Australian	Archontophoenix cunningham Pittosporum tobira	
Length of residual weed control desired Height of turf (lower cut turf may require h					azalea, flame azalea, kirishima	Rhododendron calendulaceu	southern charm m	laurel, mountain leucothoe ligustrum, Japanese	Kalmia latifolia Leucothoe fontanesiana Ligustrum japonicum	
 Exposure to high temperatures and heavy 	rainfall or irrigation (this will shorten t	plication rates if herbicide treatment history the residual preemergence performance) ing lots where residual activity may be reduce	•	with previous applications)	bamboo, heavenly barberry	Berberis thunbergii	aurea	lily, African	Agapanthus africanus	albus Peter Pan
Some target weed species (such as but n	not limited to Poa annua, goosegrass a						dwarf pigmy green kobold	lily, African blue lily of the valley lilyturf	Pieris japonica Liriope muscari	Mt. Fire evergreen giant
• For best results, apply this product within	2 to 3 days after mowing and delay mo	owing again for 2 to 3 days after the applica	ation.	ad by 1/2 inch of rainfall or sprinkler			pygmy red rose glow	illyturi	Linope muscan	lilac beauty majestic
• The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 3 days after application.			barberry, purple basket flower	Gaillardia grandiflora	atropurpurea	liviana gyaan	Liviana anicota	monroe white variegata		
Maximum Use Rates (Applies to all Turfgrass Uses) • Do not apply more than 333 lb (0.5 lb active ingredient) per acre (7.7 lb per 1000 sq ft) per application. • Do not apply more than 6 times per year and do not apply more than 1000 lb (1.5 lb active ingredient) per acre (23.1 lb per 1000 sq ft) per year if using split or sequential applications.				bearberry (common) bee balm begonia	Arctostaphylos uva-ursi Monarda didyma Begonia spp.	Massachusetts	liriope, green maple, Japanese maple, Norway	Liriope spicata Acer japonicum Acer platanoides		
. To minimize the potential for turfgrass inju	ury, do not make applications within five		.,,,,		birch, river blackeyed Susan	Betula nigra Rudbeckia hirta	goldstrum	maple, red † maple, silver	Acer rubrum Acer saccharinum	
and Suffolk counties of New York is prohibit		Control of Crabgrass	,,		blanket flower blueberry †	Gaillardia spp. Vaccinium spp.	bluecrop blue jay	maple sugar † marigold	Acer saccharum Tagetes patula	honeycomb variegata
Preemergence and Early Postemergence		rough Summer	all and a second and				jersey north blue	mock orange †	Philadelphus spp	wheeleris dwarf golden
 This product provides "preemergence" couse rates from 167 to 333 pounds of product This product also provides "early posteme 	ct (0.25 to 0.5 lb ai) per acre. ergence" control of crabgrass if applied	d in late spring or early summer prior to the	3-leaf stage of growth or first tiller as des	escribed below. This stage of growth	bottlebrush	Callistemon citrinus	northland	mondo grass moss rose	Ophiopogon japonicus	snowflake double white
	abgrass seedlings first become visible i	in the established turfgrass. The practical b			boxwood, Japanese boxwood, weller broom	Buxus sempervirens Cytisus spp.	japonica moonlight	myrtle, crape	Portulaca grandiflora Lagerstroemia indica	faurei langer
		Fall Applications st through November) at use rates from 167		trol of crabgrass through the early part of	bugle carpet	Genista pilosa	Vancouver gold			muskogee standard pink
the next spring. Fall application of this proc		ely timed application in the following spring eds Controlled or Suppres			camellia	Camellia japonica	debutante mathotiana supreme chansonette	myrtle, wax nandina	Myrica cerifera Nandina domestica	compacta
Spring Applications When applied preemergence (prior to germ		spring, this product will also control or supp			candy tuft	Camellia sasanqua Iberis spp.	snow white	narcissus oak, laurel	Narcissus spp. Quercus laurifolia	
Grasses barley	Hordeum spp.	Broadleaf Weeds bittercress †	Cardamine spp.		carex, variegated cedar, red celosia	Carex Juniperus virginiana Celosia spp.		oak, pin oak, red oak, southern	Quercus palustris Quercus rubra Quercus virginiana	
barnyardgrass bluegrass, annual	Echinochloa crus-galli Poa annua	carpetweed chickweed †	Mollugo verticillata Stellaria spp.		centaura cockscomb, plumosa	Centaurea montana Celosia cristata	scarlet plumosa	oak, willow oleander	Quercus phellos Nerium oleander	hardy red
brome	Bromus spp.	geranium, Carolina †	Geranium carolinianum		coleus columbine	Coleus blumei Aguilegia spp.	red kewpie	a cha a an a way um	October a way was fautice or way	petite pink Sister Agnes
crabgrass, large crabgrass, smooth	Digitaria sanguinalis Digitaria ischaemum	henbit knotweed, prostrate	Lamium spp. Polygonum aviculare		copper leaf coreopsis corn flower	Acalypha wilkesiana Coreopsis spp. Centaurea spp.	moonbeam	osteospermum pachysandra palm, bangalow	Osteospermum fruticosum Pachysandra terminalis	wirligig
crabgrass, Southern crowfootgrass †	Digitaria ciliaris Dactyloctenium aegyptium	lespedeza, common † marestail	Lespedeza striata Conyza canadensis		cotoneaster coyotebrush	Cotoneaster apiculatus Baccharis pilularis		pampas grass pansy	Cortaderia selloana Viola x wittrockiana	
dallisgrass (seedling) goosegrass	Paspalum dilatatum Eleusine indica	medic, black mustard	Medicago lupulina Brassica spp.		cycads cypress, bald cypress, Italian	Cycads revoluta Taxodium distichum Cupressus sempervirens	glauca	paper flower peach † periwinkle, dwarf	Bougainvillea glabra Prunus persica Vinca minor	Barbara karst
foxtail, green foxtail, yellow	Setaria verdi Setaria faberi	oxalis, buttercup pineappleweed †	Oxalis pes-caprae Matricaria matricarioides		cypress, Italian cypress, Japanese false cypress, leyland	Chamaecyparis obtusa Cupressocyparis leylandii	gracilis	petunia photinia, red tip	Petunia X hybrida Photinia X fraseri	picoti
kikuyugrass † oats, wild	Pennisetum clandestinum Avena fatua	pigweed, redroot parsley-piert †	Amaranthus retroflexus Alchemilla arvensis		daffodil daylilly	Narcissus spp. Hemerocallis spp.	King Alfred aztec gold	pieris pine, Australian	Pieris japonica Pinus nigra	
ryegrass (annual & perennial) sandbur	Cenchrus spp.	purslane, common rocket, London	Portulaca oleracea Sisymbrium irio				bright yellow (hybrid) single gold (evergreen) wilsonís yellow	pine, Japanese black pine, loblolly pine, longleaf	Pinus thunbergiana Pinus taeda Pinus palustris	
smutgrass	Sporobolus indicus	shepherdspurse speedwell, corn †	Capsella bursa-pastoris Veronica arvensis		dianthus (sweet william) delphinium	Dianthus spp. Delphinium spp.	magic fountain	pine, mugo pine, Scotch	Pinus mugho Pinus sylvestris	
		spurge, garden spurge, prostrate	Euphorbia hirta Euphorbia humistrata		dogwood dogwood, American douglas fir	Cornus florida Cornus sericea Pseudotsuga menziesii	flavarimaea	pine, slash pine, Swiss mt. pine, Virginia	Pinus elliottii Pinus mugo Pinus virginiana	
		spurge, spotted woodsorrel, creeping	Euphorbia maculata Oxalis corniculata		dusty miller elm, drake	Senecio cineraria Ulmus parvifolia		pine, white pineapple, guava †	Pinus strobus Feijoa sellowiana	
† Suppression only		woodsorrel, yellow	Oxalis stricta		euonymus	Euonymus fortunei	argenteo-variegata auereo-marginata	pieris pittosporum, Japan potentilla	Pieris taiwanensis Potentilla nepalensis	
Fall Applications							colorata emerald gaiety emerald en gold	poteritiia		abhotewood
			alona de la composición del composición de la co	ing weeds in addition to providing			emerald ën gold	privet	Potentilla fruticosa Ligustrum japonicum	abbotswood golden vicary
preemergence control of crabgrass in the fo	ollowing spring.	iali (late August tirrough November) triis pro	oduct will control or suppress the followin	g			gold edge gold princess	privet	Potentilla fruticosa	golden vicary regal texanum
bluegrass, annual Poa al		iali (late August tiriough November) triis pro	oduct will control or suppress the following				gold edge gold princess silver king tricolor		Potentilla fruticosa	golden vicary regal
bluegrass, annual Poa al bittercress Carda chickweed Stellar henbit Lamiu	nnua unine spp. ria spp. um spp.	iali (late August ilirough November) tilis pro	oduct will control or suppress the following	· · · · · · · · · · · · · · · · · · ·	fan palm, European fan palm, Mexican	Chamaerops humilis Washingtonia robusta	gold edge gold princess silver king	privet, glossy pyracantha	Potentilla fruticosa Ligustrum japonicum	golden vicary regal texanum wax yellow tipped gnome lalandei
bluegrass, annual Poa ai bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher	nnua mine spp. ria spp. ilum spp. ilum carolinianum milla arvensis	iali (late August ilirough November) illis pro	oduct will control or suppress the following	· · · · · · · · · · · · · · · · · · ·	fan palm, Mexican fern (various) fescue	Washingtonia robusta Asparagus spp. Festuca glauca	gold edge gold princess silver king tricolor vegetus	privet, glossy pyracantha	Potentilla fruticosa Ligustrum japonicum Ligustrum lucidum	golden vicary regal texanum wax yellow tipped gnome lalandei victory
bluegrass, annual Poa ai bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric	nnua mine spp. ria spp. im spp. nium carolinianum	iali (late August ilirough November) tilis pro	oduct will control or suppress the following		fan palm, Mexican fem (various) fescue fetterbush ficus fir fraser	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri	gold edge gold princess silver king tricolor vegetus rainbow nitidia	privet, glossy pyracantha	Potentilla fruticosa Ligustrum japonicum Ligustrum lucidum Pyracantha koidzumii	golden vicary regal texanum wax yellow tipped gnome lalandei victory album Cunningham white
bluegrass, annual Poa ai bittercress Carda Chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capse	nnua umine spp. ria spp. Im spp. Iium carolinianum Iimila arvensis Iaria matricarioides Iila bursa-pastoris	/eed Resistance Managem	ent		fan palm, Mexican fern (various) fescue fetterbush ficus	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf	privet, glossy pyracantha queen palm quince, Japanese †	Potentilla fruticosa Ligustrum japonicum Ligustrum lucidum Pyracantha koidzumii Arecastrum rammanzoffian	golden vicary regal texanum wax yellow tipped gnome lalandei victory album Cunningham white fashion hardy
bluegrass, annual Poa ai bittercress Carda Chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capse	minua minie spp. mia spp. mis spp. mis spp. mis spp. mis spp. mium carolinianum milla arvensis earia matricarioides ella bursa-pastoris W on® 0.15% FG turf and ornamental he		ent		fan palm, Mexican fem (various) fescue fetterbush ficus fir fraser	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri	gold edge gold princess silver king tricolor vegetus rainbow nitidia Amold dwarf	privet, glossy pyracantha queen palm quince, Japanese †	Potentilla fruticosa Ligustrum japonicum Ligustrum lucidum Pyracantha koidzumii Arecastrum rammanzoffian	golden vicary regal texanum wax yellow tipped gnome lalandei victory um album Cunningham white fashion
bluegrass, annual Poa au bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capse Lebanon ProScape Fertilizer with Dimensis system of the Weed Science Society of Am Proactively implementing diversified weed include the use of multiple herbicides with	nnua umine spp. ria spp. um spp. ilum carolinianum milla arvensis aria matricarioides ella bursa-pastoris W on® 0.15% FG turf and ornamental he nerica. control strategies to minimize selectior different modes of action and overlapp	/eed Resistance Managem erbicide, which contains the active ingredien on for weed populations resistant to one or no ping weed spectrum with or without tillage or	ent It dithiopyr is a Group 3 herbicide based	d on the mode of action classification rersified weed management program may	fan palm, Mexican ferm (various) fescue fetterbush ficus fir fraser forsythia	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping	privet, glossy pyracantha queen palm quince, Japanese † rhododendron	Potentilla fruticosa Ligustrum japonicum Ligustrum japonicum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron spp Rhododendron carolinianum Rhododendron catawbiense	golden vicary regal texanum wax yellow tipped gnome lalandei victory um album Cunningham white fashion hardy pim purple gem silvery pink
bluegrass, annual Poa ai bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capses Lebanon ProScape Fertilizer with Dimensic system of the Weed Science Society of An Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in	nnua umine spp. ria spp. im spp. ium carolinianum milla arvensis caria matricarioides ella bursa-pastoris W on® 0.15% FG turf and ornamental he merica. control strategies to minimize selection different modes of action and overlapp nportant to delay the selection for resis	Veed Resistance Managemerbicide, which contains the active ingredien on for weed populations resistant to one or no ping weed spectrum with or without tillage of stance.	ent It dithiopyr is a Group 3 herbicide based more herbicides is a best practice. A dive perations and/or other cultural practices.	d on the mode of action classification rersified weed management program may	fan palm, Mexican ferm (various) fescue fetterbush ficus fir fraser forsythia fuchsia galium gardenia	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum Gardenia jasminoides	gold edge gold princess silver king tricolor vegetus rainbow nitidia Amold dwarf bronxensis dwarf lynwood gold meadowlark	privet, glossy pyracantha queen palm quince, Japanese † rhododendron rhododendron, carolina rhododendron, catawba rockcress rhodle max (rosebay)	Potentilla fruticosa Ligustrum japonicum Ligustrum lucidum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron spp Rhododendron carolinianum Rhododendron catawbiense Arabis spp. Rhododendron maximum	golden vicary regal texanum wax yellow tipped gnome lalandei victory m album Cunningham white fashion hardy pjm purple gem silvery pink
bluegrass, annual Poa ai bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capse Lebanon ProScape Fertilizer with Dimensis system of the Weed Science Society of Am Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this produc To aid in the prevention of developing wee	nnua umine spp. ria spp. lim spp. lim spp. lim spp. lim spp. lim carolinianum milla arvensis saria matricarioides sella bursa-pastoris Won® 0.15% FG turf and ornamental he nerica. control strategies to minimize selection different modes of action and overlapp nportant to delay the selection for resis et depends on the successful implement etds resistant to this product, users shou	Veed Resistance Managemerbicide, which contains the active ingredien on for weed populations resistant to one or no ping weed spectrum with or without tillage of stance. Intation of a weed resistance management product:	ent It dithiopyr is a Group 3 herbicide based more herbicides is a best practice. A dive perations and/or other cultural practices. program.	d on the mode of action classification rersified weed management program may	fan palm, Mexican ferm (various) fescue fetterbush ficus fir fraser forsythia	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping mystery radicans	privet, glossy pyracantha queen palm quince, Japanese † rhododendron rhododendron, carolina rhododendron, catawba rockcress rhodle max (rosebay) rose † rosemary † rosemary, bog	Potentilla fruticosa Ligustrum japonicum Ligustrum lucidum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron spp Rhododendron carolinianum Rhododendron catawbiense Arabis spp. Rhododendron maximum Rosa banksiae Rosmarinus officinalis Andromeda polifolia	golden vicary regal texanum wax yellow tipped gnome lalandei victory um album Cunningham white fashion hardy pjm purple gem silvery pink snowcap luta nana
bluegrass, annual Poa ai bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capse Lebanon ProScape Fertilizer with Dimensic system of the Weed Science Society of An Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this productor in the prevention of developing wee Scout fields before application to ensure!	mnua umine spp. ria spp. ria spp. m spp. nium carolinianum milla arvensis saria matricarioides sella bursa-pastoris W on® 0.15% FG turf and ornamental he merica. control strategies to minimize selection different modes of action and overlapp nportant to delay the selection for resis at depends on the successful implement eder resistant to this product, users shou herbicides and rates will be appropriate ndown herbicide application or tillage.	Veed Resistance Management provided, which contains the active ingredient on for weed populations resistant to one or no ping weed spectrum with or without tillage ostance. Intation of a weed resistance management provided to the weed species and weed sizes presented.	ent It dithiopyr is a Group 3 herbicide based more herbicides is a best practice. A dive perations and/or other cultural practices. program.	d on the mode of action classification rersified weed management program may	fan palm, Mexican fem (various) fesscue fetterbush ficus fir fraser forsythia fuchsia galium gardenia geranium gum	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum Gardenia jasminoides Pelargonium X hortorum Eucalyptus citriodora	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping mystery radicans cockspur white crimson cloud enchantress	privet, glossy pyracantha queen palm quince, Japanese † rhododendron rhododendron, carolina rhododendron, catawba rockcress rhodie max (rosebay) rose † rosemary †	Potentilla fruticosa Ligustrum japonicum Ligustrum lucidum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron spp Rhododendron carolinianum Rhododendron catawbiense Arabis spp. Rhododendron maximum Hosa banksiae Rosmarinus officinalis	golden vicary regal texanum wax yellow tipped gnome lalandei victory album Cunningham white fashion hardy pim purple gem silvery pink snowcap luta nana rhea dragon blood red
bluegrass, annual Poa at bluegrass, annual Poa at carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capses Lebanon ProScape Fertilizer with Dimensic system of the Weed Science Society of An Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this productor of aid in the prevention of developing ween Scout fields before application to ensure 1 Start with a clean field, using either a burn 1 sing post-emergence herbicides or tar Apply full rates of Dimension 0.15% FG for Scout fields after application to detect we	nnua umine spp. ria spp. im spp. ium carolinianum milla arvensis caria matricarioides ella bursa-pastoris Won® 0.15% FG turf and ornamental he merica. control strategies to minimize selection different modes of action and overlapp riportant to delay the selection for resis et depends on the successful implement eds resistant to this product, users shounder herbicides and rates will be appropriate modown herbicide application or tillage. Ink mixes, control weeds early when the or the most difficult to control of weed in the descapes or shifts in control of weed in the descapes or shifts in control of weed red seapes or shifts in control of weed seaps weed seaps or shifts in control of weed seaps or shifts	Veed Resistance Management of the work of the weed spectrum with or without tillage of stance. Intation of a weed resistance management pulld: Ite for the weed species and weed sizes preside at the specified time to minimize with graphs.	ent It dithiopyr is a Group 3 herbicide based more herbicides is a best practice. A dive perations and/or other cultural practices. program. sent.	d on the mode of action classification rersified weed management program may	fan palm, Mexican ferm (various) fesscue fetterbush ficus fir fraser forsythia fuchsia galium gardenia geranium gum hawthorn	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum Gardenia jasminoides Pelargonium X hortorum Eucalyptus citriodora Crataegus spp.	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping mystery radicans cockspur white crimson cloud	privet, glossy pyracantha queen palm quince, Japanese † rhododendron, carolina rhododendron, catawba rockcress rhodie max (rosebay) rose † rosemary † rosemary, bog salvia sedum	Potentilla fruticosa Ligustrum japonicum Ligustrum japonicum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron carolinianum Rhododendron catawbiense Arabis spp. Rhododendron maximum Rosa banksiae Rosmarinus officinalis Andromeda polifolia Salvia farinacea Sedum spurium Antirrhinum spp.	golden vicary regal texanum wax yellow tipped gnome lalandei victory um album Cunningham white fashion hardy pim purple gem silvery pink snowcap luta nana rhea
bluegrass, annual Poa ai bittercress Carda Chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capse Lebanon ProScape Fertilizer with Dimensic system of the Weed Science Society of An Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this produc To aid in the prevention of developing wee Scout fields before application to ensure 1 start with a clean field, using either a burn 1 fl using post-emergence herbicides or tar Apply full rates of Dimension 0.15% FG for Scout fields after application to detect we Control weed escapes before they reproc	mnua umine spp. ria spp. lim s	Veed Resistance Managemerbicide, which contains the active ingredient on for weed populations resistant to one or no ping weed spectrum with or without tillage of stance. Intation of a weed resistance management product the for the weed species and weed sizes preserved are relatively small. The field at the specified time to minimize with dispecies. Ye was representative, lead to your local company representative.	ent It dithiopyr is a Group 3 herbicide based more herbicides is a best practice. A dive perations and/or other cultural practices. program. sent. eed escapes. ocal retailer, or county extension agent.	d on the mode of action classification versified weed management program may s. Research has demonstrated that using	fan palm, Mexican ferm (various) fescue fetterbush ficus fir fraser forsythia fuchsia galium gardenia geranium gum hawthorn	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum Gardenia jasminoides Pelargonium X hortorum Eucalyptus citriodora Crataegus spp.	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping mystery radicans cockspur white crimson cloud enchantress Jack Evans Washington white Mediterranean pink blue bird	privet, glossy pyracantha queen palm quince, Japanese † rhododendron rhododendron, carolina rhododendron, catawba rockcress rhodie max (rosebay) rose † rosemary † rosemary, bog salvia sedum snapdragon sourwood spiraea	Potentilla fruticosa Ligustrum Japonicum Ligustrum Japonicum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron spp Rhododendron carolinianum Rhododendron catawbiense Arabis spp. Rhododendron maximum Hosa banksiae Rosmarinus officinalis Andromeda polifolia Salvia farinacea Sedum spurium Antirrhinum spp. Oxydendrum arboreum Astilbe X arendsii	golden vicary regal texanum wax yellow tipped gnome lalandei victory um album Cunningham white fashion hardy pim purple gem silvery pink snowcap luta nana rhea dragon blood red red carpet yellow fanall
bluegrass, annual Poa at bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capse Lebanon ProScape Fertilizer with Dimensic system of the Weed Science Society of An Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this productor of the Weed Science Society of An Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this productor of the Weed Science Society of An Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of the product fields before application to ensure 1 Start with a clean field, using either a burn of the production of the Start with a clean field, using either a burn of the Start with a clean field, using either a burn of the Start with a clean field, using either a burn of the Start with a clean field, using either a burn of the Start with a clean field, using either a burn of the Start with a clean field, using either a burn of the Start with a clean field using either a burn of the Start with a clean field using either a burn of the Start with a clean field using either a burn of the Start with a clean field using either a burn of the Start with a clean field using either a burn of the Start with Dimension of the Start with	mnua umine spp. ria spp. um spp. ilum carolinianum milla arvensis aria matricarioides alla bursa-pastoris W on® 0.15% FG turf and ornamental he nerica. control strategies to minimize selection different modes of action and overlapp nportant to delay the selection for resis at depends on the successful implement eds resistant to this product, users shoutherbicides and rates will be appropriate andown herbicide application or tillage. As mixes, control weeds early when the or the most difficult to control weed in the or the most difficult to control weed in the or the most difficult to control weed in the or the product against a particular we we, crop advisor, or extension agent to this product specified for your local ce	Veed Resistance Managemerbicide, which contains the active ingredien on for weed populations resistant to one or no ping weed spectrum with or without tillage of stance. Intation of a weed resistance management puld: te for the weed species and weed sizes preside the weed species and weed sizes preside at the specified time to minimize with species.	ent It dithiopyr is a Group 3 herbicide based more herbicides is a best practice. A dive perations and/or other cultural practices. program. sent. leed escapes. ocal retailer, or county extension agent. s MOA have been found in your region. I are multiple effective modes of action for	d on the mode of action classification versified weed management program may s. Research has demonstrated that using	fan palm, Mexican ferm (various) fescue fescue fetterbush ficus fir fraser forsythia fuchsia galium gardenia geranium gum hawthom	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum Gardenia jasminoides Pelargonium X hortorum Eucalyptus citriodora Crataegus spp. Erica cinerea Tsuga canadensis Hibiscus spp.	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping mystery radicans cockspur white crimson cloud enchantress Jack Evans Washington white Mediterranean pink	privet, glossy pyracantha queen palm quince, Japanese † rhododendron rhododendron, carolina rhododendron, catawba rockcress rhodie max (rosebay) rose † rosemary † rosemary, bog salvia sedum	Potentilla fruticosa Ligustrum japonicum Ligustrum japonicum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron spp Rhododendron carolinianum Rhododendron catawbiense Arabis spp. Rhododendron maximum Rosa banksiae Rosmarinus officinalis Andromeda polifolia Salvia farinacea Sedum spurium Antirrhinum spp. Oxydendrum arboreum	golden vicary regal texanum wax yellow tipped gnome lalandei victory um album Cunningham white fashion hardy pim purple gem silvery pink snowcap luta nana rhea dragon blood red red carpet yellow
bluegrass, annual Poa au biutercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capse Lebanon ProScape Fertilizer with Dimensis system of the Weed Science Society of Am Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this productor if using post-emergence herbicides or tar Apply full rates of Dimension 0.15% FG for Scout fields after application to detect we Control weed escapes before they reproduce the control weed escapes before they reproduced the control weed escapes before they reproduced to the control weed escapes before they reproduced they are supplication as the control of the control weed to the control weed t	mnua umine spp. ria spp. um spp. im sp	Veed Resistance Managemerbicide, which contains the active ingredient on for weed populations resistant to one or no ping weed spectrum with or without tillage of stance. Intation of a weed resistance management product to the form the weed species and weed sizes preside the field at the specified time to minimize with dispecies. Interest of the weed species and weed sizes president to the field at the specified time to minimize with dispecies. In do ut if suspected resistant weeds to this orditions. Tank mix products so that there are of action other than Group 3 and/or use not service of the suspected for the services.	ent It dithiopyr is a Group 3 herbicide based more herbicides is a best practice. A dive perations and/or other cultural practices. program. sent. leed escapes. ocal retailer, or county extension agent. s MOA have been found in your region. I are multiple effective modes of action for onchemical methods to remove escapes.	d on the mode of action classification versified weed management program may s. Research has demonstrated that using	fan palm, Mexican fem (various) fescue fetterbush ficus fir fraser forsythia fuchsia galium gardenia geranium gum hawthom heather, twisted hemlock, Canada hibiscus	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum Gardenia jasminoides Pelargonium X hortorum Eucalyptus citriodora Crataegus spp. Erica cinerea Tsuga canadensis	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping mystery radicans cockspur white crimson cloud enchantress Jack Evans Washington white Mediterranean pink blue bird birlliant hula girl blue boy blue girl burlodii	privet, glossy pyracantha queen palm quince, Japanese † rhododendron rhododendron, carolina rhododendron, catawba rockcress rhodie max (rosebay) rose † rosemary † rosemary, bog salvia sedum snapdragon sourwood spiraea	Potentilla fruticosa Ligustrum Japonicum Ligustrum Japonicum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron spp Rhododendron carolinianum Rhododendron catawbiense Arabis spp. Rhododendron maximum Hosa banksiae Rosmarinus officinalis Andromeda polifolia Salvia farinacea Sedum spurium Antirrhinum spp. Oxydendrum arboreum Astilbe X arendsii	golden vicary regal texanum wax yellow tipped gnome lalandei victory um album Cunningham white fashion hardy pim purple gem silvery pink snowcap luta nana rhea dragon blood red red carpet yellow fanall Anthony waterer red dolchica froebeli pink goldenflame red snowmound white
bluegrass, annual Poa ai bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capses Lebanon ProScape Fertilizer with Dimensic system of the Weed Science Society of Am Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this productor of the Weed Science Society of Am Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this productor of the Weed Science Society of Am Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this productor of aid in the prevention of developing wee Scout fields before application to ensure 1 Start with a clean field, using either a burn of the Weed Science Society of Am Proactive India the Production of Science Society of Am Proactive India the Production of Science Society of Am Proactive India the Production of Science Society of Am Proactive India the Production of Science Society of Am Proactive India the Proactive India the India	mnua umine spp. ria spp. lim carolinianum milla arvensis caria matricarioides control strategies to minimize selection different modes of action and overlapp riportant to delay the selection for resis cat depends on the successful implement caria resistant to this product, users shou herbicides and rates will be appropriat mdown herbicides and rates will be appropriat her other most difficult to control weed in the control weeds early when the or the most difficult to control weed in the control weeds early when the or the most difficult to control of weed duce by seed or proliferate vegetatively e of this product against a particular we caria caria matricario to delay the selection or tillage. In the control of weed duce by seed or proliferate vegetatively e of this product against a particular we caria matricarioides the control weeds control weeds control weeds control weeds control weed duce by seed or proliferate vegetatively e of this product against a particular we control weeds control weeds control weeds control weeds control weeds control weeds duce by seed or proliferate vegetatively and overlappen control of weeds duce by seed or proliferate vegetatively and overlappen control of the control of weed duce by seed or proliferate vegetatively and overlappen control of the control of weed duce by seed or proliferate vegetatively and overlappen control of the	Veed Resistance Managemerbicide, which contains the active ingredient on for weed populations resistant to one or no ping weed spectrum with or without tillage of stance. Intation of a weed resistance management puld: te for the weed species and weed sizes preside to the weed species and weed sizes preside to the field at the specified time to minimize with a species. In do ut if suspected resistant weeds to this orditions. Tank mix products so that there a	ent It dithiopyr is a Group 3 herbicide based more herbicides is a best practice. A dive perations and/or other cultural practices. program. sent. leed escapes. ocal retailer, or county extension agent. s MOA have been found in your region. I are multiple effective modes of action for onchemical methods to remove escapes.	d on the mode of action classification versified weed management program may s. Research has demonstrated that using	fan palm, Mexican fem (various) fescue fetterbush ficus fir fraser forsythia fuchsia galium gardenia geranium gum hawthom heather, twisted hemlock, Canada hibiscus	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retus Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum Gardenia jasminoides Pelarganium X hortorum Eucalyptus citriodora Crataegus spp. Erica cinerea Tsuga canadensis Hibiscus spp.	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping mystery radicans cockspur white crimson cloud enchantress Jack Evans Washington white Mediterranean pink blue bird brilliant hula girl blue boy blue girl	privet, glossy pyracantha queen palm quince, Japanese † rhododendron, carolina rhododendron, catawba rockcress rhodie max (rosebay) rose † rosemary † rosemary, bog salvia sedum snapdragon sourwood spiraea spiraea spiraea, garland spruce, black hills	Potentilla fruticosa Ligustrum Japonicum Ligustrum Japonicum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron spp Rhododendron carolinianum Rhododendron catawbiense Arabis spp. Rhododendron maximum Hosa banksiae Rosmarinus officinalis Andromeda polifolia Salvia farinacea Sedum spurium Antirrhinum spp. Oxydendrum arboreum Astilbe X arendsii	golden vicary regal texanum wax yellow tipped gnome lalandei victory album Cunningham white fashion hardy pim purple gem silvery pink snowcap luta nana rhea dragon blood red red carpet yellow fanall Anthony waterer red dolchica froebeli pink goldenflame red
bluegrass, annual Poa al bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capse Lebanon ProScape Fertilizer with Dimensic system of the Weed Science Society of An Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this produc To aid in the prevention of developing wee Scout fields before application to ensure I Start with a clean field, using either a bur of If using post-emergence herbicides or tar Apply full rates of Dimension 0.15% FG for Scout fields after application to detect the Control weed escapes before they reprocementation of the Contact your local company representative Contact your local company representative the reported, use the application rates of if resistance is suspected, treat weed esc further seed production. Suspected herbicide-resistant weeds manual suspected herbicide-resistant weeds manual suspected herbicide-resistant weeds manual suspected herbicide resistant weeds manual suspected herbicides of non-controlled plant Surviving plants mixed with controlled ind	mnua umine spp. ria spp. lim spection and overlapp nortant to delay the selection for resis at depends on the successful implement spp. lim spp. li	Veed Resistance Managemerbicide, which contains the active ingredient on for weed populations resistant to one or no ping weed spectrum with or without tillage of stance. Intation of a weed resistance management pulld: It for the weed species and weed sizes preside the field at the specified time to minimize with dispecies. It ind out if suspected resistant weeds to this onditions. Tank mix products so that there are of action other than Group 3 and/or use no use applied, especially if control is achieved management practices as is practical:	ent It dithiopyr is a Group 3 herbicide based more herbicides is a best practice. A dive perations and/or other cultural practices. program. sent. leed escapes. ocal retailer, or county extension agent. s MOA have been found in your region. I are multiple effective modes of action for onchemical methods to remove escapes.	d on the mode of action classification versified weed management program may s. Research has demonstrated that using	fan palm, Mexican fem (various) fescue fetterbush ficus fir fraser forsythia fuchsia galium gardenia geranium gum hawthom heather, twisted hemlock, Canada hibiscus	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retus Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum Gardenia jasminoides Pelarganium X hortorum Eucalyptus citriodora Crataegus spp. Erica cinerea Tsuga canadensis Hibiscus spp.	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping mystery radicans cockspur white crimson cloud enchantress Jack Evans Washington white Mediterranean pink blue bird brilliant hula girl blue boy blue girl burfodii china girl compacta forsteri hellerie Japanese northem beauty	privet, glossy pyracantha queen palm quince, Japanese † rhododendron, carolina rhododendron, catawba rockcress rhodie max (rosebay) rose † rosemary † rosemary † rosemary, bog salvia sedum snapdragon sourwood spiraea spiraea spiraea spiraea, garland spruce, black hills spruce, Colorado blue spruce, norway	Potentilla fruicosa Ligustrum japonicum Ligustrum japonicum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron spp Rhododendron catawbiense Arabis spp. Rhododendron maximum Rosa banksiae Andromeda polifolia Salvia farinacea Sedum spurium Antirrhinum spp. Oxydendrum arboreum Astilbe X arendsii Spiraea spp. Picea glauca Picea glauca Picea pungens Picea abies	golden vicary regal texanum wax yellow tipped gnome lalandei victory album Cunningham white fashion hardy pim purple gem silvery pink snowcap luta nana rhea dragon blood red red carpet yellow fanall Anthony waterer red dolchica froebeli pink goldenflame red snowmound white Van Houtte white
bluegrass, annual Poa ai bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capse Lebanon ProScape Fertilizer with Dimensis system of the Weed Science Society of An Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this produc To aid in the prevention of developing wee Scout fields before application to ensure 1 Start with a clean field, using either a burn of If using post-emergence herbicides or tar Apply full rates of Dimension 0.15% FG fic Scout fields after application to detect we Control weed escapes before they reproc Report any incidence of non-performance Contact your local company representativen the seed production. Suspected herbicide-resistant weeds may a suspected herbicide-resistant weeds may Failure to control a weed species normall surviving plants mixed with controlled ind Additionally, users should follow as many of Use a broad spectrum herbicide with othe Utilize sequential applications of herbicide Notate the use of this product with non-G	nnua umine spp. ria spp. lim carolinianum milla arvensis laria matricarioides laria matricarioides laria bursa-pastoris Won® 0.15% FG turf and ornamental he nerica. control strategies to minimize selection different modes of action and overlapp nportant to delay the selection for resis at depends on the successful implement of the selection of tillage. As the selection of the selection of the selection of tillage and the selection of the selectio	Veed Resistance Managemerbicide, which contains the active ingredient on for weed populations resistant to one or no ping weed spectrum with or without tillage of stance. Intation of a weed resistance management product to for the weed species and weed sizes preside the field at the specified time to minimize with dispecies. Year relatively small. The field at the specified time to minimize with dispecies. Year of the weed species and weed sizes presided to your local company representative, length of the field at the specified time to minimize with species. Year of the weed species and weed to this orditions. Tank mix products so that there are of action other than Group 3 and/or use no one of action other than Group 3 and/or use no one of action other than Group 3 and/or use no one of action other than Group 3 and/or use no one of action other than Group 3 and/or use no one of action other than Group 3 and/or use no one of action other than Group 3 and/or use no one of action other than Group 3 and/or use no one of action other than Group 3 and/or use no one of action other than Group 3 and/or use no one of action other than Group 3 and/or use no one of action other than Group 3 and/or use no one of action other than Group 3 and/or use no one of action other than Group 3 and/or use no one of action other than Group 3 and/or use no other than Group 3 and 3 an	ent It dithiopyr is a Group 3 herbicide based more herbicides is a best practice. A dive perations and/or other cultural practices. program. sent. leed escapes. ocal retailer, or county extension agent. s MOA have been found in your region. I are multiple effective modes of action for onchemical methods to remove escapes, on adjacent weeds;	d on the mode of action classification versified weed management program may s. Research has demonstrated that using If resistant biotypes of target weeds have r each target weed. s, as practical, with the goal of preventing	fan palm, Mexican fem (various) fescue fetterbush ficus fir fraser forsythia fuchsia galium gardenia geranium gum hawthom heather, twisted hemlock, Canada hibiscus	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum Gardenia jasminoides Pelargonium X hortorum Eucalyptus citriodora Crataegus spp. Erica cinerea Tsuga canadensis Hibiscus spp. Ilex spp. Ilex X meserveae	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping mystery radicans cockspur white crimson cloud enchantress Jack Evans Washington white Mediterranean pink blue bird brilliant hula girl blue boy blue girl burfodii china girl compacta forsteri hellerie Japanese northem beauty needlepoint Nellie R. Stevens	privet, glossy pyracantha queen palm quince, Japanese † rhododendron, carolina rhododendron, catawba rockcress rhodie max (rosebay) rose † rosemary † rosemary, bog salvia sedum snapdragon sourwood spiraea spiraea spiraea spiraea, garland spruce, black hills spruce, Colorado blue spruce, norway spruce, white sweetflag, grassyy-le	Potentilla fruticosa Ligustrum japonicum Ligustrum japonicum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron spp Rhododendron carolinianum Rhododendron catawbiense Arabis spp. Rhododendron maximum Rosa banksiae Rosmarinus officinalis Andromeda polifolia Salvia farinacea Sedum spurium Antirrhinum spp. Oxydendrum arboreum Astilbe X arendsii Spiraea spp. Picea glauca Picea pungens Picea glauca Picea pungens Picea glauca Picea glauca	golden vicary regal texanum wax yellow tipped gnome lalandei victory um album Cunningham white fashion hardy pim purple gem silvery pink snowcap luta nana rhea dragon blood red red carpet yellow fanall Anthony waterer red dolchica froebeli pink goldenflame red snowmound white
bluegrass, annual Poa at bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capse Lebanon ProScape Fertilizer with Dimensis system of the Weed Science Society of Am Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this produc To aid in the prevention of developing wee Scout fields before application to ensure Istart with a clean field, using either a burn If using post-emergence herbicides or tar Apply full rates of Dimension 0.15% FG for Scout fields after application to detect of Contact your local company representative Control weed escapes before they reproduction. Suspected herbicide-resistant weed send If resistance is suspected, treat weed sof If resistance is suspected, treat weed sof If resistance is suspected, treat weeds many suspected herbicide-resistant weeds many failure to control a weed species normall A spreading patch of non-controlled plant Surviving plants mixed with controlled ind Additionally, users should follow as many of Use a broad spectum herbicide with other Utilizes sequential applications of herbicide Rotate the use of this product with non-Gavoid making more than two sequentials easeason unless mixed with an herbicide with the season unless mixe	minua umine spp. ria spp. lim carolinianum milla arvensis saria matricarioides ella bursa-pastoris Won® 0.15% FG turf and ornamental he nerica. control strategies to minimize selection different modes of action and overlapp nportant to delay the selection for resis et depends on the successful implement eds resistant to this product, users shot herbicides and rates will be appropriat modown herbicide application or tillage, nk mixes, control weeds early when the or the most difficult to control of weed duce by seed or proliferate vegetatively e of this product against a particular we eve, crop advisor, or extension agent to t this product specified for your local co capes with an herbicide having a mode ty be identified by these indicators: by be identified by the herbicide at the dos to fa particular weed species; and dividuals of the same species. of the following herbicide resistance me er mode of action as a foundation in a es with alternative modes of action. iroup 3 herbicides.	Veed Resistance Management processes applied, especially if control is achieved an agement processes applied, especially if control is achieved an agement processes applied, especially if control is achieved an agement processes and weed sizes present at the specified time to minimize with a species. Ye are relatively small, the field at the specified time to minimize with a species. Ye are the specified time to minimize with a species. Ye are the specified time to minimize with a species. Ye are the specified time to minimize with a species. Ye are the specified time to minimize with a special conditions. Tank mix products so that there are of action other than Group 3 and/or use no assess applied, especially if control is achieved management practices as is practical; weed control program, if appropriate.	ent th dithiopyr is a Group 3 herbicide based the dithiopyr is a Group 3 herbicide based the perations and/or other cultural practices. Program. In the discovery of the county extension agent. In the discovery of the county extension agent. In the multiple effective modes of action for onchemical methods to remove escapes, on adjacent weeds; In the multiple effective modes of action for onchemical methods to remove escapes, on adjacent weeds;	d on the mode of action classification rersified weed management program may s. Research has demonstrated that using If resistant biotypes of target weeds have r each target weed. s, as practical, with the goal of preventing	fan palm, Mexican fem (various) fescue fetterbush ficus fir fraser forsythia fuchsia galium gardenia geranium gum hawthom heather, twisted hemlock, Canada hibiscus	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retus Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum Gardenia jasminoides Pelarganium X hortorum Eucalyptus citriodora Crataegus spp. Erica cinerea Tsuga canadensis Hibiscus spp.	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping mystery radicans cockspur white crimson cloud enchantress Jack Evans Washington white Mediterranean pink blue bird brilliant hula girl blue boy blue girl burfodii china girl compacta forsteri hellerie Japanese northem beauty needlepoint	privet, glossy pyracantha queen palm quince, Japanese † rhododendron, carolina rhododendron, catawba rockcress rhodie max (rosebay) rose † rosemary † rosemary, bog salvia sedum snapdragon sourwood spiraea spiraea spiraea spiraea spiraea spiraea spiraea spiraee spiraea spiraea spiraee	Potentilla fruticosa Ligustrum japonicum Ligustrum japonicum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron carolinianum Rhododendron spp Rhododendron catawbiense Arabis spp. Rhododendron maximum Rosa banksiae Rosmarinus officinalis Andromeda polifolia Salvia farinacea Sedum spurium Antirrhinum spp. Oxydendrum arboreum Astilbe X arendsii Spiraea spp. Picea glauca Picea pungens Picea glauca Picea glauca Liquidambar styraciffua Platanus occidentalis Asparagus virgatus	golden vicary regal texanum wax yellow tipped gnome lalandei victory um album Cunningham white fashion hardy pim purple gem silvery pink snowcap luta nana rhea dragon blood red red carpet yellow fanall Anthony waterer red dolchica froebeli pink goldenflame red snowmound white Van Houtte white
bluegrass, annual Poa ai bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capses Lebanon ProScape Fertilizer with Dimensic system of the Weed Science Society of An Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this product To aid in the prevention of developing wee Scout fields before application to ensure 1 Start with a clean field, using either a burn of the Weed Science Society of An Start with a clean field, using either a burn of the Weed Science Society of An Start with a clean field, using either a burn of the Weed Science Society of An Start with a clean field, using either a burn of the Weed Science Society of An Start with a clean field, using either a burn of the Weed Science of Inspection to detect we Control weed escapes before they reproduction. Contact your local company representative of the seriod of the Weed Science of Inspection	minua umine spp. ria spp. lim carolinianum milla arvensis saria matricarioides sella bursa-pastoris Won® 0.15% FG turf and ornamental he nerica. control strategies to minimize selection different modes of action and overlapp nportant to delay the selection for resis et depends on the successful implement et series stant to this product, users shot herbicides and rates will be appropriat mdown herbicide application or tillage, k mixes, control weeds early when the or the most difficult to control weed in the et descapes or shifts in control of weed duce by seed or proliferate vegetatively e of this product against a particular we et, crop advisor, or extension agent to t this product specified for your local co- capes with an herbicide having a mode ty be identified by these indicators: by to including the same species, and dividuals of the same species. of the following herbicide resistance me er mode of action as a foundation in a rea with alternative modes of action. iroup 3 herbicides. spplications of Lebanon ProScape Fert h a different mode of action with an ow ractices, such as mechanical cultivatio nhance crop development and crop coi equipment before leaving fields suspe-	Veed Resistance Management processes and weed resistance management processes and weed sizes presented at the specified at the specified time to minimize we despected to your local company representative, lotted to this onditions. Tank mix products so that there are of action other than Group 3 and/or use not see applied, especially if control is achieved management practices as is practical: weed control program, if appropriate.	ent th dithiopyr is a Group 3 herbicide based the dithiopyr is a Group 3 herbicide based the perations and/or other cultural practices. Program. In the discovery of the county extension agent. In the discovery of the county extension agent. In the multiple effective modes of action for onchemical methods to remove escapes, on adjacent weeds; In the multiple effective modes of action for onchemical methods to remove escapes, on adjacent weeds;	d on the mode of action classification rersified weed management program may s. Research has demonstrated that using If resistant biotypes of target weeds have r each target weed. s, as practical, with the goal of preventing	fan palm, Mexican ferm (various) fescue fetterbush ficus fir fraser forsythia fuchsia galium gardenia geranium gum hawthorn heather, twisted hemlock, Canada hibiscus holly, Chinese	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum Gardenia jasminoides Pelargonium X hortorum Eucalyptus citriodora Crataegus spp. Erica cinerea Tsuga canadensis Hibiscus spp. Ilex Spp. Ilex X meserveae	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping mystery radicans cockspur white crimson cloud enchantress Jack Evans Washington white Mediterranean pink blue bird brilliant hula girl blue bory blue girl burdodii china girl compacta forsteri hellerie Japanese northem beauty needlepoint Nellie R. Stevens Savannah	privet, glossy pyracantha queen palm quince, Japanese † rhododendron, carolina rhododendron, catawba rockcress rhodie max (rosebay) rose † rosemary † rosemary † rosemary, bog salvia sedum snapdragon sourwood spiraea spiraea spiraea spiraea, garland spruce, lolorado blue spruce, colorado blue spruce, norway spruce, white sweetflag, grassyy-le sweetgum sycamore tree fem (tiki fem) trumpet flower, evening tulip	Potentilla fruticosa Ligustrum japonicum Ligustrum japonicum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron spp Rhododendron spp Rhododendron catawbiense Arabis spp. Rhododendron maximum Rosa banksiae Rosmarinus Oficinalis Andromeda polifolia Salvia farinacea Sedum spurium Antirrhinum spp. Oxydendrum arboreum Astilbe X arendsii Spiraea spp. Picea glauca Picea pungens Picea glauca Picea glauca Liquidambar styraciffua Platanus occidentalis Asparagus virgatus Gelsemium sempervirens Tulip, spp	golden vicary regal texanum wax yellow tipped gnome lalandei victory album Cunningham white fashion hardy pim purple gem silvery pink snowcap luta nana rhea dragon blood red red carpet yellow fanall Anthony waterer red dolchica froebeli pink goldenflame red snowmound white Van Houtte white
bluegrass, annual Poa at bluegrass, annual Poa at bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capse Pertilizer with Dimensic system of the Weed Science Society of An Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this produc To aid in the prevention of developing wee Scout fields before application to ensure I start with a clean field, using either a bur of If using post-emergence herbicides or tar Apply full rates of Dimension 0.15% FG for Scout fields after application to detect we Control weed escapes before they reproc Report any incidence of non-performance Contact your local company representative Deen they control weed escapes before they reproced the start with a classifield, using either a bur Scout fields after application to detect we Control weed escapes before they reproced in the start with a classifield, as they applicated they are control as the supplication rates of If resistance is suspected, treat weed secturther seed production. Suspected herbicide-resistant weeds may Failure to control a weed species normall A spreading patch of non-controlled plant Surviving plants mixed with controlled in Surviving plants mixed with controlled in Rotate the use of this product with non-Canton Avoid making more than two sequential aseason unless mixed with an herbicide with loncoprate non-hemical weed control pincy user species that even with an herbicide with loncoprate non-hemical weed control pincy user species that even decortrol pincy user species that even	minua umine spp. ria spp. lim carolinianum milla arvensis saria matricarioides sella bursa-pastoris Won® 0.15% FG turf and ornamental he nerica. control strategies to minimize selection different modes of action and overlapp nportant to delay the selection for resis et depends on the successful implement et series stant to this product, users shot herbicides and rates will be appropriat mdown herbicide application or tillage, k mixes, control weeds early when the or the most difficult to control weed in the et descapes or shifts in control of weed duce by seed or proliferate vegetatively e of this product against a particular we et, crop advisor, or extension agent to t this product specified for your local co- capes with an herbicide having a mode ty be identified by these indicators: by to including the same species, and dividuals of the same species. of the following herbicide resistance me er mode of action as a foundation in a rea with alternative modes of action. iroup 3 herbicides. spplications of Lebanon ProScape Fert h a different mode of action with an ow ractices, such as mechanical cultivatio nhance crop development and crop coi equipment before leaving fields suspe-	Veed Resistance Management processes and weed resistance management processes and weed sizes presented at the specified at the specified time to minimize we despected to your local company representative, lotted to this onditions. Tank mix products so that there are of action other than Group 3 and/or use not see applied, especially if control is achieved management practices as is practical: weed control program, if appropriate.	ent th dithiopyr is a Group 3 herbicide based the dithiopyr is a Group 3 herbicide based the perations and/or other cultural practices. Program. In the discovery of the county extension agent. In the discovery of the county extension agent. In the multiple effective modes of action for onchemical methods to remove escapes, on adjacent weeds; In the multiple effective modes of action for onchemical methods to remove escapes, on adjacent weeds;	d on the mode of action classification rersified weed management program may s. Research has demonstrated that using If resistant biotypes of target weeds have r each target weed. s, as practical, with the goal of preventing	fan palm, Mexican fern (various) fescue fescue fetterbush ficus fir fraser forsythia fuchsia galium gardenia geranium gum hawthorn heather, twisted hemlock, Canada hibiscus holly, Chinese holly, Japanese holly, yaupon	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum Gardenia jasminoides Pelargonium X hortorum Eucalyptus citriodora Crataegus spp. Erica cinerea Tsuga canadensis Hibiscus spp. Ilex spp. Ilex x meserveae	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping mystery radicans cockspur white crimson cloud enchantress Jack Evans Washington white Mediterranean pink blue bird brilliant hula girl blue boy blue girl burfodii china girl compacta forsteri hellerie Japanese northem beauty needlepoint Nellie R. Stevens Savannah	privet, glossy pyracantha queen palm quince, Japanese † rhododendron, carolina rhododendron, catawba rockcress rhodie max (rosebay) rose † rosemary † rosemary † rosemary, bog salvia sedum snapdragon sourwood spiraea spiraea spiraea spiraea, garland spruce, black hills spruce, Colorado blue spruce, norway spruce, white sweetflag, grassyy-le sweetgum sycamore tree fern (tiki fern) trumpet flower, evening	Potentilla fruicosa Ligustrum japonicum Ligustrum japonicum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron carolinianum Rhododendron catawbiense Arabis spp. Rhododendron maximum Rosa banksiae Rosmarinus officinalis Andromeda polifolia Salvia farinacea Sedum spurium Antirrhinum spp. Oxydendrum arboreum Astilbe X arendsii Spiraea spp. Picea glauca Picea pungens Picea glauca Liquidambar styraciflua Platanus occidentalis Asparagus virgatus Gelsemium sempervirens	golden vicary regal texanum wax yellow tipped gnome lalandei victory um album Cunningham white fashion hardy pim purple gem silvery pink snowcap luta nana rhea dragon blood red red carpet yellow fanall Anthony waterer red dolchica froebeli pink goldenflame red snowmound white Van Houtte white conica apeldoorn St. Paul
bluegrass, annual Poa ai bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capse Lebanon ProScape Fertilizer with Dimensi system of the Weed Science Society of An Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this produc To aid in the prevention of developing wee Scout fields before application to ensure 1 Start with a clean field, using either a burn of If using post-emergence herbicides or tar Apply full rates of Dimension 0.15% FG for Scout fields after application to detect we Control weed escapes before they reprod Report any incidence of non-performance Contact your local company representative been reported, use the application rates of 1f resistance is suspected, treat weed escributher seed production. Suspected herbicide-resistant weeds many control of the production of the productio	mnua umine spp. ria spp. lim carolinianum milla arvensis laria matricarioides laria matricarioides laria bursa-pastoris Won® 0.15% FG turf and ornamental he nerica. control strategies to minimize selection different modes of action and overlapp nportant to delay the selection for resis at depends on the successful implement of the selection of t	Veed Resistance Management perbicide, which contains the active ingredient on for weed populations resistant to one or no ping weed spectrum with or without tillage of stance. Intation of a weed resistance management public terror for the weed species and weed sizes preside the field at the specified time to minimize with dispecies. In the field at the specified time to minimize with species. In the field at the specified time to minimize with species. In the field at the specified time to minimize with species. In the field at the specified time to minimize with species. In the field at the specified time to minimize with species. In the field at the specified time to minimize with species. In the field at the specified time to minimize with species. In the field at the specified time to minimize with species. In the field at the specified time to minimize with species. In the field at the specified time to minimize with there are of action other than Group 3 and/or use no species. In the field at the specified time to minimize with the species. In the field at the specified time to minimize with the species. In the field at the specified time to minimize with the species. In the field at the specified time to minimize with the species. In the field at the specified time to minimize with the species. In the field at the specified time to minimize with the species. In the field at the specified time to minimize with the species. In the field at the specified time to minimize with the species. In the field at the species and weed sizes present the species. In the field at the species and weed sizes present the species and weed sizes present the species. In the field at the species and weed sizes present the species	ent It dithiopyr is a Group 3 herbicide based more herbicides is a best practice. A dive perations and/or other cultural practices. program. sent. eed escapes. ocal retailer, or county extension agent. Is MOA have been found in your region. I are multiple effective modes of action for onchemical methods to remove escapes, on adjacent weeds; omamental herbicide and any other Groud weeds. e crop seeds, as part of an integrated we uipment	d on the mode of action classification versified weed management program may s. Research has demonstrated that using s. Research has demonstrated that using s. If resistant biotypes of target weeds have reach target weed. s, as practical, with the goal of preventing sup 3 herbicides within a single growing reed control program.	fan palm, Mexican ferm (various) fescue fescue fetterbush ficus fir fraser forsythia fuchsia galium gardenia geranium gum hawthorn heather, twisted hemlock, Canada hilbiscus holly, Japanese holly, Japanese holly, Japanese holly, Jaupon honeysuckle	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum Gardenia jasminoides Pelargonium X hortorum Eucalyptus citriodora Crataegus spp. Erica cinerea Tsuga canadensis Hibiscus spp. Ilex x meserveae Ilex X attenuata Ilex corenata Ilex corenata Ilex corenata Ilex corenata Ilex comitoria Lonicera japonica	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping mystery radicans cockspur white crimson cloud enchantress Jack Evans Washington white Mediterranean pink blue bird brilliant hula girl blue bord brilliant hula girl blue bord brilliant hula girl blue bord solicia girl compacta forsteri hellerie Japanese northem beauty needlepoint Nelle R. Stevens Savannah claveyis dwarf halliana tatarian Canadian white	privet, glossy pyracantha queen palm quince, Japanese † rhododendron, carolina rhododendron, catawba rockcress rhodie max (rosebay) rose † rosemary † rosemary † rosemary , bog salvia sedum snapdragon sourwood spiraea spiraea spiraea spiraea spiraea spiraee spi	Potentilla fruticosa Ligustrum japonicum Ligustrum japonicum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron carolinianum Rhododendron catawbiense Arabis spp. Rhododendron maximum Rosa banksiae Rosmarinus officinalis Andromeda polifolia Salvia farinacea Sedum spurium Antirrhinum spp. Oxydendrum arboreum Astilbe X arendsii Spiraea spp. Picea glauca Picea pungens Picea abies Picea glauca Liquidambar styraciflua Platanus occidentalis Asparagus virgatus Gelsemium sempervirens Tulip, spp Lantana sellowiana Verbena spp.	golden vicary regal texanum wax yellow tipped gnome lalandei victory album Cunningham white fashion hardy pim purple gem silvery pink snowcap luta nana rhea dragon blood red red carpet yellow fanall Anthony waterer red dolchica froebeli pink goldenflame red snowmound white Van Houtte white conica apeldoorn St. Paul American cranberry bush arrowood common snowball
bluegrass, annual bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capse Lebanon ProScape Fertilizer with Dimensic system of the Weed Science Society of An Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this produc To aid in the prevention of developing wee Scout fields before application to ensure I Start with a clean field, using either a bun fif using post-emergence herbicides or tar Apply full rates of Dimension 0.15% FG fe Scout fields after application to detect we Control weed escapes before they reproc Report any incidence of non-performance Contact your local company representativ been reported, use the application rates of If resistance is susspected, treat weed esc further seed production. Suspected herbicide-resistant weeds may Failure to control a weed species normall A spreading patch of non-controlled plant Surviving plants mixed with controlled in Additionally, users should follow as many of Use a broad spectrum herbicide with ofte Surviving plants mixed with controlled in Additionally, users should follow as many of Use a broad spectrum herbicide with ofte Utilize sequential applications of herbicide Notate the use of this product with non-G Avoid making more than two sequential aseason unless mixed with an herbicide Incorporate non-chemical weed control p Use good agronomic principles that er Thoroughly clean plant residues from Manage weeds in and around fields to	mnua umine spp. ria spp. ria spp. m spp. mispp. miliam carolinianum milla arvensis saria matricarioides silla bursa-pastoris won® 0.15% FG turf and ornamental he merica. control strategies to minimize selection different modes of action and overlapp mortant to delay the selection for resis at depends on the successful implement eds resistant to this product, users shound herbicides and rates will be appropriate and own herbicide application or tillage. And mixes, control weeds early when the orthe most difficult to control weed in the ed escapes or shifts in control of weed use by seed or proliferate vegetatively e of this product against a particular we ve, crop advisor, or extension agent to this product specified for your local coapes with an herbicide having a mode by be identified by these indicators: by controlled by the herbicide at the dost so if a particular weed species; and dividuals of the same species. of the following herbicide resistance may remode of action as a foundation in a rese with alternative modes of action. It is a sea to the same species or educe weed seed production. Applic rotary-type spreaders designed to applanules in narrow rows or concentrated ractual applications. The derivatives and the same species in action in a result application conditions. The derivatives and the same species are seed to a polar actual application conditions. The derivatives and the same species in actual application conditions. The derivatives actual application conditions.	Veed Resistance Management perbicide, which contains the active ingredient on for weed populations resistant to one or no ping weed spectrum with or without tillage of stance. Intation of a weed resistance management public te for the weed species and weed sizes preside the field at the specified time to minimize with dispecies. In the field at the specified time to minimize with dispecies. In the field at the specified time to minimize with dispecies. In the field at the specified time to minimize with dispecies. In the field at the specified time to minimize with species. In the field at the specified time to minimize with species. In the field at the specified time to minimize with species. In the field at the specified time to minimize with species. In the field at the species and weed sizes are of action other than Group 3 and/or use not action other than Group 3 and/or use not species. In the field at the species with species are species as is practical: weed control program, if appropriate. It is the field at the species with species and species are coording to the species and species. In the field at the species are species and species are calibrated the species are according to the species and species action on setting may be marked or species action on setting may be marked or species.	ent It dithiopyr is a Group 3 herbicide based more herbicides is a best practice. A dive perations and/or other cultural practices. program. sent. eed escapes. ocal retailer, or county extension agent. Is MOA have been found in your region. I are multiple effective modes of action for onchemical methods to remove escapes, on adjacent weeds; ornamental herbicide and any other Group of the manufacturer's directions. Initial spreadows are on the manufacturer's directions. Initial spreadows or the manufacturer's directions or the manufacturer's directions. Initial spreadows or the m	d on the mode of action classification versified weed management program may s. Research has demonstrated that using If resistant biotypes of target weeds have r each target weed. s, as practical, with the goal of preventing pup 3 herbicides within a single growing reed control program.	fan palm, Mexican ferm (various) fescue fescue fetterbush ficus fir fraser forsythia fuchsia galium gardenia geranium gum hawthorn heather, twisted hemlock, Canada hibiscus holly, Chinese holly, Japanese holly, yaupon honeysuckle hosta ice plant impatiens	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum Gardenia jasminoides Pelarganium X hortorum Eucalyptus citriodora Crataegus spp. Erica cinerea Tsuga canadensis Hibiscus spp. Ilex X meserveae Ilex X meserveae Ilex cormuta Ilex cormuta Ilex cormuta Ilex cormuta Ilex cornuta Ilex cornuta Ilex cornuta Ilex corporotica Hosta spp. Hosta lancifolia Carpobrotus edulis Impatiens spp.	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping mystery radicans cockspur white crimson cloud enchantress Jack Evans Washington white Mediterranean pink blue bird brilliant hula girl blue boy blue girl burfodii china girl compacta forsteri hellerie Japanese northem beauty needlepoint Nellie R. Stevens Savannah claveyis dwarf halliana tatarian Canadian white zebelli red hosta albo marginata	privet, glossy pyracantha queen palm quince, Japanese † rhododendron, carolina rhododendron, catawba rockcress rhodie max (rosebay) rose † rosemary † rosemary † rosemary , bog salvia sedum snapdragon sourwood spiraea spiraea spiraea spiraea spiraea spiraee spi	Potentilla fruticosa Ligustrum japonicum Ligustrum japonicum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron carolinianum Rhododendron catawbiense Arabis spp. Rhododendron maximum Rosa banksiae Rosmarinus officinalis Andromeda polifolia Salvia farinacea Sedum spurium Antirrhinum spp. Oxydendrum arboreum Astilbe X arendsii Spiraea spp. Picea glauca Picea pungens Picea abies Picea glauca Liquidambar styraciflua Platanus occidentalis Asparagus virgatus Gelsemium sempervirens Tulip, spp Lantana sellowiana Verbena spp.	golden vicary regal texanum wax yellow tipped gnome lalandei victory um album Cunningham white fashion hardy pim purple gem silvery pink snowcap luta nana rhea dragon blood red red carpet yellow fanill Anthony waterer red dolchica froebeil pink goldenflame red snowmound white Van Houtte white conica speldoorm speldoorm speldoord common snowball European cranberry bush linden
bluegrass, annual Poa al bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capse Lebanon ProScape Fertilizer with Dimensic system of the Weed Science Society of An Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this produc To aid in the prevention of developing wee Scout fields before application to ensure I Start with a clean field, using either a bun of If using post-emergence herbicides or tar Apply full rates of Dimension 0.15% FG for Scout fields after application to detect we Control weed escapes before they reproduce the Control weed escapes before they reproduced to the Control weed escapes before they reproduced the Control weed escapes before they reproduced the Control weed species normall A spreading patch of non-performance Contact your local company representative esc further seed production. Suspected herbicide-resistant weeds may be a production of the Control weed section to the Control weed section of	minua umine spp. ria spp. ria spp. lim	Jeed Resistance Management procedure, which contains the active ingredient on for weed populations resistant to one or no ping weed spectrum with or without tillage of stance. Intation of a weed resistance management procedure in the field at the specified time to minimize with did at the specified time to minimize with gradient of the species. In the field at the specified time to minimize with gradient of the species. In the field at the specified time to minimize with gradient of the species. In the field at the specified time to minimize with gradient of the species. In the field at the specified time to minimize with gradient of the species. In the field at the specified time to minimize with gradient of the species. In did out if suspected resistant weeds to this onditions. Tank mix products so that there are of action other than Group 3 and/or use not one applied, especially if control is achieved management practices as is practical: weed control program, if appropriate. It is a species of the first properties of the species	ent It dithiopyr is a Group 3 herbicide based more herbicides is a best practice. A dive perations and/or other cultural practices. program. Issent. In the discovery of the county extension agent. Is MOA have been found in your region. I are multiple effective modes of action for onchemical methods to remove escapes, on adjacent weeds; In the discovery of the county and uniformly avour of the manufacturer's directions. Initial sprecorded for future reference. Apply this recorded for future reference. Apply this is recorded for future reference. Apply this is retreatment area and then applying the tereatment area and then applying the second of the county of the c	d on the mode of action classification versified weed management program may s. Research has demonstrated that using If resistant biotypes of target weeds have r each target weed. s, as practical, with the goal of preventing pup 3 herbicides within a single growing reed control program.	fan palm, Mexican fem (various) fescue fetterbush ficus fir fraser forsythia fuchsia galium gardenia geranium gum hawthom heather, twisted hemlock, Canada hibiscus holly, Chinese holly, Japanese holly, yaupon honeysuckle hosta ice plant impatiens iris	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum Gardenia jasminoides Pelargonium X hortorum Eucalyptus citriodora Crataegus spp. Erica cinerea Tsuga canadensis Hibiscus spp. Ilex Spp. Ilex X meserveae Ilex Comuta Ilex comuta Ilex comuta Ilex comuta Ilex comuta Ilex comuta Ilex omitoria Lonicera japonica Hosta spp. Hosta lancifolia Carpobrotus edulis	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping mystery radicans cockspur white crimson cloud enchantress Jack Evans Washington white Mediterranean pink blue bird brilliant hula girl blue bory blue girl burfodii china girl compacta forsteri hellerie Japanese northem beauty needlepoint Nellie R. Stevens Savannah claveyis dwarf halliana tatarian Canadian white zebelli red hosta albo marginata	privet, glossy pyracantha queen palm quince, Japanese † rhododendron, carolina rhododendron, catawba rockcress rhodie max (rosebay) rose † rosemary † rosemary † rosemary , bog salvia sedum snapdragon sourwood spiraea spiraea spiraea spiraea spiraea spiraee spi	Potentilla fruticosa Ligustrum japonicum Ligustrum japonicum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron carolinianum Rhododendron catawbiense Arabis spp. Rhododendron maximum Rosa banksiae Rosmarinus officinalis Andromeda polifolia Salvia farinacea Sedum spurium Antirrhinum spp. Oxydendrum arboreum Astilbe X arendsii Spiraea spp. Picea glauca Picea pungens Picea abies Picea glauca Liquidambar styraciflua Platanus occidentalis Asparagus virgatus Gelsemium sempervirens Tulip, spp Lantana sellowiana Verbena spp.	golden vicary regal texanum wax yellow tipped gnome lalandei victory um album Cunningham white fashion hardy pim purple gem silvery pink snowcap luta nana rhea dragon blood red red carpet yellow fanall Anthony waterer red dolchica froebeli pink goldenflame red snowmound white Van Houtte white conica apeldoorn St. Paul American cranberry bush arrowood common snowball European cranberry bush arrowood common snowball European cranberry bush arrowood
bluegrass, annual bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Jarolice	imine spp. ria spp. lium carolinianum milla arvensis saria matricarioides sella bursa-pastoris Won® 0.15% FG turf and ornamental he nerica. control strategies to minimize selection different modes of action and overlapp nportant to delay the selection for resis et depends on the successful implement eds resistant to this product, users shoutherbicides and rates will be appropriat modown herbicide application or tillage, k mixes, control weeds early when the or the most difficult to control weed in the ed escapes or shifts in control of weed use by seed or proliferate vegetatively e of this product against a particular we ev, crop advisor, or extension agent to this product specified for your local co capes with an herbicide having a mode by be identified by these indicators: by controlled by the herbicide at the dot sto of a particular weed species; and dividuals of the same species. of the following herbicide resistance me er mode of action as a foundation in a rese with alternative modes of action. iroup 3 herbicides. applications of Lebanon ProScape Fert ha different mode of action with an ow ractices, such as mechanical cultivation hance crop development and crop co requipment before leaving fields suspen- or reduce weed seed production. Applic rotary-type spreaders designed to appl anules in narrow rows or concentrated or actual application conditions. The de isually be achieved by applying one-ha tion). Avoid streaking, skips, or overla	Veed Resistance Management particide, which contains the active ingredient on for weed populations resistant to one or not ping weed spectrum with or without tillage of stance. Intation of a weed resistance management pandl: It for the weed species and weed sizes preside the field at the specified time to minimize with dispecies. It indout if suspected resistant weeds to this orditions. Tank mix products so that there are of action other than Group 3 and/or use not seed applied, especially if control is achieved management practices as is practical: weed control program, if appropriate. It ilizer with Dimension® 0.15% FG turf and correlapping spectrum for the difficult-to-contron, crop rotation, cover crops and weed-free propetitiveness. Sected to contain resistant weeds. Cation Instructions and Equal by granular herbicides. For best results, apt be bands. Calibrate the spreader according testing may be marked or all of the required amount of product over the apsiduring application. Tections for Landscape Orrections.	ent It dithiopyr is a Group 3 herbicide based more herbicides is a best practice. A dive perations and/or other cultural practices. program. sent. eed escapes. ocal retailer, or county extension agent. Is MOA have been found in your region. I are multiple effective modes of action for onchemical methods to remove escapes, on adjacent weeds; ornamental herbicide and any other Group of weeds. In crop seeds, as part of an integrated we uipment ply this product evenly and uniformly avo to the manufacturer's directions. Initial sprecorded for future reference. Apply this recorded for future reference.	d on the mode of action classification versified weed management program may s. Research has demonstrated that using s. Resear	fan palm, Mexican fem (various) fescue fetterbush ficus fir fraser forsythia fuchsia galium gardenia geranium gum hawthom healther, twisted hemlock, Canada hibiscus holly, Chinese holly, Japanese holly, yaupon honeysuckle hosta ice plant impatiens iris ivy, English jasmine, Asian	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum Gardenia jasminoides Pelargonium X hortorum Eucalyptus citriodora Crataegus spp. Erica cinerea Tsuga canadensis Hibiscus spp. Ilex X meserveae Ilex X meserveae Ilex comuta Ilex comuta Ilex comuta Ilex crenata Ilex comita Ilex crenata Ilex omitoria Lonicera japonica Hosta spp. Hosta lancifolia Carpobrotus edulis Impatiens spp. Iris spp. Iris spp. Hedera helix Trachelospermum asiaticum	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping mystery radicans cockspur white crimson cloud enchantress Jack Evans Washington white Mediterranean pink blue bird brilliant hula girl blue bory blue girl burfodii china girl compacta forsteri hellerie Japanese northem beauty needlepoint Nellie R. Stevens Savannah claveyis dwarf halliana tatarian Canadian white zebelli red hosta albo marginata dwarf blue wedgewood bulgaria thomdale	privet, glossy pyracantha queen palm quince, Japanese † rhododendron, carolina rhododendron, catawba rockcress rhodie max (rosebay) rose † rosemary † rosemary, bog salvia sedum snapdragon sourwood spiraea spiraea spiraea spiraea, garland spruce, lolorado blue spruce, colorado blue spruce, norway spruce, white sweetflag, grassyy-le sweetgum sycamore tree fem (tiki fem) trumpet flower, evening tulip verbena, shrub vervain viburnum	Potentilla fruticosa Ligustrum Japonicum Ligustrum Japonicum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron spp Rhododendron spp Rhododendron maximum Rhododendron maximum Rosa banksiae Rosmarinus officinalis Andromeda polifolia Salvia farinacea Sedum spurium Antirrhinum spp. Oxydendrum arboreum Astilbe X arendsii Spiraea spp. Picea glauca Picea pungens Picea glauca Picea glauca Liquidambar styraciffua Platanus occidentalis Asparagus virgatus Gelsemium sempervirens Tulip, spp Lantana seltowiana Verbena spp. Viburnum spp. Vinca minor Trachycarpus fortunei Xylosma congestum	golden vicary regal texanum wax yellow tipped gnome lalandei victory um album Cunningham white fashion hardy pim purple gem silvery pink snowcap luta nana rhea dragon blood red red carpet yellow fanall Anthony waterer red dolchica froebeli pink goldenflame red snowmound white Van Houtte white conica apeldoorn st. Paul American cranberry bush arrowcod common snowball European cranberry bush linden Mohican
bluegrass, annual Poa al bittercress Carda chickweed Stellar henbit Lamiu geranium, Carolina Geran parsley-piert Alcher pineappleweed Matric shepherdspurse Capse Lebanon ProScape Fertilizer with Dimensic system of the Weed Science Society of An Proactively implementing diversified weed include the use of multiple herbicides with the labeled rate and directions for use is in The continued effectiveness of this produc To aid in the prevention of developing wee Scout fields before application to ensure I Start with a clean field, using either a bur of Start with a clean field, using either a bur of I start with a clean field, using either a bur of I start with a clean field, using either a bur of I start with a clean field, using either a bur of I start with a clean field, using either a bur of I start with a clean field, using either a bur of I start with a clean field, using either a bur of I start with a clean field, using either a bur of I start with a clean field, using either a bur of I start with a clean field, using either a bur of I start with a clean field, using either a bur of I start with a clean field, using either a bur of Start with a clean field, using either a bur of I start with a clean field, using either a bur of I start with a clean field, using either a bur of I start with a clean field, using either a bur of I start with a clean field, using either a bur of I start with a start was a start with a start weed esc further seed production. Suspected herbicide-resistant weeds man of I resistance is suspected, treat weed esc further seed production. Suspected herbicide-resistant weeds man of I resistance is suspected, treat weed sec further seed production. Suspected herbicide-resistant weeds man of I resistance is suspected, treat weed esc further seed production. Suspected herbicide resistant weeds man of I resistance is suspected, treat weed esc further seed production. Suspected herbicide resistant weed sommal of the security of I resistance is suspected, treat weed esc further seed prod	imine spp. ria spp. imine spp. ria spp. imine arolinianum milla arvensis aria matricarioides alla bursa-pastoris Won® 0.15% FG turf and ornamental he nerica. control strategies to minimize selection different modes of action and overlapp nportant to delay the selection for resis at depends on the successful implement add resistant to this product, users shoutherbicides and rates will be appropriat modown herbicide application or tillage. In mixes, control weeds early when the or the most difficult to control weed in the add resistant to this product, users shoutherbicides and rates will be appropriat modown herbicide application or tillage. In this product against a particular we tor, crop advisor, or extension agent to this product specified for your local co capes with an herbicide having a mode ty be identified by these indicators: by be identified by the herbicide at the dos capes with an herbicide having a mode ty controlled by the herbicide at the dos to a particular weed species; and dividuals of the same species. of the following herbicide resistance may er mode of action as a foundation in a rese with atternative modes of action. Toroup 3 herbicides. of the following herbicide resistance may er mode of action as a foundation in a resolution of the same species. of the following herbicide resistance may er mode of action as a foundation in a resolution of the same species. of the following herbicide resistance may er mode of action as foundation in a resolution of the same species. of the following herbicide resistance may er mode of action as a foundation in a resolution of the same species. of the production of the same species of action. The descapes with an herbicide and action with an overactices, such as mechanical cultivation hance crop development and crop concentrated and a different mode of action with an overactices, such as mechanical cultivation hance crop development and crop concentrated and a splication conditions. The de susually be achieved b	Veed Resistance Management processes applied, especially if control is achieved to reaction other than Mirror program, if appropriate. Titilizer with Dimension® 0.15% FG turf and correlapping spectrum for the difficult-to-control, crop rotation, cover crops and weed-free impeditiveness. Sected to contain resistant weeds.	ent It dithiopyr is a Group 3 herbicide based more herbicides is a best practice. A dive perations and/or other cultural practices. program. sent. eed escapes. coal retailer, or county extension agent. Is MOA have been found in your region. I are multiple effective modes of action for onchemical methods to remove escapes, on adjacent weeds; commamental herbicide and any other Group of the manufacturer's directions. Initial sprecorded for future reference. Apply this retreatment area and then applying the tenamentals set of ormamentals gs of ormamental plants listed on this labe	d on the mode of action classification versified weed management program may s. Research has demonstrated that using s. Research has demonstrated that using s. If resistant biotypes of target weeds have reach target weed. s, as practical, with the goal of preventing support of the program. Total program. Total program. Total program adjustment to be preventing one-half in a different direction bel. This product may be applied in	fan palm, Mexican ferm (various) fescue fetterbush ficus fir fraser forsythia fuchsia galium gardenia geranium gum hawthorn heather, twisted hemlock, Canada hibiscus holly, Chinese holly, Japanese	Washingtonia robusta Asparagus spp. Festuca glauca Leucothoe fontanesiana Ficus retusa Abies fraseri Forsythia X intermedia Spring Glory Fuchsia spp. Galium ordoratum Gardenia jasminoides Pelargonium X hortorum Eucalyptus citriodora Crataegus spp. Erica cinerea Tsuga canadensis Hibiscus spp. Ilex X meserveae Ilex X meserveae Ilex comuta Ilex corenata Ilex corenata Ilex comuta Ilex cranata Ilex crana	gold edge gold princess silver king tricolor vegetus rainbow nitidia Arnold dwarf bronxensis dwarf lynwood gold meadowlark weeping mystery radicans cockspur white crimson cloud enchantress Jack Evans Washington white Mediterranean pink blue bird brilliant hula girl blue boy blue girl burfodii china girl compacta forsteri hellerie Japanese northem beauty needlepoint Nellie R. Stevens Savannah claveyis dwarf halliana tatarian Canadian white zebelli red hosta albo marginata dwarf blue wedgewood bulgaria	privet, glossy pyracantha queen palm quince, Japanese † rhododendron, carolina rhododendron, catawba rockcress rhodie max (rosebay) rose † rosemary † rosemary † rosemary, bog salvia sedum snapdragon sourwood spiraea spiraea spiraea, garland spruce, Colorado blue spruce, norway spruce, white sweetlag, grassyy-le sweetgum sycamore tree fem (tiki fern) trumpet flower, evening tulip verbena, shrub vervain viburnum	Potentilla fruticosa Ligustrum japonicum Ligustrum japonicum Pyracantha koidzumii Arecastrum rammanzoffian Rhododendron spp Rhododendron spp Rhododendron catawbiense Arabis spp. Rhododendron maximum Rosa banksiae Rosmarinus officinalis Andromeda polifolia Salvia farinacea Sedum spurium Antirrhinum spp. Oxydendrum arboreum Astilbe X arendsii Spiraea spp. Picea glauca Picea pungens Picea pungens Picea glauca Liquidambar styraciffua Platanus occidentalis Asparagus virgatus Gelsemium sempervirens Tulip, spp Lantana seltowiana Verbena spp. Viburnum spp. Vinca minor Trachycarpus fortunei	golden vicary regal texanum wax yellow tipped gnome lalandei victory album Cunningham white fashion hardy pim purple gem silvery pink snowcap luta nana rhea dragon blood red red carpet yellow fanil Anthony waterer red dolchica froebeli pink goldenflame red snowmound white Van Houtte white conica apeldoorn St. Paul American cranberry bush arrowood common snowball European cranberry bush linden Mohican

Terms and Conditions of Use If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of

purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the

Limitation of Remedies

directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by Buye

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to at Seller's election, one of the following:

1.Refund of purchase price paid by buyer or user for product bought, or 2.Replacement of amount of product used. To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in

writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses. The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner