PRODIAMINE | **GROUP 3** | **HERBICIDE**

LEBANON ProScape®

FERTILIZER WITH BARRICADE 0.38% PREEMERGENCE WEED CONTROL

16-0-8 WITH MESA®

0 88685 53856 8

22-53856

GUARANTEED ANALYSIS

Total Nitrogen (N) **16%**
- 2.4% Ammoniacal Nitrogen
- 1.5% Water Insoluble Nitrogen*
- 9.6% Urea Nitrogen
- 2.5% OtherWater Insoluble Nitrogen*

Soluble Potash (K_2O) **8%**

Sulfur (S) 2.8%
- 2.8% Combined Sulfur (S)

Derived From: Ammonium Sulfate, Methylene Ureas, Urea. Muriate of Potash.
*4.0% Slowly available nitrogen from Methylene Ureas.

Chlorine (Cl) not more than 7.0%

F699

SUGGESTED SPREADER SETTINGS

Spreader	3.9#/1,000 sq. ft.	4.6#/1,000 sq. ft.	5.9#/1,000 sq. ft.	Spreader	3.9#/1,000 sq. ft.	4.6#/1,000 sq. ft.	5.9#/1,000 sq. ft.
LebanonTurf	4½	4	7	Lesco (letter/Numeric)	i/17	J/21	M/26
Andersons AccuPro	K	M	S	PennMulch HVO	L	N	S
Earthway Rotary	15	16	23	ProScape SS	L	N	S
Gandy	26	28	35	Spyker	4½	4	7
Lely	5 II	6 II	7½ II	Vicon (all models)	24	26	32

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Information regarding the contents and levels of metals in this product is available on the Internet at:
http://www.regulatory-info-lebsea.com

ACTIVE INGREDIENT:
Prodiamine 0.38%
OTHER INGREDIENTS: 99.62%
TOTAL: 100.00%

(L)

EPA Reg. No. 961-365 EPA Est. No. 961-PA-01:
Superscript used is first letter of lot number.

For selective preemergence control of listed grass and broadleaf weeds as listed in:
- **established turfgrasses (excluding golf course putting greens) and lawns**
- **landscape ornamentals (including established perennials & wildflower plantings)**

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.
Before buying this product, read the Warranty Section elsewhere on this label.
If terms are not acceptable, return the unopened container to the place of purchase at once.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

FIRST AID

IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **(888)-208-1368** for emergency medical treatment information.	

Net Weight 1000 LBS. (453.6 kg)

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. **DO NOT** apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. **DO NOT** contaminate water when disposing of equipment washwaters.
If material is spilled: Contain and sweep up material of spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants not listed on this label and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds as listed in established turfgrasses (excluding golf course putting greens) and lawns, landscape ornamentals, established perennials and wildflower plantings.
This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days following application.
Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

DO NOT graze or feed livestock forage cut from areas treated with this product.
DO NOT apply this product through any type of irrigation system.
DO NOT apply aerially.
DO NOT apply to golf course putting greens.

WEED RESISTANCE MANAGEMENT

For resistance management, **Lebanon ProScape Fertilizer with BARRICADE 0.38% Preemergence Weed Control** is a Group 3 herbicide. Any weed population may contain certain or develop plants naturally resistant to **Lebanon ProScape Fertilizer with BARRICADE 0.38% Preemergence Weed Control** and other Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.
To delay herbicide resistance take one or more of the following steps:
Rotate the use of **Lebanon ProScape Fertilizer with BARRICADE 0.38% Preemergence Weed Control** or other Group 3 herbicides within a growing season with different herbicide groups that control the same weeds.
Adopt an integrated weed management program for herbicide use that includes scouting and uses historical information related to herbicide use, and that considers cultural, biological varieties and other turf management practices.
Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.
If weed population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
Contact your sales representative, pest control advisors, or local extension specialist for additional pesticide resistance management and/or integrated weed management recommendations for specific types of plants and weed biotypes.

APPLICATION DIRECTIONS

Apply uniformly with suitable, calibrated application equipment.

ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds as listed in established turfgrasses and lawns. The maximum amount of this product may be applied per year is given for each turfgrass species in the **Maximum Annual Rates** section of this label.
Most effective weed control in turfgrasses will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

CRABGRASS SEED GERMINATION DATES

- **Approximate Date**
- **After May 30**
- **After May 10**
- **After April 20**
- **After March 20**
- **January 1 to March 20**

USE PRECAUTIONS

The following precautions apply to the use of this product in turfgrasses and lawns:
Application of this product may thin emerged annual bluegrass and newly overseeded grasses.
DO NOT apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
DO NOT apply to newly set sod until the following year.
Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.
Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.
DO NOT apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds.
Maximum use rate selection must be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES

This product is for use on the turfgrass species listed in the following table. DO NOT exceed the maximum yearly rate as given in the following table:

MAXIMUM APPLICATION RATE/CALENDAR YEAR of Lebanon ProScape FERTILIZER WITH BARRICADE 0.38% PREEMERGENCE WEED CONTROL by Turfgrass Species[1)]

Turf Species:	Lbs. Product/Acre	Lbs. Product/1,000 Sq. Ft.	Lbs. AI/Acre
Bermudagrass[2)], Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turf-type), Zoysia	395	8.9	1.5
Buffalograss, Kentucky Bluegrass, Perennial Ryegrass	265	5.9	1.0
Creeping Red Fescue	195	4.6	0.75
Creeping Bentgrass	170	3.9	0.65

[1)] These are the maximum rates per calendar year by species limitations.
[2)] May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 137 lbs./Acre. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.
- **DO NOT** apply more than 395 lbs. of this product per calendar year.
- Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but **DO NOT** exceed the maximum application rates specified in the **Maximum Annual Rates Table**.

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:

Barnyardgrass	Crabgrass (large, smooth)	Henbit	Kochia	Purslane, Common	Speedwell, Persian
Bluegrass, Annual (*Poa annua*)	Crowfootgrass	Itchgrass	Lambsquarters, Common	Pusley, Florida	Sprangletop
Carpetweed	Cupgrass, Woolly	Johnsongrass (from seed)	Lovegrass	Rescuegrass[3]	Spurge, Prostrate
Chickweed, Common	Foxtail, Annual	Junglerice	Panicum (Texas, Fall, Browntop)	Shepherd's Purse[2]	Witchgrass
Chickweed, Mouseear (from seed)	Goosegrass[1]	Knotweed	Pigweed	Signalgrass, Broadleaf	Woodsorrel, Yellow (from seed)

[1] In many areas a single application of 170 to 395 lbs./Acre of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 170 to 265 lbs./Acre followed after 60 to 90 days by a second application at doses that would not exceed those given in the **Maximum Annual Rate Table**. **DO NOT** exceed the maximum rate for turfgrass species listed in the **Maximum Annual Rate Table** above.
[2] Apply this product in late summer, fall or winter prior to germination.
[3] Suppression only.
Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates listed for each turf species. All applications must be made prior to germination of the weed seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF

DO NOT apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

Lebanon ProScape FERTILIZER WITH BARRICADE 0.38%

PREEMERGENCE WEED CONTROL Rate (lb./A)		Months Before Overseeding		
Product	AI	North	Transition	South
132	.50	4	4	4
170	.66	5	4	4
195	.74	6	5	5
210	.80	---	6	6
265	1.00	---	7	7
300	1.14	---	---	9
342	1.30	---	---	10
395	1.50	---	---	12

LANDSCAPE ORNAMENTALS (including Established Perennials and Wildflower Plantings)

This product may be applied for residual preemergence weed control in ornamentals.

USE RATES

Apply this product at 170 to 395 lbs./Acre in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications are to be made so long as the total amount of product applied does not exceed the maximum annual application rate of 395 lbs./Acre.

APPLICATION TIMING AND INFORMATION

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

ESTABLISHED LANDSCAPE ORNAMENTALS

Apply this product in the fall and/or spring to established ornamentals as a broadcast, over the top or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

NEWLY TRANSPLANTED LANDSCAPE ORNAMENTALS

Apply this product in the fall and/or spring to newly transplanted ornamentals as a broadcast, over the top or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

TOLERANT ORNAMENTAL SPECIES

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. Apply this product over the top of these species.
When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

Scientific Name	Common Name	Scientific Name	Common Name
Abies spp.	Fir species** (Balsam, Fraser, Noble, etc.)	*C. Dammeri*	Bearberry Cotoneaster
Abelia grandiflora	Abelia: Sherwood	*C. horizontalis*	Rockspray Cotoneaster
Acer palmatum	Japanese Maple	*C. microphyllus*	Littleleaf Cotoneaster
A. platanoides	Norway Maple	*Cretaegus* spp.	Hawthorn
Achillea sp.	Yarrow: King Edward	*Crocosmia* sp.	Lucifer
Actinidia chinensis	Kiwi*	*Cupressus sempervirens*	Italian Cypress
Agapanthus africanus	Lily-of-the-Nile (African Lily)	*Delosperma alba*	White Trailing Ice Plant
Agapanthus orientalis		*Delosperma* sp.	Cooperi Pink
Akebia quinata	Five-Leaf or Chocolate Vine	*Delphinium* sp.	Larkspur: Blue Elf
Allium cernuum	Lady's Leek, Nodding Onion	*Dianthus deltoides*	Dianthus; Maiden Pinks "Zing"
Anemone hybrida	Japanese Anemone	*D. gratianopolitanus*	Cheddar Pink
Aquilegia sp.	Aquilegia: Red and Gold	*Dodonaea viscosa*	Hop Bush
Arctostaphylos densiflora	Vine Hill Manzanita	*Echinacea pupurea*	Coneflower, Purple; Magnus
Arctotheca calendula	Cape Weed	*Elaeagnus commutata*	Silverberry
Artemesia sp.	Wormwood; Silver Mound, Castle	*E. pungens*	Silverthorn
Aster sp.	Aster: Bonny Blue, Purple Dome	*Euonymus Fortunei*	Wintercreeper
Aster X Frikartii	Frikarti's aster	*E. japonica*	Japanese Spindle Tree (Evergreen Euonymus,Spreading Euonymus)
Athyrium Filix-femina	Lady Fern; Fern Lady		Creeping Strawberry Bush; Manhattan Euonymus
Aucuba japonica	Japanese Aucuba	*E. kiautschovica*	Japanese Aralia
Begonia sp.	Fibrous Begonia: Hardy Grandis		Border Forsythia
Berberis gladwynensis	William Penn Barberry	*Fatsia japonica*	Weeping Forsythia
B. Julianae	Wintergreen Barberry	*Forsythia x intermedia*	Greenstem Forsythia
B. mentorensis	Mentor Barberry	*Forsythia suspensa*	Gaillardia, Blanket Flower: "Goblin"
B. Thunbergii	Japanese Barberry	*F. viridissima*	Gardenia, Cape-Jasmine, Radicans
B. verruculosa	Warty Barberry	*Gaillardia* sp.	
Bergenia cordifolia	Pigsqueak	*Gardenia jasminoides*	Gentian
Boltonia asteroides	Snowbank; False Aster	*Gaura* sp.	Ashey Cranesbill
Buddleia davidii	Butterfly-Bush (Dwarf Blue); Royal Red	*Gentiana dahurica*	Gladiolus species**
		Geranium cinereum	Baby's Breath
Buxus microphylla	Japanese Boxwood	*Gladiolus* spp.	English Ivy
Callistemon viminalis	Weeping Bottlebrush	*Gypsophila repens*	Sunrose
Calluna vulgaris	Scotch Heather	*Hedera helix*	Daylily: Aztec Gold, Stella De Oro, Tender Love
Campanula carpatica	Tussock Bellflower, (White Clips)	*Helianthemum* sp.	Rose of Sharon**
Campis X Tagliabuana	Trumpet Creeper, Trumpet Flower, Madame Galen	*Hemerocallis* sp.	Mallow: Disco Belle White
		Hibiscus syriacus	Chinese Hibiscus
Carpobrotus edulis	Hottentot Fig (Ice Plant)	*Hibiscus* sp.	Hosta, Plantain Lily (Fragrant)
Cassia artemisoides	Feathery Cassia	*Hibiscus rosa-sinensis*	Hosta, "Seersucker"
Ceanothus rigidus	Wild Lilac	*Hosta plantaginea*	Chameleon Plant
Ceratostigma plumbaginoides	Hardy Plumbago	*H. Sieboldiana*	Bigleaf Hydrangea
Chamaecyparis pisifera	False Cypress	*Houttuynia cordata var. variegata*	Perny Holly
Chrysanthemum nipponicum	Nippon Daisy	*Hydrangea macrophylla*	Chinese Holly
Cleyera japonica	Cleyera	*Ilex pernyi*	Japanese Holly; Helleri
Citrus spp.	Citrus species*	*Ilex cornuta*	American Holly
Coreopsis sp.	Coreopsis (Calliopsis): Early Sunrise, Moonbeam	*I. crenata*	Yaupon Holly; Schillings
Cornus florida	Flowering Dogwood, American Dogwood	*I. opaca*	Swordleaf Inula
		I. vomitoria	Sword-Leaved Iris; Jodlesong
C. stolonifera	Red Osier Dogwood	*Inula ensifolia*	Siberian Iris; Cabernet
Cortaderia selloana	Pampas Grass	*Iris ensata*	Iris species**
Cotoneaster apiculatus	Cranberry Cotoneaster	*Iris siberica*	
C. buxifolius	Cotoneaster	*Iris* spp.	

*** DO NOT use on food producing trees, vines, or plants. ** Not for use on container grown plants.**

(continued)

For technical assistance or more information about our products visit www. LebanonTurf.com

Manufactured by:
Lebanon Seaboard Corporation
1600 E. Cumberland St. • Lebanon, PA 17042
www.LebanonTurf.com

Scientific Name	Common Name
Jasminum nudiflorum	Winter Jasmine
Juniperus chinensis	Chinese Juniper; Nick's Compact, Parsonii
J. conferta	Shore Juniper; Blue Pacific
J. davurica	Parson's Juniper
J. horizontalis	Creeping Juniper
Juglans sp.	Walnut*
Justicia brandegeana	Shrimp Plant
Lagerstroemia indica	Crepe Myrtle
Lagerstroemia indica X Fauriei	Crepe Myrtle; Tuscarora
Lantana montevidensis	Weeping Lantana
Lavender sp.	Lavender; Munstead
Leontopodium alpinum	Edelweiss
Ligustrum sinense	Chinese Privet; Variegata
Ligustrum amurense	Amur Privet
L. japonicum	Japanese Privet
L. lucidum	Glossy Privet (Waxleaf)
Lilium sp.	Lily: Jazz
Liriope muscari	Big Blue Lily; Turf; Silver Mound; Evergreen Giant
Liriope muscari var. variegata	Liriope, Variegated
Liriope spicata	Liriope, Creeping
Lobelia Cardinalis	Cardinal Flower, Indian Pink
Lonicera japonica	Japanese Honeysuckle
L. tatarica	Tatarian Honeysuckle
Loropetalum chinense	Burgundy
Lythrum sp.	Loosestrife; Modern Pink
Magnolia spp.	Magnolia species**
Malephora luteola	Rocky Point Ice Plant
Malus sp.	Crabapple*
Miscanthus sinensis	Yaku Jima, Silberfeder
Nandina domestica	Heavenly Bamboo
Narcissus spp.	Narcissus species**
Nerium oleander	Oleander
Olea europaea	Olive*
Ophiopogon japonicus	Mondo Grass
Osmanthus heterphyllus	Osmanthus (False Holly): Gulf Tide
Osteospermum fruticosum	Trailing African Daisy
Oxydendrum arboreum	Sourwood
Paeonia suffruticosa	Tree Peony; Moutan Peony
Pennisetum setaceumi	Fountain Grass (Dwarf)
Perovskia atriplicifolia	Russian Sage
Persea americana	Avocado*
Photinia Fraseri	Frasier's Photinia (Redtip)
Physostegia virginiana	Dragonhead, False; Vivid
Picea spp.	Spruce species** (Colorado Blue, Norway, etc.)
Pieris japonica	Lily-of-the-Valley Bush
Pinus brutia	Calabrian Pine
P. canariensis	Canary Island Pine
P. elliottii	Slash Pine
P. halepensis	Aleppo Pine
P. nigra	Austrian Pine
P. Palustris	Longleaf Pine
P. radiata	Monterey Pine
P. thunbergiana	Japanese Black Pine
P. strobus	Eastern White Pine
P. sylvestris	Scotch Pine
P. taeda	Loblolly Pine
P. virginiana	Virginia Pine
Pistachio sp.	Pistachio*
Pittosporum rhombifolium	Queensland Pittosporum
P. tobira	Japanese Pittosporum
Podocarpus macrophyllus	Japanese Yew
Prunus laurocerasus	English Laurel
Prunus sp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
Pseudotsuga menziesii	Douglas Fir**
Pyracantha coccinea	Firethorn (Scarlet)
P. Fortuneana	Chinese Firethorn
P. Koidzumii	Firethorn; Formosa; Taiwan
Pyrus sp.	Bradford Pear spp.
Quercus shumardii	Oak, Shumard's Red
Quercus spp.	Oak species
Raphiolepis indica	Indian Hawthorn
Raphiolepsis umbellata	Yedda Hawthorn
Rhododendron (including Azalea)	'Coral Bells'
	'Delaware Valley White'
	'Flame Creeper'
	'Formosa'
	'Girard Crimson'
	'George L. Tabor'
	'Hino-crimson'
	'PJM'
	'Roseum Elegans'
	'Wakeiebisu'
	'White Gumbo'
Rosa Banksiae	Lady Bank's Rose
Rosmarinus officinalis	Rosemary*
Rudbeckia sp.	Black-Eyed Susan; Goldsturm
Santolina virens	Santolina
Saxifraga sp.	Saxifrage; Purple Dome
Scabiosa sp.	Pincushion Flower
Sedum album	White Stonecrop
Sedum cauticola	Cliff Stonecrop; Lidakense
Sedum dasyphyllum	Corsican Stonecrop
Sedum spurium	Stonecrop; Dragon's Blood
Spiraea Bumalda	Spirea: Anthony Waterer
Syzygium paniculatum	Australian Brush cherry
Taxus cuspidata	Japanese Yew
T. media	Yew
Teucrium sp.	Germander
Thalictrum dipterocarpum	Meadow Rue
Thuja occidentalis	American Arborvitae
Trachelospermum asiaticum	Star Jasmine
Tsuga canadensis	Canada Hemlock
Tulipa spp.	Tulip species
Veronica sp.	Veronica, Speedwell; Sunny Border
Viburnum japonicum	Japanese Viburnum
V. odoratissimum	Sweet Viburnum
V. plicatum	Japanese Snowball
V. rigidum	Canary Island Viburnum
V. suspensum	Sandankwa
V. tinus	Laurustinus
V. trilobum	Cranberry Bush
V. rhytidophyllum	Leatherleaf Virburnum
V. wrightii	Wright Virburnum
Vinca major	Greater Periwinkle, Vinca
Vinca minor	Common Periwinkle, Vinca
Vitis sp.	Grape*
Weigela florida	Old Fashioned Weigela
Yucca aloifolia	Spanish Bayonet, Yucca
Y. filamentosa	Adam's Needle, Yucca

*** DO NOT use on food producing trees, vines, or plants. ** Not for use on container grown plants.**

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.
PESTICIDE STORAGE: Store product in its original container in a cool, dry, locked place out of the reach of children and animals.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.
CONTAINER HANDLING: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

WARRANTY

NOTE: To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.