

DITHIOPYR

GROUP

3

HERBICIDE

CHLORANTRANILIPROLE

GROUP

28

INSECTICIDE

Pro

Scape®

TURF FERTILIZER

with 0.067% Acelepryn® Plus 0.167% Dimension®

24-0-8

GUARANTEED ANALYSIS

Total Nitrogen (N) .....24%

24.0% Urea Nitrogen\*

Soluble Potash (K<sub>2</sub>O).....8%

Derived From: Polymer Coated Urea, Urea, Muriate of Potash.

\*9.7% Slowly available nitrogen from polymer coated urea.

Chlorine (Cl) not more than.....7.0%

SUGGESTED SPREADER SETTINGS

Spreader	3.44 lbs./1,000 sq. ft.	5.17 lbs./1,000 sq. ft.	6.88 lbs./1,000 sq. ft.	Spreader	3.44 lbs./1,000 sq. ft.	5.17 lbs./1,000 sq. ft.	6.88 lbs./1,000 sq. ft.
HVN/Prizelawn	J	L	O	Lesco (letter/Numeric)	I/17	K/22	M/26
Andersons AccuPro	L	O	R	PennMulch HVO	L	O	Q
Earthway Rotary	14	18	22	ProScape SS	L	O	Q
Gandy	26	30	34	Spyker	4½	5	6
Lely	5 II	6½ II	7½ II	Vicon (all models)	24	30	32

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage. If your spreader is not listed, or if you need a spreader setting, please call customer service at 1-800-233-0628, Monday through Friday, 8 am. To 5 pm.

Information regarding the contents and levels of metals in this product is available on the Internet at:  
<http://www.regulatory-info-lebsea.com>

Intended for Use by Commercial Applicators Only

For systemic control of white grubs, other listed pests and listed weeds infesting landscape, recreational turfgrass (including golf courses).

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes.

Not for sale, sale into, distribution and/or use in Nassau, Suffolk, Kings, and Queens Counties of New York State.

ACTIVE INGREDIENTS:

BY WT %

Chlorantraniliprole\* .....0.067%

Dithiopyr\*\* .....0.167%

OTHER INGREDIENTS: .....99.766%

TOTAL: .....100.000%

\* Chlorantraniliprole belongs to the anthranilic diamide chemical class. CAS #500008-45-7

\*\*CAS # 97886-45-8

EPA Reg. No. 961-427

EPA Est. No. 961-PA-01: 961-IL-01

Superscript used is first letter of lot number.

Acelepryn® is a trademark of Syngenta.

Acelepryn® is a trademark or registered trademark of Syngenta used under license by Lebanon Seaboard Corporation.

Dimension® is a registered trademark of Corteva Agrisciences LLC and affiliates.

ProScape®, MESA® and EXPO® are registered trademarks of Lebanon Seaboard Corporation.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you **DO NOT** understand the label, find someone to explain it to you in detail.)

NOTICE: Read the entire label. Use only according to label directions. Before buying this product, read the Warranty Section elsewhere on this label. If terms are not acceptable, return the unopened container at once.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Gloves made of any waterproof material

After the product is loaded into properly calibrated application equipment, shirt, pants, socks and shoes are sufficient Personal Protective Equipment (PPE). Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

FIRST AID

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center (1-800-222-1222) or doctor or going for treatment. You may also contact (888)-208-1368 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and highly toxic to other aquatic organisms, including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. **DO NOT** apply directly to water. **DO NOT** apply when/where conditions favor runoff. Runoff from treated turf may adversely affect aquatic organisms in water adjacent to use sites. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

Surface Water Advisory:

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of chlorantraniliprole from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Groundwater Advisory:

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

NON-TARGET ORGANISM ADVISORY

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift. For further guidance and instructions on how to minimize spray drift, refer to the Spray Drift Management section of this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

DO NOT

apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

PRODUCT INFORMATION

This product is a combination product containing Acelepryn® insecticide (active ingredient: chlorantraniliprole), and Dimension® herbicide (active ingredient: dithiopyr) which may be applied for the control of white grubs, other listed insect pests, and listed weeds infesting landscape and recreational turfgrass, including golf courses.

This product may be used on established turfgrasses that are being grown for aesthetic or recreational purposes or climatic modification in or around home lawns, commercial lawns, industrial facilities, residential dwellings, business and office complexes, shopping complexes, multi-family residential complexes, institutional buildings, airports, cemeteries, parks, playgrounds, schools, day-care facilities, golf courses (tee box areas, roughs, fairways, collars etc.), athletic fields and other landscaped areas.

This product is in GROUP 28 of the EPA's Insecticide and Acaricide Groups Based on Target Site of Action (EPA PR Notice 2001-5) and this product may be used in rotational resistance management programs.

Consult your local Cooperative Extension Service specialist or pest control advisor for regionally specific information regarding application timing.

WEED RESISTANCE MANAGEMENT

For resistance management, Lebanon ProScape Turf Fertilizer with 0.067% Acelepryn Plus 0.167% Dimension is a Group 3 herbicide. Any weed population may contain or develop certain plants naturally resistant to Lebanon ProScape Turf Fertilizer with 0.067% Acelepryn Plus 0.167% Dimension and other Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies listed below should be followed.

To delay herbicide resistance, take one or more of the following steps:

Rotate the use of Lebanon ProScape Turf Fertilizer with 0.067% Acelepryn Plus 0.167% Dimension or other Group 3 herbicides within a growing season with different herbicide groups that control the same weeds.

Fields should be scouted prior to application to identify the weed species present and their growth stage to determine if the intended application will be effective.

Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by using an alternative herbicide from a different group or by a mechanical weed removal method. Prevent movement of resistant weed seeds to other areas by cleaning equipment thoroughly. If weed population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.

INSECT RESISTANCE MANGEMENT

For resistance management, Lebanon ProScape Turf Fertilizer with 0.067% Acelepryn Plus 0.167% Dimension is a Group 28 insecticide. Any insect population may contain certain or develop naturally resistant to Lebanon ProScape Turf Fertilizer with 0.067% Acelepryn Plus 0.167% Dimension and other Group 28 insecticides. The resistant individual may dominate the insect population if this group of insecticides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay insecticide resistance, take one or more of the following steps:

Rotate the use of Lebanon ProScape Turf Fertilizer with 0.067% Acelepryn Plus 0.167% Dimension or other Group 28 insecticides within a growing season or between growing seasons, with different groups that control the same pests. Monitor after application for unexpected target pest survival. If the level of survival suggests presences of resistance, consult your local university or local extension specialist.

Contact your sales representative, pest control advisors, or local extension specialist for additional pesticide resistance management and/or integrated pest management recommendations for specific types of plants, weeds and insects.

For further information or to report suspected resistance, contact Lebanon Seaboard Corporation representatives at 1-800-532-0090 or at [www.customerservice@lebsea.com](mailto:www.customerservice@lebsea.com).

USE RESTRICTIONS

- **DO NOT** formulate this product into other end-use products.
- **DO NOT** apply this product in commercial nurseries or greenhouses.
- **DO NOT** apply this product on golf course putting greens.
- **DO NOT** apply more than 299 pounds of product/A (equivalent to 0.20 lb. of chlorantraniliprole/A and 0.5 lb. of dithiopyr/A) per application.
- **DO NOT** apply more than 449 pounds of product/A (equivalent to 0.30 lb. of chlorantraniliprole/A and 0.75 lb. of dithiopyr/A) per year in broadcast applications to turfgrass.
- **DO NOT** use clippings from treated turf for mulching around trees or fruit trees.
- Apply only to lawns and turfgrasses that are well established. **DO NOT** apply to lawns or turfgrasses that are under stress.
- **DO NOT** apply by air.
- Keep people and pets off treated areas until dust has settled.
- This product must not be applied when the soil is saturated with water because adequate distribution of the active ingredients vertically in the soil profile cannot be achieved under this condition.
- **DO NOT** apply more than 1.5 lbs. a.i./A of dithiopyr per year.
- Maximum single application rate at 0.5 lb. a.i./A.

In Pennsylvania and New Hampshire: **DO NOT** exceed 0.9 lb. of total nitrogen per 1,000 sq. ft. per application.

The following restrictions are required to permit use of this product in the State of New York:

Golf Courses:

Tee Boxes: **DO NOT** apply this product within 25 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

Fairways: **DO NOT** apply this product within 50 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

All other use sites:

**DO NOT** apply this product within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

All use sites:

**DO NOT** exceed 299 lbs. of product/A per year (equivalent of 0.5 lb. dithiopyr A.I./A per year).

This product may only be used by commercial applicators.

**DO NOT** use in Nassau, Suffolk, Kings, Queens Counties.

TOLERANT TURFGRASS SPECIES

Apply this product to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

Cool Season Turfgrass	Warm Season Turfgrass
Bentgrass, creeping*** ( <i>Agrostis palustris</i> )	Bahiagrass ( <i>Paspalum notatum</i> )
Bluegrass, Kentucky ( <i>Poa pratensis</i> )	Bermudagrass ( <i>Cynodon dactylon</i> )
Fescue, fine**** ( <i>Festuca rubra</i> )	Buffalograss***** ( <i>Buchloe dactyloides</i> )
Fescue, tall ( <i>Festuca arundinacea</i> )	Carpetgrass ( <i>Axonopus affinis</i> )
Ryegrass, perennial ( <i>Lolium perenne</i> )	Centipedegrass ( <i>Eremochloa ophiuroides</i> )
	Kikuyugrass ( <i>Pennisetum clandestinum</i> )
	St. Augustinegrass ( <i>Stenotaphrum secundatum</i> )
	Zoysiagrass ( <i>Zoysia japonica</i> )

\*\*\* Use of this product on certain varieties of creeping turfgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested.

**DO NOT** apply this product to Colonial bentgrass (*Agrostis tenuis*) varieties.

\*\*\*\* Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', Tamara', 'Tatjana', 'Waldorf', and 'Waldina'. Not all varieties of fine fescue have been tested.

\*\*\*\*\* **DO NOT** use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

**NOTE:** It is not possible to evaluate all turfgrass varieties for tolerance to dithiopyr. It is suggested that the user evaluate the response of turfgrass varieties to this product on a limited area, at the specified use rates, prior to initiating large-scale use.

Manufactured by:

Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042

800-233-0628 • (717) 273-1685

Made in the U.S.A.

[www.LebanonTurf.com](http://www.LebanonTurf.com)

(continued on reverse)

REV: EH/C-25

For technical assistance or more information about our products visit: [www.LebanonTurf.com](http://www.LebanonTurf.com)

Net Weight: 50 LBS. (22.7 kg)

**APPLICATION EQUIPMENT**

Apply this product with drop-type or rotary-type equipment. The application equipment must be properly calibrated according to the manufacturer's directions to ensure uniform coverage over the treated area. During application, check the equipment periodically to confirm that it is working properly and that the granules are being uniformly distributed. A more uniform application can be made by spreading one-half of the specified amount of this product over the turfgrass area and then spreading the remaining one-half of this product at a right angle to the previous application direction.

**APPLICATION TIMING – INSECT CONTROL**

**White Grubs:** Apply this product from early April to early September for preventative and early curative control of all of the major white grub species infesting turfgrass. The need for an application may be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. The higher rate listed for white grub control in Table 1 may be required in late August or early September when less sensitive mid-instar grubs are present at the time of application. Irrigate turf immediately after application or allow rainfall to move the product into the soil.

**Billbugs:** Apply this product when overwintered adult billbugs are first observed. This will usually occur in late April or early May in regions with cool-season turfgrasses. An application of this product at this time will also provide white grub control. The higher rates listed in Table 1 may be required for applications made after early May or when billbug species other than bluegrass billbug are present.

**European Crane Fly:** Apply this product between September and November to control European crane fly larvae in turfgrass. The higher rate listed in Table 1 may be required to achieve control when applications are made in November.

**Turf Caterpillars:** This product will provide curative caterpillar control in turfgrass. If the area being treated is maintained at a mowing height of greater than one inch, the higher rate listed in Table 1 may be required during periods of high pest pressure.

**APPLICATION TIMING – WEED CONTROL**

**Spring Application**

**Control of Crabgrass:** This product provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only and is effective up to 4 weeks after germination of crabgrass. This normally corresponds to the time when crabgrass seedlings are at the 3-leaf stage of growth and first become visible in the established turf. This product will not control crabgrass at later growth stages when crabgrass plants begin to tiller (spread). Properly timed applications will control existing crabgrass and prevent further emergence of crabgrass seedlings. **DO NOT** apply this product later than 4 weeks after crabgrass germination.

**Control or Suppression of Other Grass and Broadleaf Weeds:** For control or suppression of other grass and broadleaf weeds, this product must be applied as a **preventative** treatment prior to their germination and emergence. Only crabgrass will be controlled with applications made up to 4 weeks after weed germination.

**Fall Application**

This product can also be applied from late summer through early fall (late August through November). When applied in the fall, a sequential or second application of this product must be made the following spring to provide season-long control of crabgrass. If this product is applied in the fall, **DO NOT** reseed, overseed or sprig turfgrasses within 12 weeks after application.

**Tips for Improved Weed Control**

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
- The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.

**Weeds Controlled:** This product will provide control of crabgrass and other grass and broadleaf weeds listed below when applied prior to germination in established lawns and ornamental grasses.

Grasses	Broadleaf Weeds
Barley	Carpetweed
Barryardgrass	Henbit
Bluegrass, Annual	Knotweed, Prostrate
Brome	Marestail
Crabgrass, Large	Medic, Black
Crabgrass, Smooth	Mustard
Crabgrass, Southern	Oxalis, Buttercup
Dallisgrass	Pineappleweed
Goosegrass	Pigweed, Redroot
Foxtail, Green	Purslane, Common
Foxtail, Yellow	Rocket, London
Oats, Wild	Shepherd's Purse
Ryegrass (Annual & Perennial)	Spurge, Garden
Sandbur	Spurge, Prostrate
Smutgrass	Spurge, Spotted
	Woodsorrel, Creeping
	Woodsorrel, Yellow

**APPLICATION RATES FOR LAWNS, GOLF COURSES AND OTHER RECREATIONAL TURFGRASS AREAS**

Table 1 provides the application rates that will control the listed insect pests and will also control the listed grass and broadleaf weeds.

**Restrictions:** **DO NOT** apply more than 299 pounds of this product/acre per application and no more than 299 pounds (equivalent to 0.20 lb. of chlorantraniliprole and 0.75 lb. of dithiopyr) of product/acre per year in broadcast applications to turfgrass.

TABLE 1: TURFGRASS APPLICATION RATES				
Target Pest (Including Listed Weeds)	Product per Acre	Product per 1,000 Sq. Ft.	Lb A.I./A Acelepryn	Lb A.I./A Dimension
White Grubs (including <i>Aphodius</i> spp., Asiatic garden beetle, black turfgrass ataenius, European chafer, green June beetle, Japanese beetle, May/June beetles ( <i>Phyllophaga</i> spp.), northern masked chafer, oriental beetle and southern masked chafer)	150 to 299 pounds	3.44 to 6.88 pounds	0.1 to 0.2	0.25 to 0.50
Billbugs				
European Crane Fly				
Turf caterpillars (including armyworms, cutworms and sod webworms)				
Annual Bluegrass Weevil Spittlebugs	225 to 299 pounds	5.17 to 6.88 pounds	0.15 to 0.20	0.38 to 0.50
Chinch Bugs (suppression)	150 to 225 pounds	3.44 to 5.17 pounds	0.1 to 0.15	0.25 to 0.38

**NOTE: For 3-4 months weed control** use 150 pounds/acre of this product.

**For early postemergence control of crabgrass,** use 150 to 299 pounds of this product/acre. Use the higher rate in areas of high weed pressure.

**In the State of New York: DO NOT exceed 0.5 pound of dithiopyr/A (299 pounds product/A) per year.**

**CLEANING APPLICATION EQUIPMENT**

Thoroughly clean all application equipment after use to reduce the risk of forming hardened deposits that might become difficult to remove. Thoroughly rinse application equipment with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. **DO NOT** clean equipment near wells, or other water sources. Dispose of waste rinse water in accordance with local regulations.

**STORAGE AND DISPOSAL**

**DO NOT** contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store product in its original container in a cool, dry, place out of the reach of children and animals.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING:** Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

**WARRANTY**

**NOTE:** To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.