

Pro

LEBANON Scape®



FERTILIZER 20-0-5 MESA[®]

GUARANTEED ANALYSIS

Total Nitrogen (N)	20%
2.0% Ammoniacal Nitrogen	
1.6% Water Insoluble Nitrogen*	
13.5% Urea Nitrogen	
2.9% Other Water Soluble Nitrogen*	
Soluble Potash (K₂O)	5%
Sulfur (S)	2.3%
2.3% Combined Sulfur (S)	
Iron (Fe)	1.0%
0.1% Water Soluble Iron (Fe)	

Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash, Ferric Oxide, Ferrous Sulfate.

*4.5% Slowly available Nitrogen from Methylene Ureas.

Chlorine (Cl) not more than

8.0%
U.S.A

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)	20%
2.0% Ammoniacal Nitrogen	
1.6% Water Insoluble Nitrogen*	
13.5% Urea Nitrogen	
2.9% Other Water Soluble Nitrogen*	
Soluble Potash (K₂O)	5%
Sulfur (S)	2.3%
2.3% Combined Sulfur (S)	
Iron (Fe) (actual)	1.0%
0.1% Water Soluble Iron (Fe)	

Chlorine (Cl) (maximum)

Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash, Ferric Oxide, Ferrous Sulfate.

*4.5% Slowly available Nitrogen from Methylene Ureas.

CANADA

NOTICE: This product contains the secondary nutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash off with water.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Application Rates:
To feed at the rate of 0.9lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 4.5 lbs per 1,000 sq. ft. or 196 lbs. per acre. To feed at the rate of 0.75 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 3.75 lbs per 1,000 sq. ft. or 164 lbs. per acre.

SUGGESTED SPREADER SETTINGS		
Spreader	0.9# N /1,000 sq. ft.	0.75# N /1,000 sq. ft.
Lebanon Turf	5	4½
Andersons AccuPro	N	L
Earthway Rotary	17	15
Gandy	29	27
Lely	6 II	5½ II
Lesco (letter/numeric)	K/22	J/21
PennMulch HVO	M	L
ProScape SS	M	L
Spyker	5	4½
Vicon	27	25

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

MESA®, PennMulch® and ProScape® are registered trademarks of Lebanon Seaboard Corporation.

Warning



Hazard Statements
Causes skin irritation.
Causes eye irritation.
May be harmful if inhaled.
Suspected of causing cancer.

Precautionary Statements for handling:

Do not handle until all safety precautions have been read and understood.
Avoid breathing dust/fume/gas/mist/vapours/spray.
Wash hands and exposed skin thoroughly after handling.
Wear protective gloves/protective clothing/eye protection/face protection.
Use personal protective equipment as required.
If in eyes, rinse with water and seek medical attention if symptoms persist.
Take off contaminated clothing and wash before reuse.
IF ON SKIN: Wash with plenty of soap and water.
IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell.
If skin irritation occurs: Get medical advice/attention.
If eye irritation persists: Get medical advice/attention.
Store locked up.
Dispose of contents/container properly as required by law for waste pesticides.
If exposed or concerned, seek medical advice.
Keep out of reach of children
Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1.2 to 3.2 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:
To apply at the rate of 0.43 kg N/100 sq. meters. Equivalent to 2.1 kg of product per 100 sq. meters.
To apply at the rate of 0.36 kg N/100 sq. meters. Equivalent to 1.7 kg of product per 100 sq. meters.

SUGGESTED SPREADER SETTINGS		
Spreader	0.43 kg N /100 m ² .	0.36 kg N /100 m ² .
Lebanon Turf	5	4½
Andersons AccuPro	N	L
Earthway Rotary	17	15
Gandy	29	27
Lely	6 II	5½ II
Lesco (letter/numeric)	K/22	J/21
PennMulch HVO	M	L
ProScape SS	M	L
Spyker	5	4½
Vicon	27	25

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



0 88685 54316 6
F699 **22-54316** G/17

Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.regulatory-info-lebsea.com>

LebanonTurf

For technical assistance or more information about our products visit www.LebanonTurf.com

Manufactured by:

Lebanon Seaboard Corporation
1600 E. Cumberland St. • Lebanon, PA 17042

Made in the U.S.A.

www.LebanonTurf.com

800-233-0628 • (717-273-1685)

NET WEIGHT **50** LBS. (22.7 KG)



FERTILIZER 20-0-5

MESA[®]

P A T E N T E D

GUARANTEED ANALYSIS

Total Nitrogen (N)	20%
2.0% Ammoniacal Nitrogen	
1.6% Water Insoluble Nitrogen*	
13.5% Urea Nitrogen	
2.9% Other Water Soluble Nitrogen*	
Soluble Potash (K₂O)	5%
Sulfur (S)	2.3%
2.3% Combined Sulfur (S)	
Iron (Fe)	1.0%
0.1% Water Soluble Iron (Fe)	

Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash, Ferric Oxide, Ferrous Sulfate.

*4.5% Slowly available Nitrogen from Methylene Ureas.

Chlorine (Cl) not more than **8.0%**
U.S.A

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)	20%
2.0% Ammoniacal Nitrogen	
1.6% Water Insoluble Nitrogen*	
13.5% Urea Nitrogen	
2.9% Other Water Soluble Nitrogen*	
Soluble Potash (K₂O)	5%
Sulfur (S)	2.3%
2.3% Combined Sulfur (S)	
Iron (Fe) (actual)	1.0%
0.1% Water Soluble Iron (Fe)	

Chlorine (Cl) (maximum) **8.0%**

Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash, Ferric Oxide, Ferrous Sulfate.

*4.5% Slowly available Nitrogen from Methylene Ureas.

CANADA

NOTICE: This product contains the secondary nutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash off with water.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Application Rates:
To feed at the rate of 0.9lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 4.5 lbs per 1,000 sq. ft. or 196 lbs. per acre. To feed at the rate of 0.75 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 3.75 lbs per 1,000 sq. ft. or 164 lbs. per acre.

SUGGESTED SPREADER SETTINGS		
Spreader	0.9# N /1,000 sq. ft.	0.75# N /1,000 sq. ft.
Lebanon Turf	5	4½
Andersons AccuPro	N	L
Earthway Rotary	17	15
Gandy	29	27
Lely	6 II	5½ II
Lesco (letter/numeric)	K/22	J/21
PennMulch HVO	M	L
ProScape SS	M	L
Spyker	5	4½
Vicon	27	25

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Warning



Hazard Statements

Causes skin irritation.
Causes eye irritation.
May be harmful if inhaled.
Suspected of causing cancer.

Precautionary Statements for handling:


Do not handle until all safety precautions have been read and understood.
Avoid breathing dust/fume/gas/mist/vapours/spray.
Wash hands and exposed skin thoroughly after handling.
Wear protective gloves/protective clothing/eye protection/face protection.
Use personal protective equipment as required.
If in eyes, rinse with water and seek medical attention if symptoms persist.
Take off contaminated clothing and wash before reuse.
IF ON SKIN: Wash with plenty of soap and water.
IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell.
If skin irritation occurs: Get medical advice/ attention.
If eye irritation persists: Get medical advice/attention.
Store locked up.
Dispose of contents/container properly as required by law for waste pesticides.
If exposed or concerned, seek medical advice.
Keep out of reach of children
Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1.2 to 3.2 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:
To apply at the rate of 0.43 kg N/100 sq. meters. Equivalent to 2.1 kg of product per 100 sq. meters.
To apply at the rate of 0.36 kg N/100 sq. meters. Equivalent to 1.7 kg of product per 100 sq. meters.

SUGGESTED SPREADER SETTINGS		
Spreader	0.43 kg N /100 m ² .	0.36 kg N /100 m ² .
Lebanon Turf	5	4½
Andersons AccuPro	N	L
Earthway Rotary	17	15
Gandy	29	27
Lely	6 II	5½ II
Lesco (letter/numeric)	K/22	J/21
PennMulch HVO	M	L
ProScape SS	M	L
Spyker	5	4½
Vicon	27	25

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



F699

22-54316

G/17

Information regarding the contents and levels of metals in this product is available on the Internet at:
<http://www.regulatory-info-lebsea.com>



For technical assistance or more information about our products visit www.LebanonTurf.com

Manufactured by:

Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042

Made in the U.S.A.

www.LebanonTurf.com

800-233-0628 • (717-273-1685)

NET WEIGHT 50 LBS. (22.7 KG)