



woodace[®]

18-5-10

With

METH-EX[®] 40

longterm fertilizer

PROFESSIONAL LANDSCAPE
NUTRITION

+ Minor Elements

with

expo[®]
& EXTENDED
POTASSIUM

METH-EX[®] 40

Lebanon Seaboard Corporation

1600 East Cumberland Street • Lebanon, PA 17042

800-233-0628 • (717) 273-1685

www.LebanonTurf.com

NET WT. 50 LBS. (22.7 KG)

Nitrogen	18%
Soluble Nitrogen*	
.....	
Soluble Nitrogen*	
(O ₅)	5%
.....	
.....	10%
.....	0.5%
Soluble Magnesium (Mg)	
.....	3.2%
Sulfur (S)	
.....	1.6%
Soluble Iron (Fe)	
.....	0.3%
Soluble Manganese (Mn)	

Chlorine (Cl) not more than 2.0%

NOTICE: This product contains the secondary nutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash off with water.



F699 **22-56351** F/19

Woodace Longterm 18-5-10 now contains both Expo® and METHEX®40 slow release methylene urea nitrogen. Expo provides controlled released Potash from occluded Sulfate of Potash. This slow release product is ideal for season long, single application feeding of landscape plants, trees and ground covers. Woodace Longterm can reduce labor costs by eliminating the need for multiple applications in one season. The extensive micro package includes Magnesium, Sulfur, Iron, and Manganese. Nutrient sources are selected to reduce the salt index allowing higher rates without burning.

Woodace Longterm is designed for top-dress application. For individual plants apply Woodace Longterm evenly within the dripline or known root area. The feeder roots of large shrubs and trees may extend well beyond the dripline. Adjust the recommended rate and treated area accordingly. For densely planted areas or ground covers broadcast Woodace Longterm evenly over the entire area at a rate of 1.1 - 3.3 lbs. per 100 sq. ft., (2 - 6 lbs. N/1000 sq. ft.). Adjust the height and trajectory of the spreader where possible to minimize reflecting granules off of dense foliage. Do not broadcast fertilizer onto wet foliage. Fertilizer should be applied when plants are actively growing. Woodace Longterm can be lightly raked into the mulch or soil surface. However, do not attempt to incorporate fertilizers into the root area of established plants. Avoid piling fertilizer against the stems of plants. Never fertilize when plants are suffering from stress related to transplant, moisture, heat, insect, disease or abiotic injury. Generally spring and fall provide the optimum time to apply supplemental nutrition.

Woodace Longterm should be a part of an integrated landscape maintenance program. In addition to fertilization, pest monitoring, weed control, pruning, and water management are essential. In most landscape situations nutrients are not naturally replenished at a sufficient rate to guarantee optimum health and appearance. The objective of most landscape fertilization programs is to improve the health and appearance of plantings by providing proper fertility. Other objectives for which Woodace Longterm may be used include: supplementing nutrients known to be deficient, encouraging foliar or root growth, increasing bloom or fruiting, or to aid in recovery from injury or disease. These label directions are intended to serve as a general guideline for the use of Woodace Longterm. Other resources such as soil testing, local extension recommendations, and specific plant knowledge should be incorporated whenever possible.

HAZARD STATEMENTS

May cause respiratory irritation.



Use in accordance with recommendations of a qualified individual or institution

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

Do not apply to frozen ground

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

broadcast (for heavily planted beds)

N rate	lbs./100 sq. ft.	lbs./1000 sq. ft.
2 lbs./1000 sq. ft.	1.1 lbs.	11 lbs.
4 lbs./1000 sq. ft.	2.2 lbs.	22 lbs.
6 lbs./1000 sq. ft.	3.3 lbs.	33 lbs.

size	rate	measure
8-12"	2 oz.	¼ cup
13-18"	4 oz.	½ cup
19-24"	6 oz.	¾ cup
25-30"	8 oz.	1 cup
31-36"	12 oz.	1½ cups
37-42"	16 oz.	2 cups
> 42"	1½ cups per foot of branch spread	



Shrub size refers to branch spread for shrubs where height and spread are approximately the same. For fastigate or decumbent forms reduce the amount by approximately 1/2.

The user assumes all liability for applications of this product. For uses that are not specifically listed on the product label, it is recommended that the user perform a test on a limited basis before commencing large scale applications.

expo®, METH-EX® Slow Release Nitrogen and Woodace® are registered trademarks of Lebanon Seaboard Corporation.

www.LebanonTurf.com